


PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07C 239/14, C07D 211/20, 241/04, A61K 31/16	A1	(11) International Publication Number: WO 99/42436 (43) International Publication Date: 26 August 1999 (26.08.99)
(21) International Application Number: PCT/US98/17633 (22) International Filing Date: 26 August 1998 (26.08.98) (30) Priority Data: 09/026,372 19 February 1998 (19.02.98) US (71) Applicant: AMERICAN CYANAMID COMPANY [US/US]; Five Giralda Farms, Madison, NJ 07940-0874 (US). (72) Inventors: VENKATESAN, Aranapakam, Mudumbai; 97-07 63rd Road #9K, Rego Park, NY 11374 (US). GROSU, George, Theodore; 117 Prospect Place, Pearl River, NY 10965 (US). DAVIS, Jamie, Marie; 20 High Avenue, Nyack, NY 10960 (US). BAKER, Jannie, Lea; 127 Rockinchair Road, White Plains, NY 10607 (US). LEVIN, Jeremy, Ian; 190 Treetop Circle, Nanuet, NY 10954 (US). (74) Agents: ALICE, Ronald, W.; American Home Products Cor- poration, Patent Law Dept. - 2B, One Campus Drive, Par- sippany, NJ 07054 (US) et al.		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: N-HYDROXY-2-(ALKYL, ARYL, OR HETEROARYL SULFANYL, SULFINYL OR SUL- FONYL)-3-SUBSTITUTED-ALKYL, ARYL OR HETEROARYLAMIDES AS MATRIX METALLOPROTEINASE INHIBITORS <div style="text-align: center;">  (1) </div>		
(57) Abstract Matrix metalloproteinases (MMPs) are a group of enzymes that have been implicated in the pathological destruction of connective tissue and basement membranes. These zinc containing endopeptidases consist of several subsets of enzymes including collagenases, stromelysins and gelatinases. TNF- α converting enzyme (TACE), a pro-inflammatory cytokine, catalyzes the formation of TNF- α from membrane bound TNF- α precursor protein. It is expected that small molecule inhibitors of MMPs and TACE therefore have the potential for treating a variety of disease states. The present invention provides low molecular weight, non-peptide inhibitors of matrix metalloproteinases (MMPs) and TNF- α converting enzyme (TACE) for the treatment of arthritis, tumor metastasis, tissue ulceration, abnormal wound healing, periodontal disease, bone disease, diabetes (insulin resistance) and HIV infection having formula (I), wherein R ² and R ³ form a heterocyclic ring and A is S, S(O), or S(O) ₂ and R ¹ and R ⁴ are defined herein.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

WO 99/42436

PCT/US98/17633

- 1 -

**N-HYDROXY-2-(ALKYL, ARYL, OR HETEROARYL SULFANYL,
SULFINYL OR SULFONYL)-3-SUBSTITUTED ALKYL, ARYL OR
HETEROARYLAMIDES AS MATRIX METALLOPROTEINASE
INHIBITORS**

5

BACKGROUND OF THE INVENTION

Matrix metalloproteinases (MMPs) are a group of enzymes that have been implicated in the pathological destruction of connective tissue and basement
10 membranes. These zinc containing endopeptidases consist of several subsets of enzymes including collagenases, stromelysins and gelatinases. Of these classes, the gelatinases have been shown to be the MMPs most intimately involved with the growth and spread of tumors. It is known that the level of expression of gelatinase is elevated in malignancies, and that gelatinase can degrade the basement membrane which leads to
15 tumor metastasis. Angiogenesis, required for the growth of solid tumors, has also recently been shown to have a gelatinase component to its pathology. Furthermore, there is evidence to suggest that gelatinase is involved in plaque rupture associated with atherosclerosis. Other conditions mediated by MMPs are restenosis, MMP-mediated osteopenias, inflammatory diseases of the central nervous system, skin aging, tumor
20 growth, osteoarthritis, rheumatoid arthritis, septic arthritis, corneal ulceration, abnormal wound healing, bone disease, proteinuria, aneurysmal aortic disease, degenerative cartilage loss following traumatic joint injury, demyelinating diseases of the nervous system, cirrhosis of the liver, glomerular disease of the kidney, premature rupture of fetal membranes, inflammatory bowel disease, periodontal disease, age
25 related macular degeneration, diabetic retinopathy, proliferative vitreoretinopathy, retinopathy of prematurity, ocular inflammation, keratoconus, Sjogren's syndrome, myopia, ocular tumors, ocular angiogenesis/neo-vascularization and corneal graft rejection. For recent reviews, see: (1) Recent Advances in Matrix Metalloproteinase Inhibitor Research, R. P. Beckett, A. H. Davidson, A. H. Drummond, P. Huxley and
30 M. Whittaker, Research Focus, Vol. 1, 16-26, (1996), (2) Curr. Opin. Ther. Patents (1994) 4(1): 7-16, (3) Curr. Medicinal Chem. (1995) 2: 743-762, (4) Exp. Opin. Ther. Patents (1995) 5(2): 1087-110, (5) Exp. Opin. Ther. Patents (1995) 5(12): 1287-1196.

TNF- α converting enzyme (TACE) catalyzes the formation of TNF- α from
35 membrane bound TNF- α precursor protein. TNF- α is a pro-inflammatory cytokine that is now thought to have a role in rheumatoid arthritis, septic shock, graft rejection, cachexia, anorexia, inflammation, congestive heart failure, inflammatory disease of the

WO 99/42436


PCT/US98/17633

- 2 -

central nervous system, inflammatory bowel disease, insulin resistance and HIV infection in addition to its well documented antitumor properties. For example, research with anti- TNF- α antibodies and transgenic animals has demonstrated that blocking the formation of TNF- α inhibits the progression of arthritis. This observation has recently
5 been extended to humans as well.

It is expected that small molecule inhibitors of MMPs and TACE therefore have the potential for treating a variety of disease states. While a variety of MMP and TACE inhibitors have been identified and disclosed in the literature, the vast majority of these molecules are peptidic and peptide-like compounds that one would expect to have
10 bioavailability and pharmacokinetic problems common to such compounds that would limit their clinical effectiveness. Low molecular weight, potent, long acting, orally bioavailable inhibitors of MMPs and/or TACE are therefore highly desirable for the potential chronic treatment of the above mentioned disease states.

15 Recently, two references have appeared (U.S. 5,455,258 and European Patent Appl. 606,046) that disclose arylsulfonamido-substituted hydroxyamic acids. These documents cover compounds exemplified by CGS 27023A. These are the only non-peptide matrix metalloproteinase inhibitors disclosed to date.




CGS 27023A

20 Salah et al., Liebigs Ann. Chem. 195, (1973) discloses some aryl substituted thio and aryl substituted sulfonyl acetohydroxamic acid derivatives of general formula 1. These compounds were prepared to study the Mannich reaction. Subsequently, they
25 were tested for their fungicidal activity.

WO 99/42436

PCT/US98/17633

- 3 -



Some sulfone carboxylic acids are disclosed in U.S. patent 4,933,367. Those compounds were shown to exhibit hypoglycemic activity.

5

SUMMARY OF THE INVENTION:

The present invention relates to novel, low molecular weight, non-peptide inhibitors of matrix metalloproteinases (MMPs) and TNF- α converting enzyme (TACE) for the treatment of arthritis, tumor metastasis, tissue ulceration, abnormal wound healing, periodontal disease, bone disease, diabetes (insulin resistance) and HIV infection.

In accordance with this invention there is provided a group of compounds of general formula I

15



wherein:

- R¹ is alkyl of 1 to 18 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;
- alkenyl of 3 to 18 carbon atoms having 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;
- alkynyl of 3 to 18 carbon atoms having 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;
- aryl of 6 to 10 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;
- cycloalkyl of 3 to 8 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

WO 99/42436

PCT/US98/17633

- 4 -

- saturated or unsaturated 5 to 10 membered mono or bicyclic heterocycle
containing one heteroatom selected from O, S or NR⁷, optionally
substituted with one or two groups selected independently from R⁵;
or heteroaryl-(CH₂)₀₋₆ wherein the heteroaryl group is 5 to 6 membered with
one or two heteroatoms selected independently from O, S, and N and
may be optionally substituted with one or two groups selected
independently from R⁵;
- A is -S-, -SO- or SO₂-;
- R² and R³, taken with the carbon atom to which they are attached, form a 5 to 7
membered heterocyclic ring containing O, S or N-R⁷ optionally having one or
two double bonds;
- R⁴ is hydrogen,
alkyl of 1 to 6 carbon atoms, optionally substituted with one or two groups
selected independently from R⁵;
- alkenyl of 3 to 18 carbon atoms having 1 to 3 double bonds, optionally
substituted with one or two groups selected independently from R⁵;
- alkynyl of 3 to 18 carbon atoms having 1 to 3 triple bonds, optionally
substituted with one or two groups selected independently from R⁵;
- phenyl or naphthyl optionally substituted with one or two groups selected
independently from R⁵;
- C₃ to C₈ cycloalkyl or bicycloalkyl optionally substituted with one or two
groups selected independently from R⁵;
- saturated or unsaturated 5 to 10 membered mono or bicyclic heterocycle
containing one heteroatom selected from O, S or NR⁷, optionally
substituted with one or two groups selected independently from R⁵;
- R⁵ is H, C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁ to C₁₂ alkyl, C₂ to C₁₂ alkenyl,
C₂-C₁₂ alkynyl, F, Cl, Br, I, CN, CHO, C₁-C₆ alkoxy, aryloxy,
heteroaryloxy, C₃-C₆ alkenyloxy, C₃-C₆ alkynyloxy, C₁-C₆ alkoxyaryl,
C₁-C₆ alkoxyheteroaryl, C₁-C₆ alkylamino-C₁-C₆ alkoxy, C₁-C₂ alkylene
dioxy, aryloxy-C₁-C₆ alkyl amine, C₁-C₁₂ perfluoro alkyl, S(O)_n-C₁-C₆ alkyl,
S(O)_n-aryl where n is 0, 1 or 2; OCOO C₁-C₆ alkyl, OCOOaryl, OCONR⁶,
COOH, COO C₁-C₆ alkyl, COOaryl, CONR⁶R⁶, CONHOH, NR⁶R⁶,
SO₂NR⁶R⁶, NR⁶SO₂aryl, -NR⁶CONR⁶R⁶, NHSO₂CF₃.

WO 99/42436

PCT/US98/17633

- 5 -

- SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl, CONHSO₂aryl, SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl, CONHSO₂aryl, NH₂, OH, aryl, heteroaryl, C₃ to C₈ cycloalkyl; or saturated or unsaturated 5 to 10 membered mono or bicyclic heterocycle containing one heteroatom selected from O, S or NR⁷, wherein C₁-C₆ alkyl is straight or branched, heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;
- 10 R⁶ is H, C₁ to C₁₈ alkyl optionally substituted with OH; C₃ to C₆ alkenyl, C₃ to C₆ alkynyl, C₁ to C₆ perfluoro alkyl, S(O)_n-C₁-C₆ alkyl S(O)_n aryl where n is 0, 1 or 2, or COheteroaryl, wherein heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;
- 15 and R⁷ is C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁-C₁₂ perfluoro alkyl, S(O)_n-C₁-C₆-alkyl, S(O)_n-aryl where n is 0, 1 or 2; COO-C₁-C₆-alkyl, COOaryl, CONHR⁶, CONR⁶R⁶, CONHOH, SO₂NR⁶R⁶, SO₂CF₃, SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO-C₁-C₆-alkyl, CONHSO₂aryl, aryl, or heteroaryl, where aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected independently from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy; and heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or N-C₁-C₆ alkyl;
- 20 alkyl of 1 to 18 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;
- alkenyl of 3 to 18 carbon atoms having from 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;
- 30 alkynyl of 3 to 18 carbon atoms having from 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;
- aryllalkyl of 7 to 16 carbon atoms, wherein aryl is optionally substituted with one or two groups selected independently from R⁵;


WO 99/42436

PCT/US98/17633

- 6 -

heteroarylalkyl wherein alkyl is from 1 to 6 carbon atoms and heteroaryl contains 1 or 2 heteroatoms selected from O, S or N and is optionally substituted with one or two groups selected independently from R⁵;
 biphenylalkyl of 13 to 18 carbon atoms, wherein biphenyl is optionally substituted with one or two groups selected independently from R⁵;
 arylalkenyl of 8 to 16 carbon atoms, wherein aryl is optionally substituted with one or two groups selected independently from R⁵;
 cycloalkylalkyl or bicycloalkylalkyl of 4 to 12 carbon atoms, wherein the cycloalkyl or bicycloalkyl group is optionally substituted with one or two groups selected independently from R⁵;
 saturated or unsaturated mono or bicyclic heterocycle containing one heteroatom selected from O, S or N-C₁-C₆ alkyl, optionally substituted with one or two groups selected independently from R⁵; or
 R⁸R⁹N-C₁-C₆-alkoxyaryl-C₁-C₆-alkyl where R⁸ and R⁹ are independently selected from C₁-C₆ alkyl or R⁸ and R⁹ together with the interposed nitrogen forms a 5-7 membered saturated heterocyclic ring optionally containing an oxygen atom, wherein the aryl group is phenyl or naphthyl;
 and the pharmaceutically acceptable salts thereof.

A more preferred aspect of the present invention is the group of compounds of general formula (Ia):



wherein:

R¹ is alkyl of 1 to 18 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;
 alkenyl of 3 to 18 carbon atoms having 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;
 alkynyl of 3 to 18 carbon atoms having 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;
 aryl of 6 to 10 carbon atoms, optionally substituted with one to two groups selected independently from R⁵;

WO 99/42436

PCT/US98/17633

- 7 -

- cycloalkyl of 3 to 8 carbon atoms, optionally substituted with one to two groups selected independently from R⁵;
 saturated or unsaturated mono or bicyclic heterocycle of from 5 to 10 members containing one heteroatom selected from O, S or NR⁷, optionally substituted with one to two groups selected independently from R⁵;
 5 or heteroaryl-(CH₂)₀₋₆ wherein the heteroaryl group is 5 to 6 membered with one or two heteroatoms selected independently from O, S, and N and may be optionally substituted with one or two groups selected independently from R⁵;
- 10 A is -S-, -SO- or SO₂-;
 R² and R³, taken with the carbon atom to which they are attached, form a 5 to 7 membered heterocyclic ring containing O, S or N-R⁷ optionally having one or two double bonds;
 R⁴ is hydrogen,
- 15 alkyl of 1 to 6 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;
 alkenyl of 3 to 18 carbon atoms having 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;
 alkynyl of 3 to 18 carbon atoms having 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;
 20 phenyl or naphthyl optionally substituted with one or two groups selected independently from R⁵;
 C₃ to C₈ cycloalkyl or bicycloalkyl optionally substituted with one or two groups selected independently from R⁵;
- 25 R⁵ is H, F, Cl, Br, I, CN, CHO, C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁ to C₁₂ alkyl, C₂ to C₁₂ alkenyl, C₂-C₁₂ alkynyl, C₁-C₆ alkoxy, aryloxy, heteroaryloxy, C₃-C₆ alkenyloxy, C₃-C₆ alkynyloxy, C₁-C₆ alkoxyaryl, C₁-C₆ alkoxyheteroaryl, C₁-C₆-alkylamino-C₁-C₆ alkoxy, C₁-C₂-alkylene dioxy, aryloxy-C₁-C₆ alkylamine, C₁-C₁₂ perfluoro alkyl, S(O)_n-C₁-C₆ alkyl, S(O)_n-aryl where n is 0, 1 or 2; OCOO-C₁-C₆ alkyl, OCOOaryl, OCONR⁶, COOH, COO-C₁-C₆ alkyl, COOaryl, CONR⁶R⁶, CONHOH, NR⁶R⁶, SO₂NR⁶R⁶, NR⁶SO₂aryl, NR⁶CONR⁶R⁶, NHSO₂CF₃, SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl, CONHSO₂aryl, SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl,
- 30

WO 99/42436

PCT/US98/17633

- 8 -

CONHSO₂aryl, NH₂, OH, aryl, heteroaryl, C₃ to C₈ cycloalkyl; or saturated or unsaturated 5 to 10 membered mono or bicyclic heterocycle containing one heteroatom selected from O, S or NR⁷;

5 wherein heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected independently from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;

10 R⁶ is H, C₁ to C₁₈ alkyl optionally substituted with OH; C₃ to C₆ alkenyl, C₃ to C₆ alkynyl, C₁ to C₆ perfluoro alkyl, S(O)_n alkyl or aryl where n is 0, 1, or 2; or COheteroaryl;

15 wherein heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;

20 and R⁷ is C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁-C₁₂ perfluoro alkyl, S(O)_n-alkyl, S(O)_n-aryl where n is 0, 1 or 2; COOalkyl, COOaryl, CONHR⁶, CONR⁶R⁶, CONHOH, SO₂NR⁶R⁶, SO₂CF₃, SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO₂alkyl, CONHSO₂aryl, aryl, heteroaryl; wherein C₁-C₆ alkyl is straight or branched, heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;

25 alkyl of 1 to 18 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

alkenyl of 3 to 18 carbon atoms having from 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;

30 alkynyl of 3 to 18 carbon atoms having from 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;

arylalkyl of 7 to 16 carbon atoms, wherein aryl is optionally substituted with one or two groups selected independently from R⁵;


WO 99/42436

PCT/US98/17633

- 9 -

- heteroarylalkyl wherein alkyl is from 1 to 6 carbon atoms and heteroaryl contains 1 or 2 heteroatoms selected from O, S or N and is optionally substituted with one or two groups selected independently from R⁵;
- biphenylalkyl of 13 to 18 carbon atoms, wherein biphenyl is optionally substituted with one or two groups selected independently from R⁵;
- arylalkenyl of 8 to 16 carbon atoms, wherein aryl is optionally substituted with one or two groups selected independently from R⁵;
- cycloalkylalkyl or bicycloalkylalkyl of 4 to 12 carbon atoms, wherein cycloalkyl or bicycloalkyl is optionally substituted with one or two groups selected independently from R⁵;
- saturated or unsaturated mono or bicyclic heterocycle containing one heteroatom selected from O, S or N-C₁-C₆ alkyl, optionally substituted with one or two groups selected independently from R⁵;
- R⁸R⁹N-C₁-C₆-alkoxyaryl-C₁-C₆-alkyl where R⁸ and R⁹ are independently selected from C₁-C₆ alkyl or R⁸ and R⁹ together with the interposed nitrogen forms a 5-7 membered saturated heterocyclic ring optionally containing an oxygen atom, wherein the aryl group is phenyl or naphthyl;
- and the pharmaceutically acceptable salts thereof.

- The most preferred group of compounds are those of the following formula (Ib):



in which

- R¹ is phenyl, naphthyl, alkyl of 1-18 carbon atoms or heteroaryl such as pyridyl, thienyl, imidazolyl or furanyl, optionally substituted with C₁-C₆ alkyl, C₁-C₆ alkoxy, C₆-C₁₀ aryloxy, heteroaryloxy, C₃-C₆ alkenyloxy, C₃-C₆ alkynyloxy, halogen; or S(O)_n-C₁-C₆alkyl C₁-C₆ alkoxyaryl or C₁-C₆ alkoxyheteroaryl;
- A is -S-, -SO- or -SO₂-;
- R² and R³, taken with the carbon atom to which they are attached, form a 5 to 7 membered heterocyclic ring containing O, S or N-R⁷ optionally having one or two double bonds;

WO 99/42436

PCT/US98/17633

- 10 -

R⁴ is hydrogen,

alkyl of 1 to 6 carbon atoms, optionally substituted with one or two groups
selected independently from R⁵;

alkenyl of 3 to 18 carbon atoms having 1 to 3 double bonds, optionally

5 substituted with one or two groups selected independently from R⁵;

alkynyl of 3 to 18 carbon atoms having 1 to 3 triple bonds, optionally

substituted with one or two groups selected independently from R⁵;

phenyl or naphthyl optionally substituted with one or two groups selected
independently from R⁵;

10 C₃ to C₈ cycloalkyl or bicycloalkyl optionally substituted with one or two
groups selected independently from R⁵;

R⁵ is H, C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁ to C₁₂ alkyl, C₂ to C₁₂ alkenyl, C₂-C₁₂
alkynyl, F, Cl, Br, I, CN, CHO, C₁-C₆ alkoxy, aryloxy, heteroaryloxy, C₃-C₆
alkenyloxy, C₃-C₆ alkynyloxy, C₁-C₆ alkylamino-C₁-C₆ alkoxy, C₁-C₂
15 alkylene dioxy, aryloxy-C₁-C₆ alkyl amine, C₁-C₁₂ perfluoro alkyl, S(O)_n-C₁-
C₆ alkyl, S(O)_n-aryl where n is 0, 1 or 2; OCOO C₁-C₆ alkyl, OCOOaryl,
OCONR⁶, COOH, COO C₁-C₆ alkyl, COOaryl, CONR⁶R⁶, CONHOH,
NR⁶R⁶, SO₂NR⁶R⁶, NR⁶SO₂aryl, -NR⁶CONR⁶R⁶, NHSO₂CF₃,
SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl, CONHSO₂aryl,
20 SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl, CONHSO₂aryl, NH₂, OH, aryl,
heteroaryl, C₃ to C₈ cycloalkyl; saturated or unsaturated 5 to 10 membered
mono or bicyclic heterocycle containing one heteroatom selected from O, S or
NR⁷, wherein C₁-C₆ alkyl is straight or branched, heteroaryl is a 5-10
membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected
25 independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally
substituted by 1 or 2 groups selected from halogen, cyano, amino, nitro, C₁-C₆
alkyl, C₁-C₆ alkoxy, or hydroxy;

R⁶ is H, C₁ to C₁₈ alkyl optionally substituted with OH; C₃ to C₆ alkenyl, C₃ to C₆
alkynyl, C₁ to C₆ perfluoro alkyl, S(O)_n alkyl or aryl where n is 0, 1 or 2; or
30 COheteroaryl; wherein heteroaryl is a 5-10 membered mono or bicyclic
heteroaryl group having 1 to 3 heteroatoms selected independently from
O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1

WO 99/42436

PCT/US98/17633

- 11 -

or 2 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;

and R⁷ is C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁-C₁₂ perfluoro alkyl, S(O)_n-alkyl, S(O)_n-aryl where n is 0, 1 or 2; COOalkyl, COOaryl, CONHR⁶, CONR⁶R⁶,

5 CONHOH, SO₂NR⁶R⁶, SO₂CF₃, SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO₂alkyl, CONHSO₂aryl, aryl, or heteroaryl; where aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected independently from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy; and heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or N-
10 C₁-C₆ alkyl;

alkyl of 1 to 18 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

alkenyl of 3 to 18 carbon atoms having from 1 to 3 double bonds, optionally

15 substituted with one or two groups selected independently from R⁵;

alkynyl of 3 to 18 carbon atoms having from 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;

arylalkyl of 7 to 16 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

20 heteroarylalkyl wherein alkyl is from 1 to 6 carbon atoms and heteroaryl contains 1 or 2 heteroatoms selected from O, S or N and is optionally substituted with one or two groups selected independently from R⁵;

biphenylalkyl of 13 to 18 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

25 arylalkenyl of 8 to 16 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

cycloalkylalkyl or bicycloalkylalkyl of 4 to 12 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

30 saturated or unsaturated mono or bicyclic heterocycle containing one heteroatom selected from O, S or NR-C₁-C₆ alkyl, optionally substituted with one or two groups selected independently from R⁵;

R⁸R⁹N-C₁-C₆-alkoxyaryl-C₁-C₆-alkyl where R⁸ and R⁹ are independently selected from C₁-C₆ alkyl or R⁸ and R⁹ together with the interposed nitrogen

WO 99/42436

PCT/US98/17633

- 12 -

forms a 5-7 membered saturated heterocyclic ring optionally containing an oxygen atom, wherein the aryl group is phenyl or naphthyl; and the pharmaceutically acceptable salts thereof.

- 5 The most preferred matrix metalloproteinase and TACE inhibiting compounds of this invention are:
- 1-benzyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide,
4-(4-methoxy-benzenesulfonyl)-1-(3-methoxy-benzyl)-piperidine-4-carboxylic acid
hydroxyamide,
- 10 1-(3,4-dichlorobenzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid
hydroxamide,
4-(4-methoxy-benzenesulfonyl)-1-(4-methylbenzyl)-piperidine-4-carboxylic acid
hydroxamide,
4-(4-methoxy-benzene-sulfonyl)-1-naphthalene-2-yl-methylpiperidine-4-carboxylic acid
15 hydroxamide,
1-biphenyl-4-ylmethyl-4-(4-methoxy-benzenesulfonyl)piperidine-4-carboxylic acid
hydroxamide,
4-(4-methoxy-benzene-sulfonyl)-1-(3-methyl-but-2-enyl)piperidine-4-carboxylic acid
hydroxamide,
- 20 1-(4-bromo-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid
hydroxyamide,
4-(4-methoxy-benzenesulfonyl)-1-(3-phenyl-propyl)-piperidine-4-carboxylic acid
hydroxyamide,
1-tert-butyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid
25 hydroxyamide,
1-butyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide,
1-cyclooctyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid
hydroxyamide,
1-ethyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide,
- 30 1-isopropyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid
hydroxyamide,
1-methyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide,
1-benzyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide,
1-(4-fluoro-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid
35 hydroxyamide,
1-(4-fluoro-benzyl)-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid
hydroxyamide,

WO 99/42436

PCT/US98/17633

- 13 -

- 4-(4-methoxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid
hydroxyamide,
- 4-(4-methoxy-benzenesulfonyl)-1-[2-(4-methoxyphenyl)-ethyl]-piperidine-4-carboxylic
acid hydroxyamide,
- 5 4-(4-methoxy-benzenesulfonyl)-1-(2-phenyl-ethyl)-piperidine-4-carboxylic acid
hydroxyamide,
- 4-(4-n-butoxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid
hydroxyamide,
- 4-(4-methoxy-benzenesulfonyl)-1-(3-phenoxy-propyl)-piperidine-4-carboxylic acid
10 hydroxyamide,
- 4-(4-n-butoxy-benzenesulfonyl)-1-(3-phenoxy-propyl)-piperidine-4-carboxylic acid
hydroxyamide,
- 4-(4-methoxy-benzenesulfonyl)-1-(2-phenoxy-ethyl)-piperidine-4-carboxylic acid
hydroxyamide,
- 15 4-(4-n-butoxy-benzenesulfonyl)-1-(2-phenoxy-ethyl)-piperidine-4-carboxylic acid
hydroxyamide,
- 4-(4-methoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-
carboxylic acid hydroxyamide,
- 1-Benzyl-4-(4-benzyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid
20 hydroxyamide,
- 4-(4-Butoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-
carboxylic acid hydroxyamide,
- 4-(4-Butoxy-benzenesulfonyl)-1-[3-(2-morpholinyl-1-yl-ethoxy)-benzyl]-piperidine-4-
carboxylic acid hydroxyamide,
- 25 1-Methyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide,
1-Ethyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide,
1-n-Butyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide,
4-[4-(4-chloro-phenoxy)-benzenesulfonyl]1-methyl-piperidine-4-carboxylic acid
hydroxyamide,
- 30 4-[4-(4-chloro-phenoxy)-benzenesulfonyl]1-ethyl-piperidine-4-carboxylic acid
hydroxyamide,
- 1-Butyl-4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid
hydroxyamide,
- 1-Benzyl-4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid
35 hydroxyamide,
- 1-Benzyl-4-[4-(3-methyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid
hydroxyamide,

WO 99/42436

PCT/US98/17633

- 14 -

- 1-Butyl-4-[4-(3-methyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid
hydroxyamide,
1-Benzyl-4-[4-(2-ethyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid
hydroxyamide,
5 4-(4-butoxy-benzenesulfonyl)-1-(3-methoxy-benzyl)-piperidine-4-carboxylic acid
hydroxyamide,
4-(4-Methoxy-benzenesulfonyl)-1-(4-thiophen-2-yl-benzyl)-piperidine-4-carboxylic
acid hydroxyamide,
4-(4-methoxy-benzenesulfonyl)-1-(4-pyridin-2-yl-benzyl)-piperidine-4-carboxylic acid
10 hydroxyamide,
1-(3,4-Dichlorobenzyl)-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid
hydroxyamide,
[4-(4-Chloro-benzyloxy)-benzenesulfonyl]-1-methylpiperidine-4-carboxylic acid
hydroxamide,
15 4-(4-Butoxy-benzenesulfonyl)-1-(3-phenoxy-benzyl)-piperidine-4-carboxylic acid
hydroxamide,
[4-(4-Chloro-benzyloxy)-benzenesulfonyl]-1-(4-methylbenzyl)-piperidine-4-
carboxylic acid hydroxamide,
4-(4-Butoxy-benzenesulfonyl)-1-(4-methylbenzyl)-piperidine-4- carboxylic acid
20 hydroxamide,
4-(4-Butoxy-benzenesulfonyl)-1-(4-cyano-benzyl)-piperidine-4-carboxylic acid
hydroxamide, and
4-(4-Butoxy-benzenesulfonyl)-1-pyridin-4-ylmethyl-piperidine-4-carboxylic acid
hydroxamide, and pharmaceutical salts thereof.

25

It is understood that the definition of the compounds of formulas I, Ia and Ib,
when R¹, R², R³ and R⁴ contains asymmetric carbons, encompass all possible
stereoisomers and mixtures thereof which posses the activity discussed below. In
particular, it encompasses racemic modifications and any optical isomers which
30 possesses the indicated activity. Optical isomers may be obtained in pure form by
standard separation techniques. Where not stated otherwise, the term "alkyl" refers to a
straight or branched C₁-C₆ alkyl group and aryl is phenyl or naphthyl. The
pharmaceutically acceptable salts are those derived from pharmaceutically acceptable
organic and inorganic acids such as lactic, citric, acetic, tartaric, succinic, maleic,
35 malonic, hydrochloric, hydrobromic, phosphoric, nitric, sulfuric, methanesulfonic, and
similarly known acceptable acids.

WO 99/42436

PCT/US98/17633

- 15 -

The present invention accordingly provides a pharmaceutical composition which comprises a compound of this invention in combination or association with a pharmaceutically acceptable carrier. In particular, the present invention provides a pharmaceutical composition which comprises an effective amount of compound of this invention and a pharmaceutically acceptable carrier.

The compositions are preferably adapted for oral administration. However, they may be adapted for other modes of administration, for example, parenteral administration for patients.

In order to obtain consistency of administration, it is preferred that a composition of the invention is in the form of a unit dose. Suitable unit dose forms include tablets, capsules, and powders in sachets or vials. Such unit dose forms may contain from 0.1 to 100 mg of a compound of the invention. The compounds of the present invention can be administered orally at a dose range of about 0.01 to 100 mg per kg. Such composition may be administered from 1 to 6 times a day, more usually from 1 to 4 times a day.

The compositions of the invention may be formulated with conventional excipients, such as fillers, a disintegrating agent, a binder, a lubricant, a flavoring agent, and the like. They are formulated in conventional manner.

Also according to the present invention, there are provided processes for producing the compounds of the present invention.

PROCESS OF THE INVENTION.

The compounds of the present invention may be prepared according to one of the general processes out lined below.


The appropriately substituted mercaptan derivative was alkylated using either substituted (Scheme 1) or unsubstituted (Scheme 2) α -bromo acetic acid ester derivative in refluxing acetone using K_2CO_3 as base. The sulphide derivative thus obtained was oxidized using m-chloroperbenzoic acid in CH_2Cl_2 or by using Oxone in methanol/ water. The sulfone obtained from the above mentioned process can be either further alkylated using variety of alkyl halides to obtain the disubstituted derivative or it can be hydrolyzed using NaOH/ MeOH at room temp. However instead of using the ethyl ester, if the tertiary butyl ester is present, the hydrolysis can be carried out with TFA/ CH_2Cl_2 at room temperature. Subsequently, the carboxylic acid obtained was converted to the hydroxamic acid derivative by reaction with oxalyl chloride/ DMX (catalytic) and hydroxyl amine/ triethyl amine.

WO 99/42436

PCT/US98/17633

- 16 -

SCHEME 1

SYNTHESIS:


- a. K_2CO_3 / Acetone/ Reflux; b. m-Chloroperbenzoic acid;
c. K_2CO_3 / 18-Crown-6/ R_3Br /Acetone/ Reflux/
d. NaOH/ MeOH/ THF/ RT
e. $(COCl)_2/CH_2Cl_2/Et_3N/NH_2OH \cdot HCl$.

WO 99/42436

PCT/US98/17633

- 17 -

SCHEME 2

SYNTHESIS:

a. K_2CO_3 / Acetone/ Reflux; b. m-Chloroperbenzoic acid;
 c. K_2CO_3 / 18-Crown-6/ R_2Br /Acetone/ Reflux/
 d. R_3Br / 10 N NaOH/ $\text{BzN}(\text{Et})_3$ / CH_2Cl_2 / RT
 e. NaOH/ MeOH/ THF/ RT
 f. $(\text{COCl})_2$ / CH_2Cl_2 / Et_3N / $\text{NH}_2\text{OH.HCl}$.


As outlined in Scheme 3, the sulfide derivative can be further alkylated using
 5 lithium bis(trimethyl silyl)amide in THF at 0°C . The alkylated or mono substituted
 compound was hydrolyzed and converted to the hydroxamic acid derivative. The
 sulfinyl derivatives were prepared by oxidizing the sulfide hydroxamic acid derivatives
 with H_2O_2 in MeOH solution.

WO 99/42436

PCT/US98/17633

- 18 -

SCHEME - 3

SYNTHESIS:

a. K_2CO_3 / Acetone/ Reflux; b. R_3Br / HMDS/ THF;
 c. NaOH/ MeOH/ THF/ RT
 d. $(\text{COCl})_2/\text{CH}_2\text{Cl}_2/\text{Et}_3\text{N}/\text{NH}_2\text{OH.HCl}$.
 e. MeOH/ H_2O_2 / RT

The corresponding 1-substituted-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamides were prepared starting from diethanolamine and appropriately substituted alkyl or aryl halides (Scheme 4). The N-substituted diethanol amine derivatives were converted to the dichloro compounds using thionyl chloride. The corresponding dichlorides were reacted with substituted sulfonyl acetic acid ethyl ester derivatives in the presence of K_2CO_3 /18-Crown-6 in boiling acetone. 1-substituted-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl esters thus obtained were converted to the hydroxy amide as outlined in Scheme 4.


WO 99/42436

PCT/US98/17633

- 19 -


Alternatively these classes of compounds and other hetrocycles can be prepared as indicated in Scheme 5 and 6.

SCHEME 4




5

SCHEME 5



a. RBr / R^1SH / $CHCl_3$ / Reflux; b. Oxone/ MeOH; e. $(COCl)_2/NH_2OH.HCl/Et_3N$

SCHEME 6




Alternatively, Schemes 7 to 11 show methods for the preparation of hydroxamic acid compounds using a solid phase support (P).

WO 99/42436

PCT/US98/17633

- 20 -

Scheme 7



Reagents and Conditions: a) 2-Halo acid (3.0 eq.); 1-hydroxybenzotriazole hydrate (HOBt, 6.0 eq.); 1,3-diisopropylcarbodiimide (DIC, 4.0 eq.); DMF, 25°C; 2-16 hours. b) Thiol (5.0 eq.); sodium iodide (5.0 eq.); 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 3.0 eq.); THF; 25°C; 12-16 hours. c) 70% *tert*-butylhydroperoxide (40 eq.); benzenesulfonic acid (2.0 eq.); DCM; 25°C; 12-24 hours. d) *m*CPBA (5.0 eq.); DCM; 25°C; 12-24 hours. e) TFA : DCM (1:1); 25°C; 1 hour.

10

The 4-*O*-methylhydroxylamine-phenoxyethyl-copoly(styrene-1%-divinylbenzene)-resin (hydroxylamine resin) may be coupled with a 2-halo acid to give the hydroxamate ester resin. The coupling reaction may be carried out in the presence of carbodiimide, such as DIC, in an inert solvent such as DMF at room temperature. The halogen group may be displaced with a thiol in the presence of a base, such as DBU, in an inert solvent such as THF at room temperature. The sulfide may be oxidized to the sulfoxide by reaction with an oxidizing agent such as *tert*-butylhydroperoxide in the presence of an acid catalyst such as benzenesulfonic acid, in an inert solvent such as DCM at room temperature. Alternatively, the sulfide may be oxidized to the sulfone by reaction with an oxidizing agent such as *meta*-chloroperoxybenzoic acid, in an inert solvent such as DCM at room temperature. The sulfide, sulfoxide, or sulfone may be

WO 99/42436


PCT/US98/17633

- 21 -

treated with and acid, such as trifluoroacetic acid, in and inert solvent such as DCM to liberate the free hydroxamic acid.

Scheme 8 shows a method of preparing hydroxamic acids having alkoxy groups attached to the aromatic ring.

Scheme 8



- 10 Reagents and Conditions: a) 2-Halo acid (3.0 eq.); 1-hydroxybenzotriazole hydrate (HOBt, 6.0 eq.); 1,3-diisopropylcarbodiimide (DIC, 4.0 eq.); DMF, 25°C; 2-16 hours. b) 4-Fluorobenzenethiol (5.0 eq.); sodium iodide (5.0 eq.); 1,8-diazabicyclo[5.4.0]-undec-7-ene (DBU, 3.0 eq.); THF; 25°C; 12-16 hours. c) Alcohol (15.0 eq.); sodium hydride (15.0 eq.); DMF; 80°C; 15 hours. d) 70% *tert*-butylhydroperoxide (40 eq.); benzenesulfonic acid (2.0 eq.); DCM; 25°C; 12-24 hours. e) mCPBA (5.0 eq.); DCM; 25°C; 12-24 hours. f) TFA : DCM (1:1); 25°C; 1 hour.

The hydroxylamine resin may be coupled with the 2-halo acid and the halo group may be displaced by fluorobenzenethiol as previously described. The fluoro

WO 99/42436

PCT/US98/17633


- 22 -

group may then be displaced with an alcohol in the presence of a base such as sodium hydride, in an inert solvent such as DMF at about 80°C. The alkoxybenzenesulfanyl hydroxamate ester may then be oxidized either to the corresponding sulfinyl or sulfonyl hydroxamate ester as previously described. The free hydroxamic acids may be liberated as previously described.

Scheme 9 shows a method of preparing 2-bisarylsulfanyl-, sulfinyl-, and sulfonylhydroxamic acids.

10

Scheme 9



Reagents and Conditions: a) 2-Halo acid (3.0 eq.); 1-hydroxybenzotriazole hydrate (HOBt, 6.0 eq.); 1,3-diisopropylcarbodiimide (DIC, 4.0 eq.); DMF, 25°C; 2-16 hours.

b) 4-Bromobenzenethiol (5.0 eq.); sodium iodide (5.0 eq.); 1,8-diazabicyclo[5.4.0]-undec-7-ene (DBU, 3.0 eq.); THF; 25°C; 12-16 hours. c) 70% *tert*-butylhydroperoxide (40 eq.); benzenesulfonic acid (2.0 eq.); DCM; 25°C; 12-24 hours. d) *m*CPBA (5.0 eq.); DCM; 25°C; 12-24 hours. e) Arylboronic acid (2.0 eq.); tetrakis(triphenylphosphine) palladium(0) (0.1 eq.); 10% aqueous sodium carbonate (10.0 eq.); DME; 80°C; 8 hours. f) TFA : DCM (1:1); 25°C; 1 hour.


WO 99/42436

PCT/US98/17633

- 23 -

- The hydroxylamine resin may be coupled with the 2-halo acid and the halo group may be displaced by bromobenzenethiol as previously described. The bromobenzenesulfanyl hydroxamate ester may then be oxidized either to the corresponding sulfinyl or sulfonyl hydroxamate ester as previously described. The
- 5 bromo group may then be replaced with an aryl group by reaction with the arylboronic acid in the presence of a catalyst such as tetrakis(triphenylphosphine) palladium(0), and a base such as sodium carbonate, in an inert solvent such as DME at about 80°C. The free hydroxamic acids may be liberated as previously described.
- 10 Scheme 10 shows a method of preparing hydroxamic acids having amine groups attached to the aromatic ring.

Scheme 10



- 15 Reagents and Conditions: a) 2-Halo acid (3.0 eq.); 1-hydroxybenzotriazole hydrate (HOBt, 6.0 eq.); 1,3-diisopropylcarbodiimide (DIC, 4.0 eq.); DMF, 25°C; 2-16 hours. b) 4-Bromobenzenethiol (5.0 eq.); sodium iodide (5.0 eq.); 1,8-diazabicyclo[5.4.0]-undec-7-ene (DBU, 3.0 eq.); THF; 25°C; 12-16 hours. c) Amine (20.0 eq.);
- 20 tris(dibenzylideneacetone)-dipalladium(0) (0.2 eq.); (S)-(-)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl ((S)-BINAP, 0.8 eq.); sodium *tert*-butoxide (18.0 eq.); dioxane; 80°C, 8 hours; d) TFA : DCM (1:1); 25°C; 1 hour.

- The hydroxylamine resin may be coupled with the 2-halo acid and the halo group may be displaced by bromobenzenethiol as previously described. The bromo
- 25 group may then be displaced with an amine in the presence of a catalyst such as

WO 99/42436


PCT/US98/17633

- 24 -

tris(dibenzylideneacetone)-dipalladium(0) and a ligand such as (S)-BINAP and a base such as sodium *tert*-butoxide, in an inert solvent such as dioxane at about 80°C. The free hydroxamic acids may be liberated as previously described.

- 5 Scheme 11 shows a method of preparing hydroxamic acids having sulfonate groups attached to the aromatic ring.

Scheme 11



- 10 Reagents and Conditions: a) 2-Halo acid (3.0 eq.); 1-hydroxybenzotriazole hydrate (HOBt, 6.0 eq.); 1,3-diisopropylcarbodiimide (DIC, 4.0 eq.); DMF, 25°C; 2-16 hours. b) 4-Hydroxybenzenethiol (5.0 eq.); sodium iodide (5.0 eq.); 1,8-diazabicyclo-[5.4.0]undec-7-ene (DBU, 3.0 eq.); THF; 25°C; 12-16 hours. c) Sulfonyl chloride (5.0 eq.); triethylamine (2.0 eq.); DCM; 25°C; 8 hours. d) 70% *tert*-butylhydroperoxide (40 eq.); benzenesulfonic acid (2.0 eq.); DCM; 25°C; 12-24 hours. e) mCPBA (5.0 eq.); DCM; 25°C; 12-24 hours. f) TFA : DCM (1:1); 25°C; 1 hour.
- 15

WO 99/42436

PCT/US98/17633

- 25 -

The hydroxylamine resin may be coupled with the 2-halo acid and the halo group may be displaced by hydroxybenzenethiol as previously described. The hydroxybenzenesulfanyl hydroxamate ester may then be oxidized either to the corresponding sulfinyl or sulfonyl hydroxamate ester as previously described. The hydroxy group may then be sulfonylated by reaction with a sulfonyl chloride in the presence of a base such as triethylamine, in an inert solvent such as DCM at about room temperature. The free hydroxamic acids may be liberated as previously described.

The following examples are presented to illustrate rather than limit the scope of the invention. HPLC purity of compounds prepared by combinatorial procedures is presented as area percentage at a prescribed wavelength (% @ nm).

Example 1

N-Hydroxy-2-(4-methoxy-phenylsulfanyl)-2-methyl-3-phenyl-propionamide

To stirred solution of 4-methoxybenzenethiol (2.8 gm, 20 mmol) and anhydrous K_2CO_3 (10 gm, excess) in dry acetone (100 ml), ethyl 2-bromo-propionate (3.6 gm, 20 mmol) was added in a round bottom flask and the reaction mixture was heated at reflux for 8 hours with good stirring. At the end, reaction was allowed to cool and the potassium salts were filtered off and the reaction mixture was concentrated. The residue was extracted with chloroform and washed with H_2O and 0.5 N NaOH solution. The organic layer was further washed well with water, dried over $MgSO_4$, filtered and concentrated to afford 2-(4-methoxy-phenylsulfanyl)-propionic acid ethyl ester as a light yellow oil. Yield 4.5gms (94%); MS; 241 (M+H)⁺.

To a stirred solution of 2-(4-methoxy-phenylsulfanyl)-propionic acid ethyl ester (2.44 g, 10 mmol), in THF (100 ml) at -4°C, lithium bis(trimethylsilyl)amide (1 M solution, 15 ml, 15 mmol) was added slowly. The orange colored reaction mixture was stirred at room temperature for 15 minutes and then it was cooled to 0°C at which time it was stirred for an additional hour. The temperature of the mixture was again brought to -40°C and benzylbromide (1.72 gm, 10 mmol) was added dropwise in THF. The reaction was warmed to room temperature and stirred overnight before it was quenched with ice water, extracted with chloroform and washed with water. The organic layer was dried over $MgSO_4$, filtered and concentrated and chromatographed on a silica-gel column with 10% ethyl acetate:hexane to afford 2-(4-methoxy-phenylsulfanyl)-2-

WO 99/42436

PCT/US98/17633

- 26 -

methyl-3-phenyl-propionic acid ethyl ester as a colorless oil. Yield: 860 mg, (26%) ;

MS: 331 (M+H)⁺.

2-(4-methoxy-phenylsulfanyl)-2-methyl-3-phenyl-propionic acid ethyl ester (4.12 g,
5 12 mmol) dissolved in methanol (50 ml) and 10 N NaOH (20 ml) was added. The
reaction was allowed to stir overnight at room temperature. The reaction mixture was
concentrated, diluted with 1:1 hexane:diethyl ether and extracted with H₂O. The water
layer was cooled with ice and acidified to pH 3. The acid was then extracted with
chloroform and the organic layer was dried over MgSO₄, filtered and concentrated to
10 afford of 2-(4-methoxy-phenylsulfanyl)-2-methyl-3-phenyl-propionic acid as a low
melting solid. Yield 580 mg, 16%; MS: 303.2 (M+H)⁺.

To a stirred solution of 2-(4-methoxy-phenylsulfanyl)-2-methyl-3-phenyl-propionic
acid (0.5 g, 1.65 mmol) and DMF (2 drops) in CH₂Cl₂ (100 ml) at 0°C, oxalyl
15 chloride (1.0 gm, 8 mmol) was added in a drop-wise manner. After the addition, the
reaction mixture was stirred at room temperature for 1 hour. Simultaneously, in a
separate flask a mixture of hydroxylamine hydrochloride (2.0 gm, 29 mmol) and
triethylamine (5 ml, excess) was stirred in THF:water (5:1, 30 ml) at 0°C for 1 hour.
At the end of 1 hour, the oxalyl chloride reaction mixture was concentrated and the pale
20 yellow residue was dissolved in 10 ml of CH₂Cl₂ and added slowly to the hydroxyl-
amine at 0°C. The reaction mixture was stirred at room temperature for 24 hours and
concentrated. The residue obtained was extracted with chloroform and washed well
with water. The product obtained was purified by silica gel column chromatography
and eluted with ethyl acetate. The N-hydroxy-2-(4-methoxyphenylsulfanyl)-2-methyl-
25 3-phenyl-propionamide was isolated as a colorless solid. mp 88 °C; Yield, 300 mg,
57%; MS: 318 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.32 (s, 3H), 3.07 (d, J = 11
Hz, 1H), 3.23 (d, J = 11 Hz, 1H), 3.79 (s, 3H), 6.83-7.36 (m, 9H).

Example 2

30

N-Hydroxy-2-(4-methoxy-phenylsulfanyl)-2-phenyl-acetamide

2-(4-Methoxyphenylsulfanyl)-phenylacetic acid ethyl ester was prepared according to
the general method as outlined in Example 1. Starting from ethyl α-bromophenyl

WO 99/42436

PCT/US98/17633

- 27 -

acetate (7.18 g, 31.4 mmol) and 4-methoxythiophenol (4.4 g, 31.4 mmol), 8.5 g of the product was isolated as a light yellow oil. Yield 90%; MS: 303.1 (M+H)⁺.

5 2-(4-Methoxy-phenylsulfanyl)-2-phenyl acetic acid was prepared starting from 2-(4-methoxy-phenylsulfanyl)-phenyl-acetic acid ethyl ester (3.0 g, 10 mmol) dissolved in methanol (50 ml) and 10 N NaOH (20 ml). The resulting reaction mixture was worked up as in Example 1. Yield 1.9 g, 70%. Low melting solid. MS: 273 (M+H)⁺.

10 Starting from 2-(4-methoxy-phenylsulfanyl)-phenyl acetic acid (1.05 g, 3.83 mmol) and following the procedure as outlined in Example 1, 154 mg of N-hydroxy-2-(4-methoxy-phenylsulfanyl)-2-phenyl-acetamide was isolated as a colorless solid. mp 155°C; Yield 14%; MS: 290 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 3.72 (s, 3H), 4.68 (s, 1H), 6.86-7.44 (m, 9H).

15

Example 3

2-(4-Methoxy-phenylsulfanyl)-2,5-dimethyl-hex-4-enoic acid hydroxyamide

20 2-(4-Methoxy-phenylsulfanyl)-2,5-dimethyl-hex-4-enoic acid ethyl ester was prepared following the procedure of Example 1, second paragraph. Starting from (4-methoxy-phenylsulfanyl)-propionic acid ethyl ester (3.5 g, 14.3 mmol), and isoprenyl bromide (2.25 g, 15 mmol), 2.2 g of the product was isolated as an oil. Yield 50%; MS: 310 (M+H)⁺.

25 2-(4-Methoxy-phenylsulfanyl)-2,5-dimethyl-hex-4-enoic acid was prepared starting from 2-(4-methoxy-phenylsulfanyl)-2,5-dimethyl-hex-4-enoic acid ethyl ester (2.0 g, 6.4 mmol) dissolved in methanol (50 ml) and 10 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in Example 1. Yield is 1.9 g, 99% of low melting solid. MS: 280 (M+H)⁺.

30

Starting from 2-(4-methoxy-phenylsulfanyl)-2,5-dimethyl-hex-4-enoic acid (1.67 g, 5.8 mmol) and following the procedure as outlined in Example 1, 1.5 g of 2-(4-methoxy-phenylsulfanyl)-2,5-dimethyl-hex-4-enoic acid hydroxyamide was isolated as a colorless solid. mp 89 °C; Yield 94%; MS: 296 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.34 (s, 3H), 1.61 (s, 3H), 1.74 (s, 3H), 2.41-2.58 (m, 2H), 3.80 (s, 3H), 35 5.17 (t, J = 7.5 Hz, 1H), 6.86 (d, J = 12.4 Hz, 2H), 7.35 (d, J = 12.4 Hz, 2H).

WO 99/42436

PCT/US98/17633

- 28 -

Example 4

N-Hydroxy-2-(4-methoxy-phenylsulfanyl)-3-methyl-butyramide

- 5 2-(4-Methoxy-phenylsulfanyl)-3-methyl-butyric acid ethyl ester was prepared according to the general method of Example 1. Starting from ethyl 2-bromo-3-methyl-butanoate (20.9 g, 100 mmol) and 4-methoxybenzenethiol (14.0 g, 100 mmol), 30 g of the product was isolated. Yield 99%; Light yellow oil; MS: 271 (M+H)⁺.
- 10 2-(4-Methoxy-phenylsulfanyl)-3-methyl-butyric acid was prepared starting from 2-(4-methoxy-phenylsulfanyl)-3-methyl-butyric acid ethyl ester (5.8 g, 21.6 mmol) dissolved in methanol (50 ml) and 10 N NaOH (30 ml). The resulting reaction mixture was worked up as outlined in Example 1. Yield 5.0 g, 99%. Low melting solid. MS: 242 (M+H)⁺.
- 15 Starting from 2-(4-methoxy-phenylsulfanyl)-3-methyl-butyric acid (4.39 g, 18.3 mmol) and following the procedure as outlined in Example 1, 1.5 g of N-hydroxy-2-(4-methoxy-phenylsulfanyl)-3-methyl-butyramide was isolated as a colorless solid. mp 119 °C; Yield 33%; MS: 256 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.90-1.07
- 20 (m, 6H), 1.84-1.96 (m, 1H), 3.07 (d, J = 8.8 Hz, 1H), 3.75 (s, 3H), 6.88 (d, J = 15 Hz, 2H), 7.35 (d, J = 15 Hz, 2H).

Example 5

- 25 N-Hydroxy-2-(4-methoxy-benzenesulfanyl)-2-methyl-3-phenyl-propionamide
- N-hydroxy-2-(4-methoxy-phenylsulfanyl)-2-methyl-3-phenyl-propionamide (400 mg, 1.26 mmol) (prepared in Example 1) was dissolved in methanol (100 ml) and 30% H₂O₂ (10 ml) was added. The reaction mixture was stirred for 48 hours at room
- 30 temperature at which time it was cooled to 0° C and quenched with saturated Na₂SO₃ (20 ml) solution. The reaction mixture became cloudy. It was stirred for 4 hours before it was concentrated in a room temperature water bath, diluted with water, extracted with CHCl₃ and washed with H₂O. The organic layer was dried over MgSO₄, filtered and concentrated. The title compound was isolated by silica gel column
- 35 chromatography, eluting with 75% ethylacetate:hexane. Low melting solid. Yield: 220 mg (52%); MS: 334.1 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.11 (s, 2H), 1.22 (s, 3H), 3.84 (s, 3H), 7.00-7.61 (m, 9H).

WO 99/42436

PCT/US98/17633

- 29 -

Example 6

2-(4-Methoxy-benzenesulfinyl)-2,5-dimethyl-hex-4-enoic acid hydroxyamide

- 5 Starting from 2-(4-methoxy-benzenesulfanyl)-2,5-dimethyl-hex-4-enoic hydroxamide (900 mg, 3.0 mmol) (prepared in Example 3) and following the procedure outlined in Example 5, 2-(4-methoxy-benzenesulfinyl)-2,5-dimethyl-hex-4-enoic acid hydroxy-
amide was isolated as a colorless solid. Yield: 104 mg (10%); mp 108 °C; MS: 312 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.88 (s, 3H), 1.59 (s, 3H), 1.68 (s, 3H),
10 2.27-2.80 (m, 2H), 5.02 (t, J = 7.5 Hz, 1H), 7.09 (d, J = 9 Hz, 2H), 7.39 (d, J = 9 Hz, 2H).

Example 7

15 N-Hydroxy-2-(4-methoxy-benzenesulfinyl)-3-methyl-butyramide

- Starting from N-hydroxy-2-(4-methoxy-phenylsulfanyl)-3-methyl-butyramide (1 g, 3.9 mmol) as prepared in Example 4, and following the procedure of Example 5, N-hydroxy-2-(4-methoxy-benzenesulfinyl)-3-methyl-butyramide was isolated as a
20 colorless solid. Yield: 420mg (40%); mp 163 °C; MS: 272 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.89-1.12 (m, 6H), 1.63-1.74 (m, 1H), 3.13 (d, J = 7 Hz, 1H), 3.83 (s, 3H), 6.94-7.65 (m, 4H).

Example 8

25 N-Hydroxy-2-(4-methoxy-benzenesulfinyl)-2-phenyl-acetamide

- Starting from N-hydroxy-2-(4-methoxy-phenylsulfanyl)-2-phenyl-acetamide (240 mg, 0.83 mmol) as prepared in Example 2, and following the procedure outlined in
30 Example 5, N-hydroxy-2-(4-methoxy-benzenesulfinyl)-2-phenyl-acetamide was isolated as colorless solid. Yield: 100mg (40%); mp 135 °C; MS 304 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 3.75 (s, 3H), 4.38 (s, 1H), 6.92-7.69 (m, 9H)

Example 9

35 N-Hydroxy-2-(4-methoxy-benzenesulfonyl)-3-phenyl-propionamide.

- To a stirred solution of 4-methoxybenzenethiol (2.8 gm, 20 mmol) and anhydrous K₂CO₃ (10 gm, excess) in dry acetone (100 ml), α-bromo ethyl acetate (3.3 gm, 20
40 mmol) was added in a round bottom flask and the reaction mixture was heated at reflux

WO 99/42436

PCT/US98/17633

- 30 -

for 8 hours with good stirring. At the end, the reaction mixture was allowed to cool and the potassium salts were filtered off and the reaction mixture was concentrated. The residue was extracted with chloroform and washed with H₂O and 0.5 N NaOH solution. The organic layer was further washed well with water, dried over MgSO₄,
5 filtered and concentrated. (4-methoxy-phenylsulfonyl)-acetic acid ethyl ester was isolated as pale yellow oil. Yield: 4.4 g (100%); MS: 227 (M+H)⁺.

To a stirred solution of 60% 3-chloroperoxybenzoic acid (14.0 gm, 40 mmol) in methylene chloride (100 ml) at 0° C, (4-methoxy-phenylsulfonyl)-acetic acid ethyl ester
10 (4.4 g, 20 mmol) in CH₂Cl₂ (15 ml) was added slowly. The reaction mixture turned cloudy and was stirred at room temperature for 6 hours. The reaction mixture was then diluted with hexanes (300 ml) and stirred for 15 minutes. The solids were filtered off and Na₂SO₃ solution was added to the organic layer which was stirred for at least 3 hours before the mixture was extracted with CHCl₃ and washed with H₂O. The
15 organic layer was dried over MgSO₄, filtered and concentrated and the colorless (4-methoxy-phenylsulfonyl)-acetic acid ethyl ester was isolated as an oil. Yield: 100%; MS: 259.1 (M+H)⁺.

To stirred solution of the (4-methoxy-benzenesulfonyl)-acetic acid ethyl ester (2.5 g, 10
20 mmol), benzyl bromide (1.8 gm, 10 mmol) and 18-Crown-6 (500 mg) in acetone (250 ml) was added K₂CO₃ (10gms, excess) and the mixture was refluxed for 24 hours. At the end, the reaction mixture was filtered and the acetone layer was concentrated. The residue obtained was extracted with chloroform, washed well with water, dried over anhydrous MgSO₄, filtered and concentrated. The product obtained was purified by
25 silica-gel column chromatography, eluting with 30% ethyl acetate: hexane. The product, 2-(4-methoxy-benzenesulfonyl)-3-phenyl-propionic acid ethyl ester was isolated as a low melting solid. Yield: 3.0 gm 86%; Low melting solid; MS: 349 (M+H)⁺.

30 To a stirred solution of 2-(4-methoxy-benzenesulfonyl)-3-phenyl-propionic acid ethyl ester (348 mg, 1 mmol) in methanol (25 ml), 10 N NaOH (10 ml) was added. The reaction mixture was stirred at room temperature for 48 hours. At the end, the reaction mixture was concentrated and carefully neutralized with dilute HCl. The residue obtained was extracted with chloroform, washed well with water, dried and
35 concentrated. The product obtained was purified by silica-gel column chromatography

WO 99/42436

PCT/US98/17633

- 31 -

by eluting with ethyl acetate: methanol (95:5) to afford 2-(4-methoxy-benzenesulfonyl)-3-phenyl-propionic acid as a colorless oil. Yield: 250 mg, 89%; MS: 321 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-3-phenyl-propionic acid (200 mg, 0.625 mmol) and following the procedure as outlined in Example 1, 150 mg of N-hydroxy-2-(4-methoxy-benzenesulfonyl)-3-phenyl-propionamide was isolated as a brown solid. Yield: 71%; mp 180 °C; MS: 336 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 3.2 (m, 1H), 3.8 (s, 3H), 4.0-4.2 (m, 2H), 7.0-8.0 (m, 9H).

10

Example 10

2-(4-Methoxy-benzenesulfonyl)-hexanoic acid hydroxyamide

2-(4-Methoxy-phenylsulfanyl)-hexanoic acid ethyl ester was prepared according to the general method as outlined in Example 1. Starting from ethyl 2-bromo hexanoate (7 g, 32 mmol) and 4-methoxybenzenethiol (4.2 g, 30 mmol), 8.3 g of the product was isolated. Yield 98%; Light yellow oil; MS: 283 (M+H)⁺.

Starting from 2-(4-methoxy-phenylsulfanyl)-hexanoic acid ethyl ester. (2.8 g 10 mmol) and following the procedure as outlined in Example 9, 3 g of 2-(4-methoxy-benzenesulfonyl)-hexanoic acid ethyl ester was isolated as a colorless solid. Yield: 95%; mp 62 °C; MS: 314 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-hexanoic acid ethyl ester (2 g, 6.3 mmol) 1.5 g (83%) of 2-(4-methoxy-benzenesulfonyl)-hexanoic acid was isolated as a colorless solid by following the procedure as outlined in Example 9. Mp 116 °C; MS: 287 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-hexanoic acid (1.0 g, 3.1 mmol) and following the procedure as outlined in Example 1, 700 mg of 2-(4-methoxy-benzenesulfonyl)-hexanoic acid hydroxyamide was isolated as a colorless solid. Yield: 60%; mp 130 °C; MS: 302 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 0.786 (t, J= 7.2 Hz, 3H), 1.1 - 1.3 (m, 4H), 1.6 - 1.8 (m, 2H), 3.7 (m, 1H), 3.9 (s, 3H), 7.2 (d, J = 11 Hz, 2H), 7.8 (d, J=11 Hz, 2H), 9.3 (s, 1H), 10.9 (s, 1H).

35

WO 99/42436

PCT/US98/17633

- 32 -

Example 11

2-(4-Methoxy-benzene sulfonyl)-tetradecanoic hydroxyamide

- 5 2-(4-Methoxy-phenylsulfanyl)-tetradecanoic acid ethyl ester was prepared according to the general method as outlined in Example 1. Starting from the corresponding ethyl 2-bromomyristate (5.0 g, 14.9 mmol) and 4-methoxythiophenol (1.9 g, 13.4 mmol), 5.0 g of the product was isolated. Yield 98%; Light yellow oil; MS: 393 (M+H)⁺.
- 10 Starting from 2-(4-methoxy-phenylsulfanyl)-tetradecanoic acid ethyl ester. (3.9 g 10 mmol) and following the procedure as outlined in Example 9, 3.2 g of 2-(4-methoxy-benzenesulfonyl)-tetradecanoic acid ethyl ester was isolated as a colorless solid. yield: 76%; Oil; MS: 425 (M+H)⁺.
- 15 Starting from 2-(4-methoxy-benzenesulfonyl)-tetradecanoic acid ethyl ester (2.5 g, 5.9 mmol), 2.0 g (85%) of 2-(4-methoxy-benzenesulfonyl)-tetradecanoic acid was isolated as a colorless solid by following the procedure as outlined in Example 9. mp 82 °C; MS: 397 (M+H)⁺.
- 20 Starting from 2-(4-methoxy-benzene sulfonyl)-tetradecanoic acid (1.14 g, 2.9 mmol) and following the procedure as outlined in Example 1, 670 mg of 2-(4-methoxy-benzenesulfonyl)-tetradecanoic hydroxyamide was isolated as an off-white solid. Yield: 57%; mp 114 °C; MS: 414 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.85 (t, J = 7 Hz, 3H), 1.16-1.27 (m, 20 H), 1.66 (m, 2H), 3.62-3.70 (m, 1H), 3.87 (s, 3H), 7.12
- 25 (d, J = 15 Hz, 2H), 7.73 (d, J = 15 Hz, 2H).

Example 12

- 30 N-Hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-3-phenyl-propionamide
- To a stirred solution of 2-(4-methoxy-benzenesulfonyl)-3-phenyl-propionic acid ethyl ester (1.0 gm, 3mmol) (example 9), methyl iodide (1 ml, excess) and 18-Crown-6 (500 mg) in acetone (250 ml), K₂CO₃ (10 gm, excess) was added and the reaction mixture was refluxed for 24 hours. At the end, the reaction mixture was filtered and
- 35 the acetone layer was concentrated. The residue obtained was extracted with chloroform, washed well with water, dried over anhydrous MgSO₄, filtered and concentrated. The product obtained was purified by silica-gel column chromatography by eluting it with 30% ethyl acetate:hexanes to afford 2-(4-methoxy-benzenesulfonyl)-

WO 99/42436

PCT/US98/17633

- 33 -

2-methyl-3-phenyl-propionic acid ethyl ester as a colorless oil. Yield 1.0 g, 98%; MS: 349 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-phenyl-propionic acid ethyl ester (900 mg, 2.7 mmol), 850 mg (quantitative) of 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-phenyl-propionic acid was isolated by following the procedure as outlined in Example 9. Colorless oil, MS 335 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-phenyl-propionic acid (900 mg, 2.7 mmol) and following the procedure as outlined in Example 1, 450 mg of N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-3-phenyl-propionamide was isolated as a brown solid. Yield: 48%; mp 58 °C; MS: 350 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.4 (s, 3H), 3.1 (d, J=9 Hz, 1H), 3.6 (d, J= 9 Hz, 1H), 3.9 (s, 3H), 6.8 - 7.8 (m, 9H).

15

Example 13

2-(4-Methoxy-benzenesulfonyl)-2,5-dimethyl-hex-4-enoic acid hydroxyamide

Starting from 2-(4-methoxy-phenylsulfanyl)-propionic acid ethyl ester (Example 1) (12 g; 50 mmol) and following the procedure as outlined in Example 9, 12 g of 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester was isolated as a semi-solid. yield 100%; MS: 256.1 (M+H)⁺.

Following the procedure as outlined in Example 12, 2-(4-methoxy-benzenesulfonyl)-2,5-dimethyl-hex-4-enoic acid ethyl ester was prepared, starting from (1 g, 3.6 mmol) of 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and isoprenyl bromide (1.0 g, 6 mmol). Yield 1.0 g, 81%; Colorless oil; MS: 341 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2,5-dimethyl-hex-4-enoic acid ethyl ester (900 mg, 2.6 mmol) 800 mg (96%) of 2-(4-methoxybenzenesulfonyl)-2,5-dimethyl-hex-4-enoic acid was isolated as a semi solid by following the procedure as outlined in Example 9. MS: 313 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2,5-dimethyl-hex-4-enoic acid (1.0 g, 3.2 mmol) and following the procedure as outlined in Example 1, 700 mg of 2-(4-methoxy-benzenesulfonyl)-2,5 dimethyl-hex-4-enoic acid hydroxyamide was isolated as a low melting solid. Yield: 67%; MS: 328 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ

WO 99/42436

PCT/US98/17633

- 34 -

1.3 (s, 3H), 1.5 (d, J=6.2 Hz, 6H), 2.5 -3.0 (m, 2H), 3.9 (s, 3H), 7.0 (d, J= 11 Hz, 2H), 7.8 (d, J= 11 Hz, 2H).

Example 14

5 3-(Biphenyl-4-yl)-N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionamide

Following the procedure as outlined in Example 12, 3-(biphenyl-4-yl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionic acid ethyl ester was prepared, starting from (2.7
10 g, 10 mmol) of 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and 4-(chloromethyl)biphenyl (2.5 g, 12 mmol). Yield 4.0 g, 91%; Colorless oil; MS: 438 (M+H)⁺.

Starting from 3-(biphenyl-4-yl)-2-(4-methoxy-benzenesulfonyl)-2-methyl propionic
15 acid ethyl ester (3 g, 6.8 mmol), 2.5 g (89%) of 3-(biphenyl-4-yl)-2-(4-methoxy-benzenesulfonyl)-2-methyl propionic acid was isolated as a colorless solid by following the procedure as outlined in Example 9. mp 161 °C; MS: 411 (M+H)⁺.

Starting from 3-(biphenyl-4-yl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionic
20 acid (2.0 g, 4.8 mmol) and following the procedure as outlined in Example 1, 1.2 g of 3-(biphenyl-4-yl)-N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionamide was isolated as colorless solid. Yield: 58%; mp 177 °C; MS: 426 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.4 (s, 3H), 3.2 (d, J=9 Hz, 1H), 3.7 (d, J= 9 Hz, 1H), 3.9 (s, 3H), 7.0 - 7.8 (m, 13H), 9.7 (bs, 1H).

25

Example 15

2-(4-methoxy-benzenesulfonyl)-2,5,9-trimethyl-deca-4,8-dienoic acid hydroxyamide

30 Following the procedure as outlined in Example 12, 2-(4-methoxy-benzenesulfonyl)-2,5,9-trimethyl-deca-4,8-dienoic acid ethyl ester was prepared, starting from (2.7 g, 10 mmol) of 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and geranyl bromide (3.0 g, 13 mmol). Yield 4.0 g, 98%; Colorless oil; MS: 409 (M+H)⁺.

35 Starting from 2-(4-methoxy-benzenesulfonyl)-2,5,9-trimethyl-deca-4,8-dienoic acid ethyl ester (3 g, 7.4 mmol), 2.8 g (96%) of 2-(4-methoxy-benzenesulfonyl)-2,5,9-trimethyl-deca-4,8-dienoic acid was isolated as a colorless oil by following the procedure as outlined in Example 9. MS: 379 (M-H)⁻.

WO 99/42436

PCT/US98/17633

- 35 -

Starting from 2-(4-methoxy-benzenesulfonyl)-2,5,9-trimethyl-deca-4,8-dienoic acid (2.0 g, 5.2 mmol) and following the procedure as outlined in Example 1, 1.8 g of 2-(4-methoxy-benzenesulfonyl)-2,5,9-trimethyl-deca-4,8-dienoic acid hydroxyamide was isolated as a colorless oil. Yield: 88%; MS: 396 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃):

5 δ 1.4 (s, 3H), 1.6 (s, 3H), 1.65 (s, 3H), 1.7 (s, 3H), 2.0-3.1 (m, 6 H), 3.9 (s, 3H), 5.5 (m, 2H), 6.98 (d, J= 9.0 Hz, 2H), 7.7 (d, J= 9.0 Hz, 2H).

Example 16

10 3-Cyclohexyl-N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionamide

Following the procedure as outlined in Example 12, 3-cyclohexyl-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionic acid ethyl ester was prepared, starting from (2.7 g, 10 mmol) of 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and bromo-

15 methylcyclohexane (1.8 g, 10 mmol). Yield 3.5 g, 95%; Yellow oil; MS: 369 (M+H)⁺.

Starting from 3-cyclohexyl-2-(4-methoxy-benzenesulfonyl)-2-methyl propionic acid ethyl ester (3 g, 8.1 mmol) 2.5 g (90%) of 3-cyclohexyl-2-(4-methoxy-benzenesulfonyl)-2-methyl propionic acid was isolated as colorless solid by following the

20 procedure as outlined in Example 9. mp 116 °C; MS: 341 (M+H)⁺.

Starting from 3-cyclohexyl-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionic acid (2.0 g, 5.8 mmol) and following the procedure as outlined in Example 1, 1.1 g of 3-cyclohexyl-N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionamide was

25 isolated as colorless solid. Yield: 55%; mp 58 °C; MS: 356 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.4 (s, 3H), 2.3 -1.0 (m, 13 H), 3.9 (s, 3H), 7.0 (d, 8.8 Hz, 2H), 7.69 (d, 9.0 Hz, 2H).

Example 17

30 N-Hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-propionamide

Following the procedure as outlined in example 12, 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid ethyl ester was prepared,

35 starting from (2.7 g, 10 mmol) of 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and the 4-(2-piperidin-1-yl-ethoxy)-benzyl chloride (2.9 g, 10 mmol). Yield 4.8 g, 98%; Brown oil; MS: 490 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 36 -

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid ethyl ester (4.0 gm, 7.9 mmol) 3.5 g (Yield: 94 %) of 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid was isolated as colorless crystals by following the procedure as outlined in example 9. Mp 106 °C; MS: 462.5 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid (2.0 g, 4.2 mmol) and following the procedure as outlined in example 1, 1 g of N-hydroxy- 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-propionamide was isolated as colorless solid. Yield: 1 g, 48%; mp 98°C; MS: 477 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.2 (s, 3H), 3.5 - 1.5 (m, 16 H), 3.9 (s, 3H), 4.4 (m, 1H), 6.5 - 7.8 (m, 8H), 10.8 (bs, 1H).

Example 18

2-[4-(2-Azepan-1-yl-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-propionic acid hydroxyamide

Following the procedure as outlined in example 12, 2-[4-(2-azepan-1-yl-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester was prepared, starting from (2.7 g, 10 mmol) of 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and the 1-[2-(4-chloromethyl-phenoxy)ethyl]-azepane (3.03 g, 10 mmol). Yield 4.5 g, 90%; Brown oil; MS: 504 (M+H)⁺.

Starting from 2-[4-(2-azepan-1-yl-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester (4.0 gm, 7.9 mmol) 3.5 g (Yield: 94 %) of 2-[4-(2-azepan-1-yl-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-propionic acid was isolated as semi-solid by following the procedure as outlined in example 9. MS: 476 (M+H)⁺.

Starting from 2-[4-(2-azepan-1-yl-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-propionic acid (2.0 g, 4.2 mmol) and following the procedure as outlined in example 1, 1 g of 2-[4-(2-azepan-1-yl-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-propionic acid hydroxyamide was isolated as colorless solid. Yield: 1.8 g, 87%; mp 68°C; MS: 491 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.23 (s, 3H), 3.5 - 1.7 (m, 18 H), 3.8 (s, 3H), 4.2 (m, 1H), 6.4 - 7.89 (m, 8H), 10.9 (bs, 1H).

WO 99/42436

PCT/US98/17633

- 37 -

Example 19

2-[4-(2-Azepan-1-yl-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-
pentanoic acid hydroxyamide

5 2-[4-(2-Azepan-1-yl-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-pentanoic acid ethyl ester was prepared according to the general method as outlined in example 12. Starting from 2-(4-methoxy-benzenesulfonyl)-pentanoic acid ethyl ester (3.5 g, 11.7 mmol) and 1-[2-(4-chloromethyl-phenoxy)-ethyl]-azepane (3.9 g, 12.8 mmol). Yield
10 2.58 g (42%); brown oil; MS: 532.4 (M+H)⁺.

2-[4-(2-Azepan-1-yl-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-pentanoic acid was prepared starting from 2-[4-(2-azepan-1-yl-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-pentanoic acid ethyl ester (2 g, 3.76 mmol) dissolved in methanol
15 (300 ml) and 10 N NaOH (15 ml). The resulting mixture was worked up as outlined in example 1. Yield 830 mg (44%); brown solid; mp 55 °C; MS: 504.4 (M+H)⁺.

Starting from 2-[4-(2-azepan-1-yl-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-pentanoic acid (690 mg, 1.37 mmol) and following the procedure as outlined in
20 example 1, 240 mg of 2-[4-(2-azepan-1-yl-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-pentanoic acid hydroxyamide was isolated as a yellow solid. Yield 34%; mp 85 °C; MS: 519.2 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.71 (t, J = 7.3 Hz, 3H), 0.78-1.77 (m, 16 H), 3.04-3.46 (m, 4H), 3.87 (s, 3H), 4.26 (m, 2H) 6.87 (d, J = 8.7 Hz, 2H), 7.14 (m, 4H), 7.71 (d, J = 9 Hz, 2H), 9.07 (s, 1H), 10 (s, 1H).

25

Example 20

N-Hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diisopropyl amino-ethoxy)-phenyl]-propionamide

30 Following the procedure as outlined in example 12, 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diisopropyl amino-ethoxy)-phenyl]-propionic acid ethyl ester was prepared, starting from (5.4 g, 20 mmol) of 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and the 4-(2-N,N-diisopropyl amino-ethoxy)-benzyl chloride (6.1 g, 20 mmol). Yield 8.9 g, 88%; Yellow oil; MS: 506.5 (M+H)⁺.

35

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diisopropyl amino-ethoxy)-phenyl]-propionic acid ethyl ester (4.0 gm, 7.9 mmol) 3.5 g (Yield: 92%) of 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diisopropyl amino-

WO 99/42436

PCT/US98/17633

- 38 -

ethoxy)-phenyl]-propionic acid was isolated as colorless crystals by following the procedure as outlined in example 9. Mp 68 °C; MS: 478.6 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diisopropyl amino-ethoxy)-phenyl]-propionic acid (2.0 g, 4.1 mmol) and following the procedure as outlined in example 1, 1 g of 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diisopropyl amino-ethoxy)-phenyl]-propionamide was isolated as colorless solid. Yield: 1 g, 49%; mp 98°C (HCl Salt); MS: 493 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.2 (s, 3H), 1.3 (d, 6H), 1.4 (d, 6H), 3.5 - 1.5 (m, 6 H), 3.9 (s, 3H), 4.4 (s, 2H), 6.5 - 7.8 (m, 8H), 10.8 (bs, 1H).

Example 21

N-Hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionamide

Following the procedure as outlined in example 12, 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionic acid ethyl ester was prepared, starting from (5.4 g, 20 mmol) of 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and the 4-(2-N,N-diethyl amino-ethoxy)-benzyl chloride (5.5 g, 20 mmol). Yield 8.5 g, 89%; Brown oil; MS: 478.6 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionic acid ethyl ester (3.5 gm, 7.7 mmol) 3.0 g (Yield: 85 %) of 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionic acid was isolated as colorless crystals by following the procedure as outlined in example 9. Mp 96-98 °C; MS: 450.5 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionic acid (2.0 g, 4.4 mmol) and following the procedure as outlined in example 1, 1 g of 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionamide was isolated as colorless solid. Yield: 1g, 48%; mp 56-59°C (HCl Salt); MS: 465.5 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.1 (t, 6H), 1.3 (s, 3H), 3.2 - 3.9 (m, 8 H), 3.9 (s, 3H), 4.3 (s, 2H), 6.5 - 7.8 (m, 8H), 10.8 (bs, 1H).

WO 99/42436

PCT/US98/17633

- 39 -

Example 22

N-Hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-piperidin-1-yl-ethoxy)-phenyl]-propionamide

5 Following the procedure as outlined in example 12, 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid ethyl ester was prepared, starting from (5.2 g, 20 mmol) of 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and the 3-(2-piperidin-1-yl-ethoxy)-benzyl chloride (6.0 g, 20 mmol). Yield 8.2 g, 83%; Brown oil; MS: 490 (M+H)⁺.

10

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid ethyl ester (6.0 gm, 12.2 mmol) 4.9 g (Yield: 79 %) of 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid was isolated as colorless crystals by following the procedure as outlined in

15 example 9. Mp 112 °C; MS: 462.5 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid (3.0 g, 6.5 mmol) and following the procedure as outlined in example 1, 1.8 g of 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-piperidin-1-yl-ethoxy)-phenyl]-propionamide was isolated as colorless solid. Yield: 1.8 g, 58%; mp 74°C; MS: 477 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.25 (s, 3H), 1.6-1.8 (m, 6 H), 2.5 - 3.7 (m, 8H), 3.9 (s, 3H), 4.4 (t, 2H), 6.7 - 7.8 (m, 8H), 10.8 (bs, 1H).

20

Example 23

25

3-(4-{3-[4-(3-Chloro-phenyl)-piperazin-1-yl]-propoxy}-phenyl)-N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionamide

Following the procedure as outlined in example 12, 3-(4-{3-[4-(3-chloro-phenyl)-piperazin-1-yl]-propoxy}-phenyl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionic acid ethyl ester was prepared, starting from (2.72 g, 10 mmol) of 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and the 1-[2-(4-chloromethyl-phenoxy)-ethyl]-4-(3-chloro-phenyl)-piperazine (4.2 g, 11 mmol). Yield 5.5 g, 89%; Brown oil; MS: 616 (M+H)⁺.

30

35 Starting from 3-(4-{3-[4-(3-chloro-phenyl)-piperazin-1-yl]-propoxy}-phenyl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionic acid ethyl ester (4.0 gm, 6.5 mmol) 3.0 g (Yield: 78 %) of 3-(4-{3-[4-(3-chloro-phenyl)-piperazin-1-yl]-propoxy}-phenyl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionic acid was isolated as colorless

WO 99/42436

PCT/US98/17633

- 40 -

crystals by following the procedure as outlined in example 9. Mp 196 °C; MS: 588.1 (M+H)⁺.

Starting from 3-(4-{3-[4-(3-chloro-phenyl)-piperazin-1-yl]-propoxy}-phenyl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionic acid (3.0 g, 5.1 mmol) and following the procedure as outlined in example 1, 1.8 g of 3-(4-{3-[4-(3-chloro-phenyl)-piperazin-1-yl]-propoxy}-phenyl)-N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionamide was isolated as pale yellow solid. Yield: 1.8 g, 55%; mp 122°C (HCl Salt); MS: 640 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.2 (s, 3H), 3.4 - 1.5 (m, 14 H), 3.9 (s, 3H), 4.5 (m, 2H), 6.5 - 8.2 (m, 12H), 10.3 (bs, 1H).

Example 24

2-(4-Methoxy-benzenesulfonyl)-5-methyl-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-hex-4-enoic acid hydroxyamide

To a stirred solution of (4-methoxy-benzenesulfonyl)-acetic acid ethyl ester (5.16 g, 20 mmol), isoprenyl bromide (3.0 g, 20 mmol) and 18-Crown-6 (500 mg) in acetone (250 ml) was added K₂CO₃ (10 gms, excess) and the mixture refluxed for 24 hours. At the end, the reaction mixture was filtered and the acetone layer was concentrated. The residue obtained was extracted with chloroform, washed well with water, dried over anhydrous MgSO₄, filtered and concentrated. The product obtained was purified by silica-gel column chromatography, eluting with 30% ethyl acetate: hexane. The product 2-(4-methoxy-benzenesulfonyl)-5-methyl-hex-4-enoic acid ethyl ester was isolated as a colourless oil. Yield: 3.0 g, 93%.

Following the procedure as outlined in example 12, 2-(4-Methoxy-benzenesulfonyl)-5-methyl-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-hex-4-enoic acid ethyl ester was prepared, starting from (3.26 g, 10 mmol) of 2-(4-methoxy-benzenesulfonyl)-5-methyl-hex-4-enoic acid ethyl ester and 4-(2-morpholin-1-yl-ethoxy)-benzyl chloride (3.0 g, 11 mmol). Yield 4.5 g, 82%; Brown oil; MS: 546 (M+H)⁺.

Starting from 2-(4-Methoxy-benzenesulfonyl)-5-methyl-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-hex-4-enoic acid ethyl ester (3.0 gm, 5.5 mmol) 2.1 g (Yield: 75 %) of 2-(4-Methoxy-benzenesulfonyl)-5-methyl-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-hex-4-enoic acid was isolated as semi-solid by following the procedure as outlined in example 9. MS: 518.6 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 41 -

Starting from 2-(4-Methoxy-benzenesulfonyl)-5-methyl-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-hex-4-enoic acid (1.0 g, 1.9 mmol) and following the procedure as outlined in example 1, 450 mg of 2-(4-Methoxy-benzenesulfonyl)-5-methyl-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-hex-4-enoic acid hydroxyamide was isolated as pale yellow solid. Yield: 450 mg, 45%; mp 92°C (HCl Salt); MS: 570 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.3 (d, 3H), 1.65 (d, 2H), 3.5 - 1.8 (m, 14 H), 3.9 (s, 3H), 4.5 (m, 2H), 5.4 (m, 1H), 6.5 - 7.9 (m, 8H), 11.5 (bs, 1H).

Example 25

10 N-Hydroxy-2-(4-ethoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionamide

To a stirred solution of 4-hydroxy thiophenol (12.6 g, 100 mmol) and triethyl amine (15.0 g, 150 mmol) in chloroform (400 ml) 2-bromo ethylpropionate (18.2 g, 100 mmol) was added drop wise. The reaction mixture was refluxed for 1 hr and cooled to room temperature. The reaction mixture was washed with water, dried and concentrated. 2-(4-hydroxy-phenylsulfanyl)-propionic acid ethyl ester was isolated as colorless oil. Yield: 22.0 g, 99%, MS: 227 (M+H).

20 To stirred solution of 2-(4-hydroxy-phenylsulfanyl)-propionic acid ethyl ester (11.3 g, 50 mmol), and K₂CO₃ (50 g, excess) in acetone (300 ml) ethyl iodide (20 ml, excess) was added and refluxed for 8 hrs. At the end, reaction mixture was filtered and concentrated. The residue obtained was extracted with chloroform and washed well with water. It was dried and concentrated. The product, 2-(4-ethoxy-phenylsulfanyl)-propionic acid ethyl ester was isolated as colorless oil. Yield: 12.0 g, 98%; MS: 255 (M+H).

2-(4-Ethoxy-phenylsulfanyl)-propionic acid ethyl ester was converted to 2-(4-ethoxy-phenylsulfonyl)-propionic acid ethyl ester by following the procedure as described in example 9, paragraph 2.

30 Following the procedure as outlined in example 12, 2-(4-ethoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionic acid ethyl ester was prepared, starting from (3.5 g, 12.2 mmol) of 2-(4-ethoxy-benzenesulfonyl)-propionic acid ethyl ester and the 4-(2-N,N-diethyl amino-ethoxy)-benzyl chloride (3.5 g, 12.2 mmol). Yield 4.8 g, 80%; Brown oil; MS: 492.6 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 42 -

Starting from 2-(4-ethoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionic acid ethyl ester (4.0 gm, 8.1 mmol) 3.2 g (Yield: 80 %) of 2-(4-ethoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionic acid was isolated as colorless semi-solid by following the procedure as outlined in example 9. MS: 464.5 (M+H)⁺.

Starting from 2-(4-ethoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionic acid (2.0 g, 4.3 mmol) and following the procedure as outlined in example 1, 1.2 g of 2-(4-ethoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionamide was isolated as colorless low melting solid. Yield: 1.2 g, 57%; (HCl Salt); MS: 478.5 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 0.9 (t, 3H), 1.1 (t, 6H), 1.3 (s, 3H), 3.2 - 3.9 (m, 8 H), 3.9 (s, 3H), 4.3 (s, 2H), 6.5 - 7.8 (m, 8H), 10.8 (bs, 1H).

15

Example 26

(4E)-2-(4-Methoxy-benzenesulfonyl)-5,9-dimethyl-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-deca-4,8-dienoic acid hydroxyamide

To a stirred solution of (4-methoxy-benzenesulfonyl)-acetic acid ethyl ester (5.16 g, 20 mmol), geranyl bromide (4.2g, 20 mmol) and 18-Crown-6 (500 mg) in acetone (250 ml) was added K₂CO₃ (10 gms, excess) and the mixture refluxed for 24 hours. At the end, the reaction mixture was filtered and the acetone layer was concentrated. The residue obtained was extracted with chloroform, washed well with water, dried over anhydrous MgSO₄, filtered and concentrated. The product obtained was purified by silica-gel column chromatography, eluting with 30% ethyl acetate: hexane. The product 2-(4-methoxy-benzenesulfonyl)-5,9-dimethyl-deca-4,8-dienoic acid ethyl ester was isolated as a colourless oil. Yield: 7.0 g, 89%.

Following the procedure as outlined in example 12, 2-(4-Methoxy-benzenesulfonyl)-5,9-dimethyl-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-deca-4,8-dienoic acid ethyl ester was prepared, starting from (1.0 g, 2.5 mmol) of 2-(4-methoxy-benzenesulfonyl)-5,9-dimethyl-deca-4,8-dienoic acid ethyl ester and 4-(2-morpholin-1-yl-ethoxy)-benzyl chloride (800 mg, 2.5 mmol). Yield 1.2 g, 76%; Brown oil; MS: 614 (M+H)⁺.

Starting from 2-(4-Methoxy-benzenesulfonyl)-5,9-dimethyl-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-deca-4,8-dienoic acid ethyl ester (2.0 gm, 3.2 mmol) 1.5 g (Yield: 80%) of 2-(4-Methoxy-benzenesulfonyl)-5,9-dimethyl-2-[4-(2-morpholin-4-yl-

WO 99/42436

PCT/US98/17633

- 43 -

ethoxy)-benzyl]-deca-4,8-dienoic acid was isolated as semi-solid by following the procedure as outlined in example 9. MS: 586.6 (M+H)⁺.

Starting from 2-(4-Methoxy-benzenesulfonyl)-5,9-dimethyl-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-deca-4,8-dienoic acid (1.0 g, 1.7 mmol) and following the procedure as outlined in example 1, 550 mg of (4E)-2-(4-Methoxy-benzenesulfonyl)-5,9-dimethyl-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-deca-4,8-dienoic acid hydroxyamide was isolated as pale yellow solid. Yield: 550 mg, 53%; mp 61°C (HCl Salt); MS: 638 (M+H)⁺.

10

Example 27

2-[4-(2-Diethylamino-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-hexanoic acid hydroxyamide

2-[4-(2-Diethylamino-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-hexanoic acid ethyl ester was prepared according to the general method as outlined in example 12. Starting from 2-(4-methoxy-benzenesulfonyl)-hexanoic acid ethyl ester (4 g, 12.7 mmol) and [2-(4-chloromethyl-phenoxy)-ethyl]-diethylamine (3.38 g, 14 mmol). Yield 8.21 g crude (100%); brown oil; MS: 520.4 (M+H)⁺.

20

2-[4-(2-Diethylamino-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-hexanoic acid was prepared starting from 2-[4-(2-diethylamino-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-hexanoic acid ethyl ester (8 g, 15.4 mmol) dissolved in methanol (200 ml) and 10 N NaOH (30 ml). The resulting mixture was worked up as outlined in example 1. Yield 3.88 g crude (51%); brown oil; MS: 492 (M+H)⁺.

25

Starting from 2-[4-(2-diethylamino-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-hexanoic acid (3.88 g, 7.89 mmol) and following the procedure as outlined in example 1, 800 mg of 2-[4-(2-diethylamino-ethoxy)-benzyl]-2-(4-methoxy-benzenesulfonyl)-hexanoic acid hydroxyamide was isolated as a yellow powder. Yield 20%; mp 67 °C; MS: 507.4 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.75 (t, J = 7.1 Hz, 3H), 1.05 (m, 2 H), 1.23 (t, J = 7.2 Hz, 6H) 1.37-1.91 (m, 2H), 3.13 (m, 4H), 3.38-3.51 (m, 4H), 3.87 (s, 3H), 4.3 (t, J = 4.8 Hz, 2H), 6.88 (d, J = 8.7 Hz, 2H), 7.15 (m, 4H), 7.7 (d, J = 9 Hz, 2H), 9.07 (s, 1H), 10.1 (s, 1H)

35

WO 99/42436

PCT/US98/17633

- 44 -

Example 28

N-Hydroxy-2-(4-n-butoxy-benzenesulfonyl)-2-methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-propionamide

5 Following the procedure as outlined in example 12, 2-(4-n-butoxy-benzenesulfonyl)-2-methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid ethyl ester was prepared, starting from (3.1 g, 10 mmol) of 2-(4-n-butoxy-benzenesulfonyl)-propionic acid ethyl ester (Prepared from 2-(4-hydroxy-phenylsulfonyl)-propionic acid ethyl ester and n-butylbromide following the procedure outlined in example 27)the 4-(2-piperidin-1-yl-ethoxy)-benzyl chloride (3.0 g, 10.1 mmol). Yield 4.5 g, 84%; Brown oil; MS: 532.7 (M+H)⁺.

Starting from 2-(4-n-butoxy-benzenesulfonyl)-2-methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid ethyl ester (5.0 gm, 9.4 mmol) 4.2 g (Yield: 88 %) of
15 2-(4-n-butoxy-benzenesulfonyl)-2-methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid was isolated as colorless solid by following the procedure as outlined in example 9. MS: 504.6 (M+H)⁺

Starting from 2-(4-n-butoxy-benzenesulfonyl)-2-methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid (3.0 g, 5.9 mmol) and following the procedure as outlined in example 1, 1.3 g of 2-(4-n-butoxy-benzenesulfonyl)-2-methyl-3-[4-(2-piperidine-1-yl-ethoxy)-phenyl]-propionamide was isolated as colorless solid. MP. 65°C, Yield: 1.3 g, 42%; (HCl Salt); MS: 478.5 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 0.9 (t, 3H), 1.2 (s, 3H), 1.3 - 1.9 (m, 10H), 2.8 - 4.5 (m, 12 H), 6.8 -
25 7.8 (m, 8H), 10.8 (bs, 1H).

Example 29

N-Hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionamide

Following the procedure as outlined in example 12, 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionic acid ethyl ester was prepared, starting from (5.0 g, 18 mmol) of 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and the 3-(2-N,N-diethyl amino-ethoxy)-benzyl chloride (4.9 g, 18
35 mmol). Yield 8.1 g, 93%; Brown oil; MS: 478.1 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionic acid ethyl ester (8.1 gm, 16.9 mmol) 6.7 g (Yield: 88 %) of

WO 99/42436

PCT/US98/17633

- 45 -

2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionic acid was isolated as colorless semi-solid by following the procedure as outlined in example 9. MP: 78-81; MS: 450.1 (M+H)⁺.

- 5 Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionic acid (6.7 g, 15 mmol) and following the procedure as outlined in example 1, 1.5 g of 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionamide was isolated as colorless low melting solid. Yield: 1.5 g, 21%; (HCl Salt); MS: 450.5 (M+H)⁺; ¹H NMR (300
- 10 MHz, DMSO-d₆): δ 1.21 (t, 6H), 1.26 (s, 3H), 3.18-3.24 (m, 2H), 3.38 (m, 4H), 3.43-3.46 (m, 2H), 3.80 (s, 3H), 4.30 (s, 2H), 6.76-6.78 (d, 2H), 6.84-7.2 (m, 6H), 10.3 (bs, 1H).

Example 30

15

N-Hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-morpholin-1-yl-ethoxy)-phenyl]-propionamide

- Following the procedure as outlined in example 12, 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-morpholin-1-yl-ethoxy)-phenyl]-propionic acid ethyl ester was
- 20 prepared, starting from (5.2 g, 20 mmol) of 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and the 3-(2-morpholin-1-yl-ethoxy)-benzyl chloride (6.0 g, 20 mmol). Yield 9.1 g, 93%; Brown oil; MS: 492 (M+H)⁺.

- 25 Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-morpholin-1-yl-ethoxy)-phenyl]-propionic acid ethyl ester (10.0 gm, 20.3 mmol) 8.0 g (Yield: 86 %) of 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-morpholin-1-yl-ethoxy)-phenyl]-propionic acid was isolated as colorless crystals by following the procedure as outlined in example 9.; MS: 464.5 (M+H)⁺.

30

- Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-morpholin-1-yl-ethoxy)-phenyl]-propionic acid (4.55 g, 9.8 mmol) and following the procedure as outlined in example 1, 440 mg of 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[3-(2-morpholin-1-yl-ethoxy)-phenyl]-propionamide was isolated as colorless solid. Yield:
- 35 440 mg, 9%; mp 63°C; MS: 479.5 (M+H)⁺; ¹H NMR (300 Mhz, DMSO-d₆): δ 1.26 (s, 3H), 3.18- 3.8 (m, 12H), 3.9 (s, 3H), 4.4 (m, 2H), 6.7 - 8.8 (m, 8H), 10.8 (bs, 1H).

WO 99/42436

PCT/US98/17633

- 46 -

Example 31

6-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-hexanoic acid hydroxyamide

5 Following the procedure as outlined in Example 9, 6-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-hexanoic acid ethyl ester was prepared, starting from (5.0 g, 20 mmol) of 2-(4-methoxy-benzenesulfonyl)-acetic acid ethyl ester and 4-phthalimido bromobutane (5.66 g, 20 mmol). Yield 8.4 g, 97%; Colorless
10 oil; MS: 474 (M+H).

Starting from 6-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-hexanoic acid ethyl ester (8.4 g, 17.7 mmol) 6.95 g (88%) of 6-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-
15 hexanoic acid was isolated as colorless oil by following the procedure as outlined in Example 9. MS: 446 (M-H).

Starting from 6-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-hexanoic acid (4.9 g, 11 mmol) and following the procedure as
20 outlined in Example 1, 3.1 g of 6-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-hexanoic acid hydroxyamide was isolated as a light brown solid; Yield: 46%; mp 146- 148 °C; MS: 461.2 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 1.55 (s, 3H), 1.61- 3.77 (m, 8H), 3.82 (s, 3H), 6.92-8.21 (m, 8H), 10.70 (bs, 1H), 11.20 (bs, 1H).

25

Example 32

3-[4-(2-Diethylamino-ethoxy)-phenyl]-2-(4-furan-2-yl-benzenesulfonyl)-N-hydroxy-2-methyl-propionamide

30 To a stirred solution of 4-bromo thiophenol (19.0 g, 100 mmol) and triethyl amine (15.0 g, 150 mmol) in chloroform (400 ml) 2-bromo ethylpropionate (18.2 g, 100 mmol) was added drop wise. The reaction mixture was refluxed for 1 hr and cooled to room temperature. The reaction mixture was washed with water, dried and concentrated. 2-(4-bromo-phenylsulfanyl)-propionic acid ethyl ester was isolated as
35 colorless oil. Yield: 28.0 g, 99%, MS: 290 (M+H).

WO 99/42436

PCT/US98/17633

- 47 -

2-(4-bromo-phenylsulfanyl)-propionic acid ethyl ester was converted to 2-(4-bromo-phenylsulfonyl)-propionic acid ethyl ester by following the procedure as described in example 9, paragraph 2.

- 5 A mixture of 2-(4-bromo-phenylsulfonyl)-propionic acid ethyl ester (6.4 g, 20 mmol), 2-(tributyl stannyl)furan (7.5g, 21 mmol) and (Ph₃P)₄Pd (500 mg) was refluxed in degassed toluene (250 ml) for 8 hrs. At the end reaction mixture was filtered through Celite and concentrated. The product was purified by silica gel column chromatography by eluting it with 50% ethylacetate : hexane. Colorless oil. Yield: 5.9 g, 95%,
10 MS: 309 (M+H).

- Following the procedure as outlined in example 12, 2-(4-(2-furanyl-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionic acid ethyl ester was prepared, starting from (3.08 g, 10.0 mmol) of 2-(4-(2-furanyl-benzenesulfonyl)-
15 propionic acid ethyl ester and the 4-(2-N,N-diethyl amino-ethoxy)-benzyl chloride (3.5 g, 12.2 mmol). Yield 5.0 g, 97%; Brown oil; MS: 514.6 (M+H)⁺.

- Starting from 2-(4-(2-furanyl-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionic acid ethyl ester (5.1 gm, 10.0 mmol) 3.8 g (Yield: 78 %) of
20 2-(4-(2-furanyl-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionic acid was isolated as colorless solid by following the procedure as outlined in example 9. MP: 58 °C, MS: 486.5 (M+H)⁺.

- Starting from 2-(4-(2-furanyl-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionic acid (5.0 g, 10.3 mmol) and following the procedure as outlined in example 1, 1.2 g of 2-(4-ethoxy-benzenesulfonyl)-2-methyl-3-[4-(2-N,N-diethyl amino-ethoxy)-phenyl]-propionamide was isolated as colorless low melting solid. Yield: 3.2 g, 62%; (HCl Salt); MS: 502 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃):
25 δ 1.23 (t, 6H), 1.4 (s, 2H), 2.8 (q, 4H), 3.0 (t, 2 H), 4.1 (t, 2H), 6.5 - 8.0 (m,
30 7H).

Example 33

- N-Hydroxy-2-(4-methoxy-benzenesulfonyl)-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-butyramide
35

2-(4-Methoxy-phenylsulfanyl)-butyric acid ethyl ester was prepared according to the general method as outlined in example 9. Starting from ethyl 2-bromobutyrate (10.71 g, 55 mmol) and 4-methoxythiophenol (7 g, 50 mmol), 5.19 g (40%); clear oil; MS: 255.2 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 48 -

2-(4-Methoxy-benzenesulfonyl)-butyric acid ethyl ester was prepared according to the general method as outlined in example 9. Starting from 2-(4-methoxy-phenylsulfanyl)-butyric acid ethyl ester (5 g, 20 mmol). Yield 5.74 g (100%); clear oil; MS: 287.1

5 (M+H)⁺.

Following the procedure as outlined in example 12, 2-(4-Methoxy-benzenesulfonyl)-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-butyric acid ethyl ester was prepared, starting from (3.5 g, 12.2 mmol) of 2-(4-methoxy-benzenesulfonyl)-butyric acid ethyl ester and
10 the 4-[2-(chloromethyl-phenoxy)-ethyl]-morpholine (2.34 g, 6.7 mmol). Yield 5.7 g, 100%; Brown oil; MS: 506.4 (M+H)⁺.

Starting from 2-(4-Methoxy-benzenesulfonyl)-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-butyric acid ethyl ester (5.54 gm, 11 mmol) 2.9 g (Yield: 55 %) of 2-(4-Methoxy-benzenesulfonyl)-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-butyric acid was isolated as
15 colorless semi-solid by following the procedure as outlined in example 9. MS: 478.3 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-butyric acid (2.6 g, 5.4 mmol) and following the procedure as outlined in example 1,
20 510 mg of N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-butyramide was isolated as a brown solid. Yield 2%; mp 51 °C; MS: 493.3 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.90 (t, J = 7.2 Hz, 3H), 1.69-1.96 (m, 4 H), 2.67 (t, 2H), 3.34 (m, 8H), 3.87 (s, 3H), 4.04 (m, 2H) 6.8 (d, J = 8.7
25 Hz, 2H), 7.14 (m, 4H), 7.73 (d, J = 4.7 Hz, 2H), 9.08 (s, 1H), 10.8 (s, 1H).

Example 34

30 N-Hydroxy-2-(4-methoxy-benzenesulfonyl)-2-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-butyramide

Following the procedure as outlined in example 12, 2-(4-Methoxy-benzenesulfonyl)-2-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-butyric acid ethyl ester was prepared, starting from (1.0 g, 3.33 mmol) of 2-(4-methoxy-benzenesulfonyl)-butyric acid ethyl ester and the 1-[2-(4-chloromethyl-phenoxy)-ethyl]-piperidine (0.85 g, 3.36 mmol). Yield 1.07
35 g, 62%; Brown oil; MS: 504.4 (M+H)⁺.

Starting from 2-(4-Methoxy-benzenesulfonyl)-2-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-butyric acid ethyl ester (3.7 gm, 7.3 mmol) 2.2 g (Yield: 63 %) of 2-(4-Methoxy-

WO 99/42436

PCT/US98/17633

- 49 -

benzenesulfonyl)-2-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-butyric acid was isolated as colorless semi-solid by following the procedure as outlined in example 9. MS: 476 (M+H)⁺.

- 5 Starting from 2-(4-Methoxy-benzenesulfonyl)-2-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-butyric acid (2.2 g, 4.63 mmol) and following the procedure as outlined in example 1, 360 mg of N-Hydroxy-2-(4-methoxy-benzenesulfonyl)-2-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-butyramide was isolated as a brown solid. Yield 16%; mp 75 °C; MS: 491.3 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.90 (t, J = 7.1 Hz, 3H), 1.36-1.96 (m, 4
10 H), 2.4-2.63 (m, 14H), 3.87 (s, 3H), 4.01 (t, J = 5.9 Hz, 2H) 6.8 (d, J = 8.5 Hz, 2H), 7.11 (m, 4H), 7.71 (d, J = 8.8 Hz, 2H), 9.09 (s, 1H), 10.8 (s, 1H)

Example 35

- 15 2-(4-Methoxy-benzenesulfonyl)-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-pentanoic acid hydroxyamide

- 2-(4-Methoxy-phenylsulfanyl)-pentanoic acid ethyl ester was prepared according to the general method as outlined in example 9. Starting from ethyl 2-bromovalerate (8.23 g, 39.3 mmol) and 4-methoxythiophenol (5 g, 35.7 mmol), 10.46 g (100%); clear oil;
20 MS: 269 (M+H)⁺.

- 2-(4-Methoxy-benzenesulfonyl)-pentanoic acid ethyl ester was prepared according to the general method as outlined in example 9. Starting from 2-(4-methoxy-phenyl-sulfanyl)-pentanoic acid ethyl ester (6.9 g, 27.4 mmol). Yield 7.07 g (86%); clear oil;
25 MS: 300.9 (M+H)⁺.

- Following the procedure as outlined in example 12, 2-(4-Methoxy-benzenesulfonyl)-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-pentanoic acid ethyl ester was prepared, starting from (3.0 g, 10.8 mmol) of 2-(4-methoxy-benzenesulfonyl)-pentanoic acid ethyl ester and the 4-[2-(chloromethyl-phenoxy)-ethyl]-morpholine (3.45 g, 11.9 mmol). Yield
30 3.08 g, 62%; Brown oil; MS: 520.4 (M+H)⁺.

- Starting from 2-(4-Methoxy-benzenesulfonyl)-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-pentanoic acid ethyl ester (2.73 gm, 5.27 mmol) 1.45 g (Yield: 56 %) of 2-(4-methoxy-benzenesulfonyl)-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-pentanoic acid was
35 isolated as colorless semi-solid by following the procedure as outlined in example 9. MS: 492.3 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 50 -

Starting from 2-(4-methoxy-benzenesulfonyl)-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-pentanoic acid (1.01 g, 2.05 mmol) and following the procedure as outlined in example 1, 190 mg of 2-(4-methoxy-benzenesulfonyl)-2-[4-(2-morpholin-4-yl-ethoxy)-benzyl]-pentanoic acid hydroxyamide was isolated as a brown solid. Yield 18%; mp 101 °C;

5 MS: 507.4 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.71 (t, J = 7 Hz, 3H), 1.58-1.82 (m, 4 H), 3.12-3.98 (m, 12H), 3.87 (s, 3H), 4.35 (t, 2H) 6.89 (d, J = 8.7 Hz, 2H), 7.15 (m, 4H), 7.74 (d, J = 8.9 Hz, 2H), 9.08 (s, 1H).

Example 36

10

2-[4-(2-Azepan-1-yl-ethoxy)-benzyl]-2-(4-Methoxy-benzenesulfonyl)-octanoic acid hydroxyamide

2-(4-Methoxy-phenylsulfanyl)-octanoic acid ethyl ester was prepared according to the general method as outlined in example 9. Starting from ethyl 2-bromooctanoate (11.8 g, 47.3 mmol) and 4-methoxythiophenol (6 g, 43 mmol). Yield: 7.24 g (57%); clear oil; MS: 311.2 (M+H)⁺,

15

2-(4-Methoxy-benzenesulfonyl)-octanoic acid ethyl ester was prepared according to the general method as outlined in example 9. Starting from 2-(4-methoxy-phenylsulfanyl)-octanoic acid ethyl ester (4.0 g, 13.6 mmol). Yield 3.7 g (83%); clear oil; MS: 343.3 (M+H)⁺.

20

Following the procedure as outlined in example 12, 2-[4-(2-Azepan-1-yl-ethoxy)-benzyl]-2-(4-Methoxy-benzenesulfonyl)-octanoic acid ethyl ester was prepared, starting from (1.69 g, 5.18 mmol) of 2-(4-methoxy-benzenesulfonyl)-octanoic acid ethyl ester and the 1-[2-(4-chloromethyl-phenoxy)-ethyl]-azepane (1.73 g, 6.0 mmol). Yield 4.86 g, 99%; Brown oil; MS: 574.5 (M+H)⁺.

25

Starting from 2-[4-(2-Azepan-1-yl-ethoxy)-benzyl]-2-(4-Methoxy-benzenesulfonyl)-octanoic acid ethyl ester (4.8 gm, 8.37 mmol) 1.55 g (Yield: 34 %) of 2-[4-(2-Azepan-1-yl-ethoxy)-benzyl]-2-(4-Methoxy-benzenesulfonyl)-octanoic acid was isolated as colorless semi-solid by following the procedure as outlined in example 9. MS: 551 (M+H)⁺.

30

Starting from 2-[4-(2-Azepan-1-yl-ethoxy)-benzyl]-2-(4-Methoxy-benzenesulfonyl)-octanoic acid (1.09 g, 2.0 mmol) and following the procedure as outlined in example 1, 300 mg of 2-[4-(2-Azepan-1-yl-ethoxy)-benzyl]-2-(4-Methoxy-benzenesulfonyl)-octanoic acid hydroxyamide was isolated as a yellow solid. Yield 27%; mp 65 °C; MS:

35

WO 99/42436

PCT/US98/17633

- 51 -

561.6 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.81 (t, J = 6.6 Hz, 3H), 1.08-1.82 (m, 14H), 3.13-3.51 (m, 12H), 3.87 (s, 3H), 4.33 (t, 2H), 6.88 (d, J = 8.7 Hz, 2H), 7.14 (m, 4H), 7.7 (d, J=9Hz, 2H), 9.06 (s, 1H), 10.28 (s, 1H).

5

Example 37

2-(4-Methoxy-benzenesulfanyl)-octanoic acid hydroxyamide

2-(4-Methoxy-phenylsulfanyl)-octanoic acid ethyl ester was prepared according to the general method as outlined in example 9. Starting from ethyl 2-bromooctanoate (11.8 g, 47.3 mmol) and 4-methoxythiophenol (6 g, 43 mmol). Yield: 7.24 g (57%); clear oil; MS: 311.2 (M+H)⁺.

Starting from 2-(4-Methoxy-benzenesulfanyl)-octanoic acid ethyl ester (3.1 gm, 10 mmol) 2.55 g (Yield: 90 %) of 2-(4-Methoxy-benzenesulfanyl)-octanoic acid was isolated as colorless semi-solid by following the procedure as outlined in example 9. MS: 283 (M+H)⁺.

Starting from 2-(4-Methoxy-benzenesulfanyl)-octanoic acid (4.25 g, 16 mmol) and following the procedure as outlined in example 1, 3.64 g of 2-(4-Methoxy-benzenesulfanyl)-octanoic acid hydroxyamide was isolated as colorless solid. Yield: 76%, MP: 90°C; MS: 298.2 (M+H).

25

Example 38

2-(4-Fluoro-phenylsulfanyl)-octanoic acid hydroxyamide

2-(4-Fluoro-phenylsulfanyl)-octanoic acid ethyl ester was prepared according to the general method as outlined in example 9. Starting from ethyl 2-bromooctanoate (6.47 g, 24.7 mmol) and 4-fluorothiophenol (3 g, 23.4 mmol). Yield: 6.31 g (90%); clear oil; MS: 299 (M+H)⁺.

Starting from 2-(4-fluoro-benzenesulfanyl)-octanoic acid ethyl ester (3.1 gm, 10 mmol) 2.89 g (Yield: 100 %) of 2-(4-fluoro-benzenesulfanyl)-octanoic acid was isolated as colorless semi-solid by following the procedure as outlined in example 9. MS: 268.9 (M+H)⁺.

Starting from 2-(4-fluoro-benzenesulfanyl)-octanoic acid (2.49 g, 9.2 mmol) and following the procedure as outlined in example 1, 2.72 g of 2-(4-fluoro-benzene-

WO 99/42436

PCT/US98/17633

- 52 -

sulfanyl)-octanoic acid hydroxyamide was isolated as colorless solid. Yield: 99%, MP: 58°C; MS: 284(M-H).

Example 39

5

2-(1-methyl-1H-imidazol-2-ylsulfanyl)-octanoic acid hydroxyamide

2-(1-methyl-1H-imidazol-2-ylsulfanyl)-octanoic acid ethyl ester was prepared according to the general method as outlined in example 9. Starting from ethyl 2-bromooctanoate (12.1 g, 48 mmol) and 1-methyl-2-mercapto imidazole (5 g, 43.8 mmol). Yield: 12 g (96%); clear oil; MS: 285 (M+H)⁺.

Starting from 2-(1-methyl-1H-imidazol-2-ylsulfanyl)-octanoic acid ethyl ester (12 gm, 42.2 mmol) 10.2 g (Yield: 95 %) of 2-(1-methyl-1H-imidazol-2-ylsulfanyl)-octanoic acid was isolated as colorless solid by following the procedure as outlined in example 9. MP: 95 °C, MS: 257.1 (M+H)⁺.

Starting from 2-(1-methyl-1H-imidazol-2-ylsulfanyl)-octanoic acid (7.84 g, 30.6 mmol) and following the procedure as outlined in example 1, 2.77 g of 2-(1-methyl-1H-imidazol-2-ylsulfanyl)-octanoic acid hydroxyamide was isolated as colorless solid. Yield: 33%, MP: 125 °C; MS: 272.2 (M+H).

Example 40

25 N-Hydroxy-2-(4-methoxy-benzenesulfonyl)-3-naphthalen-2-yl-propionamide

Following the procedure as outlined in Example 9, 2-(4-methoxy-benzenesulfonyl)-3-naphthalen-2-yl-propionic acid ethyl ester was prepared, starting from (5.0 g, 20 mmol) of 2-(4-methoxy-benzenesulfonyl)-acetic acid ethyl ester and 2-bromomethyl naphthalene (4.4 g, 20 mmol). Yield 7.2 g, 91%; Colorless oil; MS: 399 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-3-naphthalen-2-yl-propionamide ethyl ester (3.7 g, 9 mmol) 3.3g (96%) of 2-(4-methoxy-benzenesulfonyl)-3-naphthalen-2-yl-propionamide was isolated as colorless oil by following the procedure as outlined in Example 9. MS: 369.1 (M-H)⁻.

Starting from 2-(4-methoxy-benzenesulfonyl)-3-naphthalen-2-yl-propionic acid (2.2 g, 5.9 mmol) and following the procedure as outlined in Example 1, 820 mg of N-hydroxy-2-(4-methoxy-benzenesulfonyl)-3-naphthalen-2-yl-propionamide was isolated as a light brown solid; Yield: 36%; mp 161- 163 °C; MS: 385.9 (M+H)⁺; ¹H NMR

WO 99/42436

PCT/US98/17633

- 53 -

(300 MHz, CDCl_3): δ 3.32 (d, $J=7.0$ Hz, 1H), 3.69 (d, $J=7.0$ Hz, 1H), 3.82 (s, 3H), 5.02 (s, 1H), 6.92-7.89 (m, 11H).

Example 41

5

N-Hydroxy-2-(4-methoxy-phenylmethanesulfonyl)-2-methyl-3-phenyl propionic acid hydroxamide

10 A mixture of 4-methoxybenzyl mercaptan (7.0g, 45 mmol), ethyl 2-bromopropionate (8.2 g, 46 mmol) and powdered oven dried potassium carbonate (10g, 72 mmol) in 150 mL of acetone was heated at reflux for 18 h. The mixture was cooled, filtered, and the filtrate concentrated. The residue was taken up in 150 mL of methylene chloride, washed with water (150 mL), dried over anhydrous sodium sulfate and evaporated to yield 12 g (99%); colorless liquid; MS 255.1 (M+H). This product is used without
15 further purification.

To an ice cold (5 °C) solution of 2-(4-methoxy-phenylmethanesulfanyl)-propionic acid ethyl ester (5.7g, 21 mmol) in 100 mL CH_2Cl_2 was added portionwise (7.2g, 40 mmol) of m-chloroperbenzoic acid and the mixture was stirred for 1 h. The reaction
20 was diluted with hexanes (500 mL) and stirred at 25 °C for 30 minute at room temperature. The mixture was filtered and the organic layer treated with saturated aqueous sodium bisulfite (200 mL). The hexanes solution containing the product was washed with water, dried (Na_2SO_4) and concentrated. Yield 5.5g (91%); colorless oil; MS 287.1 (M+H)⁺.

25

Following the procedure as outlined in Example 9, 2-(4-Methoxy-phenylmethanesulfonyl)-2-methyl-3-phenyl-propionic acid ethyl ester was prepared, starting from 2-(4-Methoxy-phenylmethanesulfonyl)-propionic acid ethyl ester (2g, 7 mmol) and benzyl bromide (1.3g, 7.7 mmol). Yield 3.0 g, 100%; Low melting solid; MS: 377
30 (M+H)⁺.

2-(4-Methoxy-phenylmethanesulfonyl)-2-methyl-3-phenyl-propionic acid was prepared starting from 2-(4-Methoxy-phenylmethanesulfonyl)-2-methyl-3-phenyl-propionic acid ethyl ester (3.5 g, 9.0 mmol) dissolved in methanol (50 ml) and 10 N NaOH (30 ml).
35 The resulting reaction mixture was worked up as outlined in Example 9. Yield 930 mg, 31%. Colorless solid, mp: 106-108 °C; MS: 347 (M-H)⁺.

WO 99/42436

PCT/US98/17633

- 54 -

Starting from 2-(4-Methoxy-phenylmethanesulfonyl)-2-methyl-3-phenyl-propionic acid (2.7 g, 7.0 mmol) and following the procedure as outlined in example 1, 266 mg of N-Hydroxy-2-(4-methoxy-phenylmethanesulfonyl)-2-methyl-3-phenyl propionic acid hydroxamide was isolated as light colorless solid; Yield: 10%; mp 58-59 °C; MS: 364.2

5 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 1.28 (s, 3H), 2.84-2.88 (d, 1H), 3.75 (s, 3H), 3.81-3.86 (d, 1H), 4.59-4.63 (d, 1H), 4.69-4.74 (d, 1H), 6.94-6.98 (d, 2H), 7.19 (m, 2H), 7.29-7.33 (d, 4H), 9.24 (s, 1H), 10.88 (s, 1H).

Example 42

10

5-Methyl-2-(3-methyl-but-2-enyl)-2-(toluene-4-sulfonyl)-hex-4-enoic acid hydroxamide

15 5-Methyl-2-(3-methyl-but-2-enyl)-2-(toluene-4-sulfonyl)-hex-4-enoic acid ethyl ester was prepared according to general method as outlined in example 9. Starting from ethyl α-(p-tolylsulfonyl)acetate (2.9g, 10.9 mmol and 4-bromo-2-methyl butene (3.42g, 23 mmol). Yield 4.6g ; tan oil; MS 379.2 (M+H)⁺.

20 5-methyl-2-(3-methyl-but-2-enyl)-2-(toluene-4-sulfonyl)-hex-4-enoic acid was prepared according to general method as outlined in example 9. Starting from 5-methyl-2-(3-methyl-but-2-enyl)-2-(toluene-4-sulfonyl)-hex-4-enoic acid ethyl ester (4.5g, 11 mmol), ethanol (15 mL) and 10 N sodium hydroxide.

25 Starting from 5-methyl-2-(3-methyl-but-2-enyl)-2-(toluene-4-sulfonyl)-hex-4-enoic acid (4.1 g, 11 mmol) and following the procedure as outlined in example 1, 1.07 g of 5-Methyl-2-(3-methyl-but-2-enyl)-2-(toluene-4-sulfonyl)-hex-4-enoic acid hydroxy-amide was isolated as colorless solid; Yield: 30%; mp 108-110 °C; MS: 366.2 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 1.49 (s, 6H), 1.62 (s, 6H), 2.41 (s, 3H), 2.53-2.63 (m, 4H), 5.00-5.05 (t, 2H), 7.40-7.43 (d, 2H), 7.59-7.62 (d, 2H), 9.04 (s, 30 1H), 10.80 (s, 1H).

Example 43

35

2-Methyl-2-(2-methyl-furan-3-sulfonyl)-3-phenyl-propionic acid hydroxamide

2-Methyl-2-(2-methyl-furan-3-sulfonyl)-3-phenyl-propionic acid ethyl ester (Prepared from 3-mercapto-2-methylfuran) was prepared according to the general method as outlined in example 9. Starting from 2-(2-methyl-furan-3-ylsulfonyl)-propionic acid

WO 99/42436

PCT/US98/17633

- 55 -

ethyl ester (2.9g, 11.9 mmol), benzyl bromide (2.22g, 13 mmol) and potassium carbonate (10g) in acetone (75 mL). Yield (99 %); amber oil; MS 337.1 (M+H)⁺.

- 2-Methyl-2-(2-methyl-furan-3-sulfonyl)-3-phenyl-propionic acid was prepared according to the general method as outlined in example 9. Starting from 2-(2-methyl-furan-3-ylsulfonyl)-propionic acid ethyl ester (4.8g, 14.3 mmol), dissolved in ethanol (25 mL) and 10 N sodium hydroxide (10 mL). Yield 3.7g (84 %), white solid, MS 307.4 (M-H).
- Starting from 2-Methyl-2-(2-methyl-furan-3-sulfonyl)-3-phenyl-propionic acid (3.58 g, 12 mmol) and following the procedure as outlined in example 1, 1.078 g of 2-Methyl-2-(2-methyl-furan-3-sulfonyl)-3-phenyl-propionic acid hydroxyamide was isolated as orange color solid; Yield: 29%; mp 68-70 °C; MS: 324 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 1.27 (s, 3H), 2.81-2.86 (d, 1H), 3.33 (s, 3H), 3.61-3.66 (d, 1H), 6.66 (s, 1H), 7.19-7.25 (m, 5H), 7.76 (s, 1H), 9.09 (s, 1H), 10.81 (s, 1H)

Example 44

- 2-Methyl-2-(2-methyl-furan-3-sulfonyl)-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid hydroxamide

- 2-Methyl-2-(2-methyl-furan-3-sulfonyl)-3-[4-(2-piperidin-yl-ethoxy)-phenyl]-propionic acid ethyl ester was prepared according to the general method as outlined in example 9. Starting from 2-(2-methyl-furan-3-sulfonyl)-propionic acid ethyl ester (2.4g, 9.8 mmol) and 1-[2-(4-chloromethylphenoxy)-ethyl]-piperidine (2.96g, 10.7 mmol); Yield 2.4g (92%); amber oil; MS 464.2 (M+H)⁺.
- 2-Methyl-2-(2-methyl-furan-3-sulfonyl)-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid was prepared according to the general method as outlined in example 1. Starting from 2-methyl-2-(2-methyl-furan-3-sulfonyl)-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid ethyl ester (2.01g, 4.5 mmol), dissolved in ethanol (20 mL) and 10 N sodium hydroxide (10 mL). The resulting mixture was worked up as outline in example 9. Yield 2.03g; amber crystals mp 66-68 °C; MS 434 (M-H).
- Starting from 2-Methyl-2-(2-methyl-furan-3-sulfonyl)-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid (2.03 g, 6.0 mmol) and following the procedure as outlined in example 1, 1.36 g of 2-Methyl-2-(2-methyl-furan-3-sulfonyl)-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-propionic acid hydroxyamide was isolated as amber color solid; Yield:

WO 99/42436

PCT/US98/17633

- 56 -

32%; mp 115-117 °C; MS: 451.1 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 1.15-1.22 (m, 2H), (1.75 (s, 3H), 1.78 (s, 3H) 2.98-3.03 (m, 2H), 3.42-3.47 (m, 2H), 3.5 (s, 3H), 6.65 (s, 1H), 6.87-6.90 (d, 2H), 7.12-7.17 (d, 2H), 10.35 (s, 1H), 10.60 (s, 1H), 11.70 (s, 1H).

5

Example 45

2-Methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl-2-(thiophene-2-sulfonyl)-propionic acid hydroxamide

10

2-Methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl-2-(thiophene-2-sulfonyl)-propionic acid ethyl ester was prepared according to the general method as outlined in example 9. Starting from 2-(thiophene-2-sulfonyl)-propionic acid ethyl ester (prepared from 2-mercaptothiophene and 2-bromopropionic acid ethylester) (4.4g, 17.7 mmol) and 1-[2-(4-chloromethylphenoxy)-ethyl]-piperidine (5.3g, 19.5 mmol); Yield (96%); semi-solid; MS 466.

15

2-Methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl-2-(thiophene-2-sulfonyl)-propionic acid was prepared according to the general method as outlined in example 9. Starting from 2-methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl-2-sulfonyl)-propionic acid ethyl ester (9.8g, 20 mmol), dissolved in ethanol (20 mL) and 10 N sodium hydroxide (20 mL). The resulting mixture was worked up as outline in example 1. Yield 4.5g (49 %); white solid mp 170-172 °C; MS 436.3 (M-H).

20

25 Starting from 2-Methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl-2-(thiophene-2-sulfonyl)-propionic acid (3.6 g, 8.0 mmol) and following the procedure as outlined in example 1, 345 mg of 2-Methyl-3-[4-(2-piperidin-1-yl-ethoxy)-phenyl-2-(thiophene-2-sulfonyl)-propionic acid hydroxyamide was isolated as light colorless solid; Yield: 10%; mp 115-118 °C; MS: 451.2 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 1.29 (s, 3H), 1.66-1.78 (m, 6H), 2.81-2.86 (d, 1H), 2.96-3.99 (m, 4H), 3.39-3.47 (m, 2H), 3.51-3.59 (d, 1H), 4.32 (m, 2H), 6.72-6.74 (d 1H), 6.87-6.96 (d, 2H), 7.01-7.20 (m, 3H), 7.31-7.33 (m, 1H), 7.69-7.72 (m, 1H), 7.83-7.84 (m, 1H), 8.07-8.08 (dd, 1H), 8.17 (dd, 1H), 9.0 (s, 1H) 10.0 (s, 1H), 10.78 (s, 1H).

30

WO 99/42436

PCT/US98/17633

- 57 -

Example 46

2-(octane-1-sulfonyl)-3-[4-(2-piperidin-yl-ethoxy)-phenyl]propionic acid hydroxamide

5 2-(Octane-1-sulfonyl)-3-[4-(2-piperidin-yl-ethoxy)-phenyl]-propionic acid ethyl ester was prepared according to the general method as outlined in example 9. Starting from 2-(octane-1-sulfonyl)-propionic acid ethyl ester (5.0g, 18 mmol) and 1-[2-(4-chloromethylphenoxy)-ethyl]-piperidine (5.6g, 19.7 mmol); Yield 8.9g (96%); amber oil, MS 495.

10 2-(Octane-1-sulfonyl)-3-[4-(2-piperidin-yl-ethoxy)-phenyl]-propionic acid was prepared according to the general method as outlined in example 9. Starting from 2-(octane-1-sulfonyl)-3-[4-(2-piperidin-yl-ethoxy)-phenyl]-propionic acid ethyl ester (8.9g, 18 mmol), ethanol (25 mL) and 10 N sodium hydroxide (25 mL). Yield 6.0g
15 (72 %).

Starting from 2-(Octane-1-sulfonyl)-3-[4-(2-piperidin-yl-ethoxy)-phenyl]-propionic acid (3.6 g, 7.7 mmol) and following the procedure as outlined in example 1, 3.3 g of 2-(Octane-1-sulfonyl)-3-[4-(2-piperidin-yl-ethoxy)-phenyl]-propionic acid hydroxy-
20 amide was isolated as tan solid; Yield: 89%; mp 69-70 °C; MS: 483.2 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ .687 (t, 3H), 1.27-1.69 (m, 15H), 2.71-2.75 (d, 1H), 3.51 (s, 3H), 3.65-3.69 (d, 1H), 6.86-6.89 (d, 2H), 7.08-7.11 (d, 2H), 9.16 (s, 1H), 10.70 (s, 1H).

25 Example 47

3-Biphenyl-4-yl-2-methyl-2-(1-methyl-1H-imidazole-2-sulfonyl)-propionic acid hydroxyamide

30 3-Biphenyl-4-yl-2-methyl-2-(1-methyl-1H-imidazole-2-sulfonyl)-propionic acid ethyl ester was prepared according to the general method as outlined in example 9. Starting from 2-methyl-(1-methyl-1H-imidazolesulfonyl)-propionic acid ethyl ester Prepared from (1-Methyl-2-mercapto imidazole and 2-bromo ethyl propionate) (3.0g, 12.2 mmol) and 4-chloromethylbiphenyl (2.97g, 15 mmol). Yield 5.0g (99 %); low melting
35 solid; MS 413 (M+H)⁺.

3-Biphenyl-4-yl-2-methyl-2-(1-methyl-1H-imidazole-2-sulfonyl)-propionic acid was prepared according to the general method as outlined in example 9. Starting from 3-biphenyl-4-yl-2-methyl-2-(1-methyl-1H-imidazole-2-sulfonyl)-propionic acid ethyl ester

WO 99/42436

PCT/US98/17633

- 58 -

(5.0g, 11.9 mmol), ethanol (15 mL) and 10 N sodium hydroxide (10 mL). Yield 2.8g (61 %); brown solid mp 119-122 °C; MS 385.2 (M+H).

Starting from 3-Biphenyl-4-yl-2-methyl-2-(1-methyl-1H-imidazole-2-sulfonyl)-propionic acid (2.8 g, 7.0 mmol) and following the procedure as outlined in example 1, 112 mg of 3-Biphenyl-4-yl-2-methyl-2-(1-methyl-1H-imidazole-2-sulfonyl)-propionic acid hydroxyamide was isolated as tan colored solid; Yield: 4%; mp 112 °C; MS: 399.0 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.911 (s, 3H), 3.3 (s, 3H), 3.5 (d, 1H), 4.2 (d, 1H), 6.8 (d, 1H), 6.9 (d, 1H), 7.18-7.66 (m, 5H), 7.30-7.33 (d, 2H), 7.55-7.58 (d, 2H).

Example 48

2-Methyl-3-phenyl-2-(thiophene-2-sulfonyl)-propionic acid hydroxamide

2-Methyl-3-phenyl-2-(thiophene-2-sulfonyl)-propionic acid ethyl ester was prepared according to the general method as outlined in example 9. Starting from 2-(thiophen-2-sulfonyl)-propionic acid ethyl ester (3.0g, 12 mmol) and benzyl bromide (2.48g, 15 mmol). Yield 5.2 g (%); tan oil; MS 339.1 (M+H).

2-Methyl-3-phenyl-2-(thiophene-2-sulfonyl)-propionic acid was prepared according to the general method as outlined in example 9. Starting from 2-methyl-3-phenyl-2-(thiophen-2-sulfonyl)-propionic acid ethyl ester (5.0 g, 15 mmol), ethanol (30 mL) and 10 N sodium hydroxide (10 mL). Yield 5.6g MS 310.0 (M+H).

Starting from 2-Methyl-3-phenyl-2-(thiophene-2-sulfonyl)-propionic acid (5.0 g, 16 mmol) and following the procedure as outlined in example 1, 1.8 g of 2-Methyl-3-phenyl-2-(thiophene-2-sulfonyl)-propionic acid hydroxyamide was isolated as colorless solid; Yield: 40%; mp 116-117 °C; MS: 325.9 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 1.29 (s, 3H), 3.33 (d, 1H), 3.69 (d 1H), 7.18-7.30 (m, 5H), 7.74 (m, 1H), 8.22 (m, 1H), 9.13 (s, 1H), 10.80 (s, 1H).

Example 49

2-[8-(1-carboxy-ethanesulfonyl)-octane-1-sulfonyl]-propionic acid hydroxyamide

2-[8-(1-Carboxyl-ethanesulfonyl)-octane-1-sulfonyl]-propionic acid ethyl ester was prepared according to the general method as outlined in example 9. Starting from 2-[8-

WO 99/42436

PCT/US98/17633

- 59 -

- (1-ethoxycarbonyl-ethylsulfanyl)-octylsulfanyl]-propionic acid ethyl ester (10.2g, 26 mmol) and sodium peroxymonopersulfate (64g, 104 mmol). Yield 9.87g (86%); colorless liquid; MS 442.9 (M+H).
- 5 2-[8-(1-Carboxy-ethanesulfonyl)-octane-1-sulfonyl]-propionic acid was prepared according to general method as outline in example 1. Starting from 2-[8-(1-carboxy-ethanesulfonyl)-octane-1-sulfonyl]-propionic acid ethyl ester (3.0g, 6.8 mmol), ethanol (15 mL) and 10 N sodium hydroxide (15 mL). Yield 2.7g (98 %); white solid mp 99-102 °C; MS 387 (M+NH₃)⁺.
- 10 Starting from 2-[8-(1-Carboxy-ethanesulfonyl)-octane-1-sulfonyl]-propionic acid (2.5 g, 6.5 mmol) and following the procedure as outlined in example 1, 641 mg of 2-[8-(1-Carboxy-ethanesulfonyl)-octane-1-sulfonyl]-propionic acid hydroxyamidewas isolated as amber coloured oil.; Yield: 23%; MS: 434.0 (M+NH₄)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 1.27-3.23 (m, 22H), 3.33 (m, 2H), 8.9 (s, 1H), 9.28 (s, 1H).
- 15

Example 50

- 20 2-(4-Bromo-benzenesulfonyl)-2-methyl-3-[4-(2-piperidine-1-yl-ethoxy)-phenyl]-propionic acid hydroxamide
- 2-(4-Bromo-benzenesulfonyl)-2-methyl-3-[4-(2-piperidine-1-yl-ethoxy)-phenyl]-propionic acid ethyl ester was prepared according to general method as outlined in example 9. Starting from ethyl α-(4-bromophenyl-sulfonyl) acetate (5.0g, 16 mmol) and 1-[2-(4-chloromethylphenoxy)-ethyl]-piperidine (4.97g, 16 mmol). Yield 6.1g (71 %); tan oil; MS 541.1 (M+H)⁺.
- 25
- 2-(4-Bromo-benzenesulfonyl)-2-methyl-3-[4-(2-piperidine-1-yl-ethoxy)-phenyl]-propionic acid was prepared according to general method as outlined in example 9.
- 30 Starting from 2-(4-bromo-benzenesulfonyl)-2-methyl-3-[4-(2-piperidine-1-yl-ethoxy)-phenyl]-propionic acid ethyl ester (6.5g, 20 mmol), ethanol (30 mL) and 10 N sodium hydroxide (15 mL). Yield 6.3g (100 %); yellow solid mp 125-127 °C; MS 512.5 (M+H)⁺.
- 35 Starting from 2-(4-Bromo-benzenesulfonyl)-2-methyl-3-[4-(2-piperidine-1-yl-ethoxy)-phenyl]-propionic acid (6.1 g, 612 mmol) and following the procedure as outlined in example 1, 1.07 g of 2-(4-Bromo-benzenesulfonyl)-2-methyl-3-[4-(2-piperidine-1-yl-

WO 99/42436

PCT/US98/17633

- 60 -

ethoxy)-phenyl]-propionic acid hydroxyamide was isolated as light yellow solid; Yield: 17%; MS: 525.4 (M+H)⁺.

Example 51

5

3-(4-Bromo-phenyl)-N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionamide

Following the procedure as outlined in Example 9, 3-(4-bromo-phenyl)-2-(4-methoxy-benzensulfonyl)-2-methyl-propionic acid ethyl ester was prepared, starting from (3.0 g, 11 mmol) 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and 4-bromo-benzyl bromide (3.0 g, 12 mmol). Yield 4.67 g, 96%; Colorless oil; MS: 441 (M+H)⁺.

3-(4-Bromo-phenyl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionic acid was prepared starting from 3-(4-bromo-phenyl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionic acid ethyl ester (4.0 g, 9.0 mmol) dissolved in methanol (50 ml) and 10 N NaOH (30 ml). The resulting reaction mixture was worked up as outlined in Example 9. Yield 3.0 g, 78%. Low melting solid. MS: 413 (M+H)⁺.

Starting from 3-(4-bromo-phenyl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionic acid (2.7 g, 6.5 mmol) and following the procedure as outlined in example 1, 2.26 g of 3-(4-bromophenyl)-N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionamide was isolated as light colorless solid; Yield: 81%; mp 86-88 °C; MS: 429.8 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.42 (s, 3H), 1.77 (bs, 1H), 3.26 (d, J=7.0 Hz, 1H), 3.68 (d, J= 7.0 Hz, 1H), 3.85 (s, 3H), 7.01 -7.76 (m, 8H), 9.71 - 9.88 (bs, 1H).

Example 52

30 N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-3-naphthalen-2-yl-propionamide

Following the procedure as outlined in Example 9, 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-naphthalen-2-yl-propionic acid ethyl ester was prepared, starting from (5.4 g, 20 mmol) 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and 2-bromo-methyl naphthalene (4.4 g, 20 mmol). Yield 8.0 g, 97%; Colorless crystals, mp 182-184 °C; MS: 413 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 61 -

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-naphthalen-2-yl-propionic acid ethyl ester (4.6 g, 11 mmol) 4.2g (98%) of 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-naphthalen-2-yl-propionic acid was isolated as colorless crystals by following the procedure as outlined in Example 9. mp 144-146 °C; MS: 384.9 (M+H)⁺.

- 5 Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-naphthalen-2-yl-propionic acid (2.4 g, 6.2 mmol) and following the procedure as outlined in Example 1, 1.6 g of N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-3-naphthalen-2-yl-propionamide was isolated as a light colorless solid; Yield: 64%; mp 185 -187 °C; MS: 400.2
- 10 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.56 (s, 3H), 3.28 (d, J= 8.0 Hz, 1H), 3.81 (d, J=8Hz, 1H), 3.93 (s, 3H), 4.88 (bs, 1H), 7.02 - 7.92 (m, 11H).

Example 53

- 15 N-Hydroxy-2-(4-methoxy-benzenesulfonyl)-3-methyl-butyramide

2-(4-Methoxy-phenylsulfanyl)-3-methyl-butyric acid ethyl ester was prepared according to the general method as outlined in Example 1. Starting from ethyl 2-bromo-3-methyl-butanoate (20.9 g, 100 mmol) and 4-methoxybenzenethiol (14.0 g, 100

20 mmol), 30 g of 2-(4-methoxy-phenylsulfanyl)-3-methyl-butyric acid ethyl ester was isolated. Yield 99%; Light yellow oil; MS: 269 (M+H)⁺.

Starting from 2-(4-methoxy-phenylsulfanyl)-3-methyl-butyric acid ethyl ester. (2.68 g 10 mmol) and following the procedure as outlined in Example 9 for oxidation, 3 g of

25 2-(4-methoxy-benzenesulfonyl)-3-methyl-butyric acid ethyl ester was isolated as a colorless solid. yield: 99%; mp 53 °C; MS: 273 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-3-methyl-butyric acid ethyl ester (3 g, 10 mmol) 2.7 g (96%) of 2-(4-methoxy-benzenesulfonyl)-3-methyl-butyric acid was

30 isolated as a colorless solid by following the procedure as outlined in Example 9. Mp 96 °C; MS: 273 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-3-methyl-butyric acid (2.0 g, 7.34 mmol) and following the procedure as outlined in Example 9, 590 mg of N-hydroxy-2-

35 (4-methoxy-benzenesulfonyl)-3-methyl-butyramide was isolated as a colorless solid. Mp 220 °C; Yield 28%; MS: 288 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.88 (d,

WO 99/42436

PCT/US98/17633

- 62 -

J = 6.7 Hz, 3H), 1.07 (d, J = 6.7 Hz, 3H), 2.09-2.20 (bs, 1H), 3.53 (d, J = 9, 1H), 7.12-7.17 (m, 2H), 7.74-7.79 (m, 2H).

Example 54

5

1-(4-Methoxy-benzenesulfonyl)-cyclopentanecarboxylic acid hydroxyamide

Following the procedure as outlined in Example 9, 1-(4-methoxy-benzenesulfonyl)-cyclopentanecarboxylic acid ethyl ester was prepared, starting from (3.0 g, 11.6 mmol) of 2-(4-methoxy-benzenesulfonyl)-acetic acid ethyl ester and 1,4-dibromobutane (2.4 g, 7.6 mmol). Yield 2.4 g, 78%; Colorless solid, mp 86-88 °C; MS: 313 (M+H)⁺.

1-(4-Methoxy-benzenesulfonyl)-cyclopentanecarboxylic acid was prepared starting from 1-(4-methoxy-benzenesulfonyl)-cyclopentanecarboxylic acid ethyl ester (2.2 g, 7.0 mmol) dissolved in methanol (50 ml) and 10 N NaOH (30 ml). The resulting reaction mixture was worked up as outlined in Example 9. Yield 1.66 g, 83%.

Colorless solid; mp 112-115 °C; MS: 285 (M+H)⁺.

Starting from 1-(4-methoxy-benzenesulfonyl)-cyclopentanecarboxylic acid (442 mg, 1.5 mmol) and following the procedure as outlined in Example 1, 410 mg of 1-(4-methoxy-benzenesulfonyl)cyclopentanecarboxylic acid hydroxyamide was isolated as a colorless solid. mp 89-91 °C; Yield 88%; MS: 300 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.65-1.82 (m, 4H), 2.17-2.42 (m, 4H), 3.87 (s, 3H), 7.0 (d, J = 8Hz, 2H), 7.7 (bs, 1H), 7.72 (d, J=8 Hz, 2H), 9.73 (bs, 1H).

25

Example 55

3-(2-Bromo-phenyl)-N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionamide

30

Following the procedure as outlined in Example 9, 3-(2-bromo-phenyl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionic acid ethyl ester was prepared, starting from (2.0 g, 7.3 mmol) of 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and 2-(bromo)benzyl bromide (2.0 g, 8 mmol). Yield 3.1 g, 87%; Colorless oil; MS:

35 441 (M+H)⁺.

3-(2-Bromo-phenyl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionic acid was prepared starting from 3-(2-bromo-phenyl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-

WO 99/42436

PCT/US98/17633

- 63 -

propionic acid ethyl ester (3.0 g, 68 mmol) dissolved in methanol (50 ml) and 10 N NaOH (30 ml). The resulting reaction mixture was worked up as outlined in Example

9. Yield 1.7 g, 63%. Waxy solid; MS: 414 (M+H)⁺.

- 5 Starting from 3-(2-bromo-phenyl)-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionic acid (470 mg, 1.1 mmol) and following the procedure as outlined in Example 9, 380 mg of 3-(2-bromo-phenyl)-N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-propionamide was isolated as a colorless solid. mp 93-96 °C; Yield 77%; MS: 429 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.3 (s, 3H), 3.32 (d, J=7.0 Hz, 1H), 3.69 (d, J= 7.0 Hz, 1H), 3.82 (s, 3H), 6.92-7.89 (m, 8H).
- 10

Example 56

- 15 2-(4-methoxy-benzenesulfonyl)-2-methyl-5-phenyl-pent-4-enoic acid hydroxyamide
- Following the procedure as outlined in Example 9, 2-(4-methoxy-benzenesulfonyl)-2-methyl-5-phenyl-pent-4-enoic acid ethyl ester was prepared, starting from (3.0 g, 11 mmol) 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and cinnamyl bromide (2.1 g, 11 mmol). Yield 3.51 g, 82%; Colorless oil; MS: 389 (M+H)⁺.
- 20 2-(4-Methoxy-benzenesulfonyl)-2-methyl-5-phenyl-pent-4-enoic acid was prepared starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-5-phenyl-pent-4-enoic acid ethyl ester (3.0 g, 11 mmol) dissolved in methanol (50 ml) and 10 N NaOH (30 ml). The resulting reaction mixture was worked up as outlined in Example 9. Yield 1.9 g, 68%; yellowish oil; MS: 361 (M+H)⁺.
- 25

- Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-5-phenyl-pent-4-enoic acid (440 mg, 1.2 mmol) and following the procedure as outlined in Example 1, 420 mg of 2-(4-methoxy-benzenesulfonyl)-2-methyl-5-phenyl-pent-4-enoic acid hydroxyamide
- 30 was isolated as a colorless solid. mp 162-164 °C; Yield 92%; MS: 376 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.41 (s, 3H), 3.0-3.16 (m, 1H), 3.30 (d, J= 11 Hz, 2H), 3.92 (s, 3H), 5.9 - 6.1 (m, 1H), 6.53 (d, J=11Hz, 1H), 7.1-7.72 (m, 9H), 9.12 (bs, 1H).

WO 99/42436

PCT/US98/17633

- 64 -

Example 57

2-(4-methoxy-benzenesulfonyl)-5-phenyl-2-(3-phenyl-propyl)-pentanoic acid hydroxyamide

- 5 Following the procedure as outlined in Example 9, 2-(4-methoxy-benzenesulfonyl)-5-phenyl-2-(3-phenyl-propyl)-pentanoic acid ethyl ester was prepared, starting from (4.0 g, 15.8 mmol) 2-(4-methoxy-benzenesulfonyl)-acetic acid ethyl ester and 3-bromo-propyl benzene (6.4 g, 32 mmol). Yield 3.7 g, 47%; Colorless oil; MS: 495 (M+H)⁺.
- 10 2-(4-Methoxy-benzenesulfonyl)-5-phenyl-2-(3-phenyl-propyl)-pentanoic acid was prepared starting from 2-(4-methoxy-benzenesulfonyl)-5-phenyl-2-(3-phenyl-propyl)-pentanoic acid ethyl ester (2.0 g, 4 mmol) dissolved in methanol (50 ml) and 10 N NaOH (30 ml). The resulting reaction mixture was worked up as outlined in Example
- 15 9. Yield 1.18 g, 63%. Waxy solid; MS: 449.2 (M+H-H₂O)⁺.

- Starting from 2-(4-methoxy-benzenesulfonyl)-5-phenyl-2-(3-phenyl-propyl)-pentanoic acid (600 mg, 1.2 mmol) and following the procedure as outlined in Example 1, 420 mg of 2-(4-methoxy-benzenesulfonyl)-5-phenyl-2-(3-phenyl-propyl)-pentanoic acid
- 20 hydroxyamide was isolated as a colorless solid. Mp 118-120 °C; yield 68%; MS: 482 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.52 - 1.68 (m, 2H), 1.74 - 1.92 (m, 2H), 1.98-2.20 (m, 4H), 2.58 - 2.72 (m, 4H), 3.86 (s, 3H), 6.93 (d, J= 11 Hz, 2H), 7.02-7.63 (m, 10H), 7.81 (d, J=11 Hz, 2H).

- 25 Example 58

2-allyl-2-(4-methoxy-benzenesulfonyl)-pent-4-enoic acid hydroxyamide

- Following the procedure as outlined in Example 9, 2-allyl-2-(4-methoxy-benzene-sulfonyl)-pent-4-enoic acid ethyl ester was prepared, starting from (3.0 g, 11.6 mmol)
- 30 2-(4-methoxy-benzenesulfonyl)-acetic acid ethyl ester and allyl bromide (4 ml, excess). Yield 3.6 g, 92%; Yellow oil; MS: 338 (M+H)⁺.

- 2-Allyl-2-(4-methoxy-benzenesulfonyl)-pent-4-enoic acid was prepared starting from
- 35 2-allyl-2-(4-methoxy-benzenesulfonyl)-pent-4-enoic acid ethyl ester (2.2 g, 6.5 mmol) dissolved in methanol (50 ml) and 10 N NaOH (30 ml). The resulting reaction mixture was worked up as outlined in Example 9. Yield 1.76 g, 87%; yellowish oil; MS: 311 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 65 -

Starting from 2-allyl-2-(4-methoxy-benzenesulfonyl)-pent-4-enoic acid (1.5 g, 4.8 mmol) and following the procedure as outlined in Example 1, 1.5 g of 2-allyl-2-(4-methoxy-benzenesulfonyl)-pent-4-enoic acid hydroxyamide was isolated as colorless solid. Mp 114-116 °C; Yield 99%; MS: 326 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃):

5 δ 1.62 (s, 1H), 2.70 - 2.80 (m, 4H), 3.9 (s, 3H), 5.16 - 5.27 (m, 4H), 5.81-5.94 (m, 2H), 7.12 (d, J=8 Hz, 2H).

Example 59

10

2-(4-methoxy-benzenesulfonyl)-2-propyl-pentanoic acid hydroxyamide

2-allyl-2-(4-methoxy-benzenesulfonyl)-pent-4-enoic acid hydroxyamide (326 mg, 1.0 mmol) (example 26) was dissolved in methanol (50 ml) and hydrogenated over 10% Pd/C (100 mg) at room temperature, under 49 psi pressure for 4 hours. At the end, the

15 reaction mixture was filtered and methanol was removed. The resulting solid was crystallized from methanol. Yield: 250 mg, 75%; MS: 330 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 0.92 (t, J = 4.0 Hz, 6H), 1.27-1.59 (m, 4H), 1.78-2.02 (m, 4H), 3.86 (s, 3H), 6.04 (bs, 1H), 6.97 (d, J=9Hz, 2H), 7.76 (d, J=9 Hz, 2H).

20

Example 60

2-benzyl-N-hydroxy-2-(4-methoxy-benzenesulfonyl)-3-phenyl-propionamide

25 Following the procedure as outlined in Example 9, 2-benzyl-2-(4-methoxy-benzenesulfonyl)-3-phenyl-propionic acid ethyl ester was prepared, starting from (1.0 g, 3.8 mmol) of 2-(4-methoxy-benzenesulfonyl)-acetic acid ethyl ester and benzylbromide (4 ml, excess). Yield 1.2 g, 72%; Yellow oil; MS: 439 (M+H)⁺.

30 2-Benzyl-2-(4-methoxy-benzenesulfonyl)-3-phenyl-propionic acid was prepared starting from 2-benzyl-2-(4-methoxy-benzenesulfonyl)-3-phenyl-propionic acid ethyl ester (1.0 g, 2.2 mmol) dissolved in methanol (50 ml) and 10 N NaOH (30 ml). The resulting reaction mixture was worked up as outlined in Example 9. Yield: 580 mg, 62%; Waxy solid; MS: 409 (M-H)⁻.

35

Starting from 2-benzyl-2-(4-methoxy-benzenesulfonyl)-3-phenyl-propionic acid (410 mg, 1 mmol) and following the procedure as outlined in Example 1, 225 mg of 2-benzyl-N-hydroxy-2-(4-methoxy-benzenesulfonyl)-3-phenyl-propionamide was

WO 99/42436

PCT/US98/17633

- 66 -

isolated as a waxy solid. Yield 52%; MS: 426 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃):
δ 3.25 (d, J=14 Hz, 2H), 3.52 (d, J= 14 Hz, 2H), 3.9 (s, 3H), 6.93 (d, J=8Hz, 2H),
7.02 - 7.26 (m, 9H), 7.61 (d, J=8Hz, 2H), 7.87 (d, J=4Hz, 1H), 9.58 (bs, 1H).

5

Example 61

N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-3-pyridin-3-yl-propionamide

To a stirred solution of 2-(4-methoxy-benzenesulfonyl)propionic acid ethyl ester (2.7
10 gm, 10 mmol), 3-picolyl chloride hydrochloride (3.2 g, 20 mmol). and triethyl benzyl-
ammonium chloride (1 g) in methylene chloride (400 ml), 10 N NaOH (30 ml) was
added. The reaction was continued at room temp for 48 hours. At the end, the organic
layer was separated and washed well with water. The organic layer was dried, filtered
and concentrated. The crude product obtained was purified by silica-gel column
15 chromatography. The column was eluted with 50% ethyl acetate: hexane. 2-(4-
Methoxy-benzenesulfonyl)-2-methyl-3-pyridin-3-yl-propionic acid ethyl ester was
isolated as brown oil. Yield 3.0 g, 82%; Brown oil; MS: 364 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-pyridin-3-yl-propionic acid
20 ethyl ester (2.5 g, 6.8 mmol) 1.8 g (79%) of 2-(4-methoxy-benzenesulfonyl)-2-methyl-
3-pyridin-3-yl-propionic acid was isolated as a colorless solid by following the
procedure as outlined in Example 9. mp 58 °C; MS: 336 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-pyridin-3-yl-propionic acid
25 (410 mg, 1 mmol) and following the procedure as outlined in Example 1, 225 mg of
N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-3-pyridin-3-yl-propionamide
was isolated as a colorless solid. Yield 52%; mp 98 °C; MS: 351 (M+H)⁺; ¹H NMR
(300 MHz, CDCl₃): δ 1.4 (s, 3H), 3.1 (d, J=9.0, 1H), 3.65 (d, J= 9.1, 1H), 3.9 (s,
3H), 7-8.5 (m, 8H).

30

Example 62

2-(4-Methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-decanoic acid hydroxyamide

35 Starting from 2-(4-methoxy-benzenesulfonyl)-acetic acid ethyl ester (7.5 g, 29 mmol)
and 1-bromooctane (6.7 g, 35 mmol) 8 g of the mono octylated compound 2-(4-

WO 99/42436

PCT/US98/17633

- 67 -

methoxy-benzenesulfonyl)-decanoic acid ethyl ester was isolated by following the procedure outlined in Example 9. Yield: 8.0 g 74%; MS: 370 (M+H)⁺.

- Following the procedure as outlined in example 29, 2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-decanoic acid ethyl ester was prepared, starting from (8.0 g, 21.6 mmol) of 2-(4-methoxy-benzenesulfonyl)-decanoic acid ethyl ester and 3-picolyl chloride hydrochloride (4.1 g, 25 mmol). Yield 6.5 g, 68%; Brown oil; MS: 462 (M+H)⁺.
- 10 Starting from 2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-decanoic acid ethyl ester (5.0 g, 11 mmol), 4.5g (91%) of 2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-decanoic acid was isolated as a colorless solid by following the procedure as outlined in Example 9. Mp 159 °C; MS: 434 (M+H)⁺.
- 15 Starting from 2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-decanoic acid (2.5 g, 5.7 mmol) and following the procedure as outlined in Example 1, 1.4 g of 2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-decanoic acid hydroxyamide was isolated as colorless solid. Yield: 50%; mp 62 °C; MS: 448 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 0.86 (t, 6.9 Hz, 3H), 1.25-2.17 (m, 14 H), 3.3 (d, J=14 Hz, 1H), 3.5 (d, J= 14 Hz, 1H), 3.9 (s, 3H), 6.8 - 8.6 (m, 8H).
- 20

Example 63

- 25 2-(4-Methoxy-benzenesulfonyl)-5-methyl-2-pyridin-3-ylmethyl-hex-4-enoic acid hydroxyamide

- Following the procedure as outlined in Example 9, 2-(4-methoxy-benzenesulfonyl)-5-methyl-hex-4-enoic acid ethyl ester was prepared, starting from (6.0 g, 23 mmol) 2-(4-methoxy-benzenesulfonyl)-acetic acid ethyl ester and isoprenyl bromide (3.0 g, 20 mmol). Yield 6.52 g, 86%; Colorless oil; MS: 327 (M+H)⁺.
- 30

- Following the procedure as outlined in Example 29, 2-(4-methoxy-benzenesulfonyl)-5-methyl-2-pyridin-3-ylmethyl-hex-4-enoic acid ethyl ester was prepared, starting from (4.0 g, 12.2 mmol) of 2-(4-methoxy-benzenesulfonyl)-5-methyl-hex-4-enoic acid ethyl ester and 3-picolylchloride hydrochloride (2.1 g, 13 mmol). Yield 4.14 g, 81%; Brown oil; MS: 418 (M+H)⁺.
- 35

WO 99/42436

PCT/US98/17633

- 68 -

2-(4-Methoxy-benzenesulfonyl)-5-methyl-2-pyridin-3-ylmethyl-hex-4-enoic acid was prepared starting from 2-(4-methoxy-benzenesulfonyl)-5-methyl-2-pyridin-3-ylmethyl-hex-4-enoic acid ethyl ester (4.0 g, 9.5 mmol) dissolved in methanol (50 ml) and 10 N NaOH (30 ml). The resulting reaction mixture was worked up as outlined in Example

5 9. Yield 3.2 g, 87%; ivory solid; mp 117-119 °C; MS: 390 (M+H)+.

Starting from 2-(4-methoxy-benzenesulfonyl)-5-methyl-2-pyridin-3-ylmethyl-hex-4-enoic acid (2.1 g, 5.4 mmol) and following the procedure as outlined in Example 1, 1.82 g of 2-(4-methoxy-benzenesulfonyl)-5-methyl-2-pyridin-3-ylmethyl-hex-4-enoic
10 acid hydroxyamide was isolated as a colorless solid. Yield: 82%; mp 89 - 92 °C; MS: 405 (M+H)+; ¹H NMR (300 MHz, CDCl₃): δ 1.63 (s, 3H), 1.76 (s, 3H), 2.62-2.78 (m, 2H), 3.3 (d, J=4.0 Hz, 1H), 3.63 (d, J= 4.0 Hz, 1H), 3.82 (s, 3H), 5.26 (m, 1H), 7.12-7.88 (m, 6H), 8.27-8.33 (m, 2H).

15 Example 64

2-Benzyl-4-diisopropylamino-N-hydroxy-2-(4-methoxy-benzenesulfonyl)-butyramide

Following the procedure as outlined in Example 29, 2-benzyl-4-diisopropylamino-2-(4-methoxy-benzenesulfonyl)-butyric acid ethyl ester was prepared, starting from (3.0 g,
20 8.5 mmol) of 2-(4-methoxy-benzenesulfonyl)-3-phenyl-propionic acid ethyl ester (Example 9) and 2-diisopropylaminoethyl chloride hydrochloride (4.0 g, 20 mmol). Yield 3.2 g, 79%; Ivory solid, mp 89-91 °C; MS: 476.4 (M+H)+.

25 Starting from 2-benzyl-4-diisopropylamino-2-(4-methoxy-benzenesulfonyl)-butyric acid ethyl ester (3.53 gm, 7.5 mmol) 2.8 g (86%) of 2-benzyl-4-diisopropylamino-2-(4-methoxy-benzenesulfonyl)-butyric acid was isolated as colorless crystals by following the procedure as outlined in Example 9. Mp 136-138 °C; MS: 448.5 (M+H)+.

30 Starting from 2-benzyl-4-diisopropylamino-2-(4-methoxy-benzenesulfonyl)-butyric acid (1.85 g, 4.1 mmol) and following the procedure as outlined in Example 1, 1.3 g of 2-benzyl-4-diisopropylamino-N-hydroxy-2-(4-methoxy-benzenesulfonyl)-butyramide was isolated as a low melting waxy solid; Yield: 68%; MS: 463.3 (M+H)+; ¹H NMR
35 (300 MHz, CDCl₃): δ 0.98 (d, J = 11 Hz, 6H), 1.16 (d, J=11 Hz, 6H), 1.92 (m, 2H), 2.46 (m, 2H), 2.71 (m, 2H), 3.18 (m, 1H), 3.48 (m, 1H), 3.86 (s, 3H), 6.98 (d, J=8 Hz, 2H), 7.18 -7.22 (m, 5H), 7.92 (d, J=8 Hz, 2H), 8.12 (s, 1H).

WO 99/42436

PCT/US98/17633

- 69 -

Example 65

3-Cyclohexyl-N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-propionamide

5 Following the procedure as outlined in Example 9, 3-cyclohexyl-2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester was prepared, starting from (4.0 g, 15 mmol) 2-(4-methoxy-benzenesulfonyl)-acetic acid ethyl ester and 1-bromomethyl cyclohexane (2.7 g, 15 mmol). Yield 5.0 g, 94%; Colorless oil; MS: 355 (M+H)⁺.

10 Following the procedure as outlined in Example 29, 3-cyclohexyl-2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-propionic acid ethyl ester was prepared, starting from 3-cyclohexyl-2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester (1.5 g, 4.2 mmol) and 3-picolyl chloride (1.0 g, 6 mmol). Yield 1.0 g, 38%; Colorless oil; MS 446 (M+H)⁺.

15 Starting from 3-cyclohexyl-2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-propionic acid ethyl ester (1.3 g, 2.9 mmol) 1.0 g (83%) of 3-cyclohexyl-2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-propionic acid was isolated as colorless crystals by following the procedure as outlined in Example 9. Mp 92 °C; MS: 417.5 (M+H)⁺

20 Starting from 3-cyclohexyl-2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-propionic acid (1.0 g, 2.4 mmol) and following the procedure as outlined in Example 1, 80 mg of 3-cyclohexyl-N-hydroxy-2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-propionamide was isolated as a colorless hydrochloride salt; Yield: 71%; mp 57-60 °C; MS: 433 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 0.8-2.08 (m, 13 H), 3.3 (d, J=14 Hz, 1H), 3.7 (d, J= 14 Hz, 1H), 3.9 (s, 3H), 7.0 - 8.5 (m, 8H).

30 Example 66

2-(4-Methoxy-benzenesulfonyl)-4-methyl-2-pyridin-3-ylmethyl-pentanoic acid hydroxyamide

35 Following the procedure as outlined in Example 9, 2-(4-methoxy-benzenesulfonyl)-4-methyl-pentanoic acid ethyl ester was prepared, starting from (5.0 g, 20 mmol) 2-(4-methoxy-benzenesulfonyl)-acetic acid ethyl ester and 1-bromo-2-methyl propane (2.6 g, 20 mmol). Yield 6.0 g, 95%; Colorless oil; MS: 315 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 70 -

Following the procedure as outlined in Example 29, 2-(4-methoxy-benzenesulfonyl)-4-methyl-2-pyridin-3-ylmethyl-pentanoic acid ethyl ester was prepared, starting from (3.1 g, 10 mmol) of 2-[(4-methoxy-benzenesulfonyl)-4-methyl pentanoic acid ethyl ester and 3-picoyl chloride hydrochloride (1.8 g, 11 mmol). Yield 3.0 g, 75%; Colorless oil; MS: 406 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-4-methyl-2-pyridin-3-ylmethyl-pentanoic acid ethyl ester (1.2 g, 2.9 mmol) 1.0g (91%) of 2-(4-methoxy-benzenesulfonyl)-4-methyl-2-pyridin-3-ylmethyl-pentanoic acid was isolated as colorless crystals by following the procedure as outlined in Example 9. Mp 188-186 °C; MS: 378 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-4-methyl-2-pyridin-3-ylmethyl-pentanoic acid (800 mg, 2.1 mmol) and following the procedure as outlined in Example 1, 180 mg of 2-(4-methoxy-benzenesulfonyl)-4-methyl-2-pyridin-3-ylmethyl-pentanoic acid hydroxyamide was isolated as a colorless solid; Yield: 21%; mp 78 °C; MS: 393.4 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 0.65 (d, 6.3 Hz, 3H), 0.89 (d, J=6.2 Hz, 3H), 1.7 (m, 1H), 2.06 (m, 2H), 3.85 (s, 3H), 6.8 -8.5 (m, 10H).

Example 67

N-Hydroxy-2-(4-methoxy-benzenesulfonyl)-2-methyl-3-quinolin-6-yl-propionamide

Following the procedure as outlined in Example 29, 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-quinolin-6-yl-propionic acid ethyl ester was prepared, starting from (5.2 g, 20 mmol) of 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and 7-bromomethyl quinoline (4.4 g, 20 mmol). Yield 4.5 g, 54%; Pale yellow solid; mp 86°C; MS: 414 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-quinolin-6-yl-propionic acid ethyl ester (3.0 gm, 7.2 mmol) 2.5g (90%) of 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-quinolin-6-yl-propionic acid was isolated as colorless crystals by following the procedure as outlined in Example 9. mp 106-108 °C; MS: 386.4 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-quinolin-6-yl-propionic acid (2.0 gm, 5.2 mmol) and following the procedure as outlined in Example 1, 1.2 g of N-hydroxy-2-(methoxy-benzenesulfonyl)-2-methyl-3-quinolin-6-yl-propionamide was isolated as a colorless solid; Yield: 57%; mp 206 °C; MS: 401.4 (M+H)⁺; ¹H NMR

PCT/US98/17633

(300 MHz, CDCl₃): δ 1.4 (s, 3H), 3.19 (m, 1H), 3.8 -4.0 (m, 4H), 7.1 -8.95 (m, 12H).

5 2-(4-Methoxy-benzenesulfonyl)-6-phenoxy-2-pyridin-3-ylmethyl-hexanoic acid
hydroxyamide

Following the procedure as outlined in Example 9, 2-(4-methoxy-benzenesulfonyl)-6-
15 phenoxy-2-pyridin-3-ylmethyl-hexanoic acid ethyl ester was prepared, starting from
(3.1 g, 10 mmol) 2-(4-methoxy-benzenesulfonyl)-6-phenoxy-hexanoic acid ethyl ester
and 3-picolyl chloride (1.8 g, 11 mmol). Yield 3.5 g, 71%; Colorless oil; MS: 498
(M+H)⁺.

25 Starting from 2-(4-methoxy-benzenesulfonyl)-6-phenoxy-2-pyridin-3-ylmethyl-hexanoic acid (2.0 g, 4.3 mmol) and following the procedure as outlined in Example 1, 1.5 g of 2-(4-methoxy-benzenesulfonyl)-6-phenoxy-2-pyridin-3-ylmethyl-hexanoic acid hydroxyamide was isolated as a colorless solid; Yield: 72%; mp 68 °C; MS: 485.5

30 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.5 - 2.5 (m, 8H), 3.4 (bs, 2H), 3.8 (s, 3H), 6.8 - 8.7 (m, 13H).

WO 99/42436

PCT/US98/17633

- 72 -

Example 69

2-(4-Methoxy-benzenesulfonyl)-5-methyl-2-pyridin-3-ylmethyl-hexanoic acid hydroxyamide

5

Following the procedure as outlined in Example 9, 2-(4-methoxy-benzenesulfonyl)-5-hexanoic acid ethyl ester was prepared, starting from (10.0 g, 39 mmol) 2-(4-methoxy-benzenesulfonyl)-acetic acid ethyl ester and 1-bromo-3-methyl butane (6.0 g, 40 mmol). Yield 8.5 g, 62%; Colorless oil; MS: 329 (M+H)⁺.

10

Following the procedure as outlined in Example 9, 2-(4-methoxy-benzenesulfonyl)-5-methyl-2-pyridin-3-ylmethyl-hexanoic acid ethyl ester was prepared, starting from (6.0 g, 18 mmol) of 2-(4-methoxy-benzenesulfonyl)-5-methyl-hexanoic acid ethyl ester and picolyl chloride hydrochloride (4.1 g, 25 mmol). Yield 4.5 g, 60%; Brown oil; MS:

15

420 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-5-methyl-2-pyridin-3-ylmethyl-hexanoic acid ethyl ester (3.0 g, 7.1 mmol) 2.6g (92%) of 2-(4-methoxy-benzenesulfonyl)-5-methyl-2-pyridin-3-ylmethyl-hexanoic acid was isolated as a colorless solid by

20

following the procedure as outlined in Example 9. Mp: 173 C; MS: 392 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-5-methyl-2-pyridin-3-ylmethyl-hexanoic acid (1.0 g, 2.5 mmol) and following the procedure as outlined in Example 1, 800 mg of 2-(4-methoxy-benzenesulfonyl)-5-methyl-2-pyridin-3-ylmethyl-hexanoic acid

25

hydroxyamide was isolated as a colorless solid; The hydrochloride was prepared by passing hydrogen chloride gas through methanol solution of the hydroxyamide. Yield: 72%; mp 62 °C (HCl salt); MS: 408 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 0.76 (m, 6H), 1.2 -2.0 (m, 5H), 3.5 (bq, 2H), 7.1 - 8.8 (m, 8H), 11.1 (bs, 1H).

30

Example 70

2-(4-Methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-hexanoic acid hydroxyamide

35

(4-Methoxy-phenylsulfanyl) -acetic acid tert-butyl ester was prepared according to the general method as outlined in Example 1. Starting from the corresponding 1-bromo tert-butyl acetate (5.3 g, 27 mmol) and 4-methoxybenzenethiol (3.7 g, 27 mmol), 6.4 g of the product was isolated. Yield 98%; Light yellow oil; MS: 255 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 73 -

2-(4-Methoxy-benzenesulfonyl)-acetic acid tert-butyl ester was prepared according to the general method as outlined in Example 9. Starting from 2-(4-methoxy-benzenesulfonyl)-acetic acid tert-butyl ester (5.0 g, 20 mmol) and 3-chloroperoxybenzoic acid 57% (12.0g, 40 mmol), 5.3 g of the product was isolated. Yield 92%; Waxy solid;

5 MS: 287.1 (M+H)⁺.

2-(4-Methoxy-benzenesulfonyl)-pyridin-3-ylpropionic acid tert-butyl ester was prepared according to the procedure as outlined in Example 9. Starting from 2-(4-methoxy-benzenesulfonyl)acetic acid tert-butyl ester (20.0 g, 70.0 mmol) and 3-picolyl chloride (7.28 g, 44.4 mmol), 10.5 g of the product was isolated by silica gel chromatography (50% ethyl acetate: hexane). Yield 63%; white solid; mp 93-94 °C; MS: 378.0 (M+H)⁺.

15 2-(4-Methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-hexanoic acid tert-butyl ester was prepared according to the procedure as outlined in Example 9. Starting from 2-(4-methoxy-benzenesulfonyl)-pyridin-3-ylpropionic acid tert-butyl ester (2.0 g, 5.3 mmol) and n-butyl bromide (0.73 g, 5.3 mmol), 1.20 g of the product isolated. Yield 52%; yellowish gum; MS: 434.3 (M+H)⁺.

20 A mixture of the 2-(4-Methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-hexanoic acid tert-butyl ester (1.1 g, 2.5 mmol), in methylene chloride/ TFA (1:1) was stirred at room temperature for about 2 hours. The solvents were then evaporated and the 2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-hexanoic acid was purified by silica gel chromatography (30% methanol/methylene chloride). Yield 0.90 g, 94%; white solid; mp 70 °C; MS: 376.1 (M-H)⁻.

2-(4-Methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-hexanoic acid hydroxyamide was prepared according to the method as outlined in Example 1. Starting from 2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-hexanoic acid (0.31 g, 0.81 mmol) and hydroxylamine hydrochloride (0.70 g, 10 mmol), 0.13 g of the product isolated. Yield 37%; pale yellowish solid; mp 65 °C; MS: 392.9 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 0.80 (t, J = 7.2 Hz, 3H), 1.10-1.25 (m, 2H), 1.25-1.50 (m, 2H), 1.70-2.00 (m, 2H), 3.53 (d, J = 14.4 Hz, 1H), 3.62 (d, J = 14.4 Hz, 1H), 3.88 (s, 3H), 7.15 (d, J = 8.9 Hz, 2H), 7.71 (d, J = 8.9 Hz, 2H), 7.90-8.00 (m, 1H), 8.40-8.45 (m, 1H), 8.70-8.85 (m, 2H), 11.0 (brs, 1H); IR (KBr, cm⁻¹): 3064m, 2958s, 2871m, 1671m.

WO 99/42436

PCT/US98/17633

- 74 -

Example 71

2-(4-methoxy-benzenesulfonyl)-2-oct-2-ynyl-dec-4-ynoic acid hydroxyamide.

5

The title compound was prepared according to the procedure as outlined in example 9. Starting from 2-(4-methoxy-benzenesulfonyl)-acetic acid tert-butyl ester (2.86 g, 10 mmol) and 1-bromo-2-octyne (3.80 g, 20 mmol), 4.4 g of the product isolated. Yield 100%; yellowish gum; MS: 446.9 (M+H)⁺.

10

2-(4-Methoxy-benzenesulfonyl)-2-oct-2-ynyl-dec-4-ynoic acid was prepared according to the method as outlined in example 70. Starting from 2-(4-methoxy-benzenesulfonyl)-2-oct-2-ynyl-dec-4-ynoic acid tert-butyl ester (4.40 g, 10.0 mmol), 2.0 g of the product isolated. Yield 49%; white solid; mp 61 °C; MS: 345.1 (M-H)⁻.

15

2-(4-Methoxy-benzenesulfonyl)-2-oct-2-ynyl-dec-4-ynoic acid hydroxyamide was prepared according to the method as outlined in example 1. Starting from 2-(4-methoxy-benzenesulfonyl)-2-oct-2-ynyl-dec-4-ynoic acid (0.36 g, 0.81 mmol) and hydroxylamine hydrochloride (0.70 g, 10 mmol), 0.25 g of the product isolated. Yield

20

62%; white solid; mp 83-84 °C; 462.0 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 0.82-0.90 (m, 6H), 1.15-1.45 (m, 12H), 1.90-2.05 (m, 4H), 2.86 (brd, J = 17.0 Hz, 2H), 3.00 (brd, J = 17.0 Hz, 2H), 3.87 (s, 3H), 7.15 (d, J = 10.0 Hz, 1H), 7.71 (d, J = 10.0 Hz, 1H), 9.20 (brs, 1H), 10.90 (brs, 1H); IR (KBr, cm⁻¹): 3344s, 3208m, 2930m, 2870m, 1677s, 1592s;

25

Anal. Calc'd for C₂₅H₃₅NO₅S: C, 65.05; H, 7.64; N, 3.03.

Found:

C, 65.26; H, 7.68; N, 2.90.

Example 72

30

2-(4-Methoxy-benzenesulfonyl)-2-but-2-ynyl-hex-4-ynoic acid hydroxyamide

2-(4-Methoxy-benzenesulfonyl)-2-but-2-ynyl-hex-4-ynoic acid tert-butyl ester was prepared according to the procedure as outlined in Example 9. Starting from 2-(4-methoxy-benzenesulfonyl)-acetic acid tert-butyl ester (2.86 g, 10 mmol) and 1-bromo-

35

2-butyne (2.68 g, 20 mmol), 3.50 g of the product was isolated. Yield 90%; white solid; mp 85-87 °C; MS: 391.0 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 75 -

2-(4-Methoxy-benzenesulfonyl)-2-but-2-ynyl-hex-4-ynoic acid was prepared according to the procedure as outlined in example 70. Starting from 2-(4-methoxy-benzenesulfonyl)-2-but-2-ynyl-hex-4-ynoic acid tert-butyl ester (3.0 g, 7.7 mmol), 2.5 g of the product isolated. Yield 97%; white solid; mp 141-143 °C; MS: 333.1 (M-H)⁻.

- 5 2-(4-Methoxy-benzenesulfonyl)-2-but-2-ynyl-hex-4-ynoic acid hydroxyamide was prepared according to the method as outlined in example 1. Starting from 2-(4-methoxy-benzenesulfonyl)-2-but-2-ynyl-hex-4-ynoic acid (0.27 g, 0.81 mmol) and hydroxylamine hydrochloride (0.70 g, 10 mmol), 0.23 g of the product was isolated.
- 10 Yield 89%; white solid; mp 135-137°C; MS: 349.9 (M+H)⁺1; ¹H NMR (300 MHz, DMSO-d₆) δ 1.67 (s, 6H), 2.70-3.10 (m, 4H), 3.88 (s, 3H), 7.15 (d, J = 10.0 Hz, 2H), 7.71 (d, J = 10.0 Hz, 2H), 9.20 (brs, 1H), 10.90 (brs, 1H); IR (KBr, cm⁻¹): 3301s, 3161m, 2922m, 1640m, 1595s, 1500m.

15

Example 73

2-(4-Methoxy-benzenesulfonyl)-2-prop-2-ynyl-pent-4-ynoic acid hydroxyamide

- 20 2-(4-Methoxy-benzenesulfonyl)-2-prop-2-ynyl-pent-4-ynoic acid tert-butyl ester was prepared according to the procedure as outlined in Example 9. Starting from 2-(4-methoxy-benzenesulfonyl)-acetic acid tert-butyl ester (2.0 g, 7.0 mmol) and propargyl bromide (1.77 g, 15 mmol), 1.90 g of the product was isolated. Yield 75%; white solid; mp 113-115°C; MS: 362.1 (M+H)⁺.
- 25 2-(4-Methoxy-benzenesulfonyl)-2-prop-2-ynyl-pent-4-ynoic acid was prepared according to the procedure as outlined in Example 70. Starting from 2-(4-methoxy-benzenesulfonyl)-2-prop-2-ynyl-pent-4-ynoic acid tert-butyl ester (1.70 g, 4.7 mmol), 1.30 g of the product isolated. Yield 90%; white solid; mp 156°C; MS: 305.1 (M-H)⁻.
- 30 2-(4-Methoxy-benzenesulfonyl)-2-prop-2-ynyl-pent-4-ynoic acid hydroxyamide was prepared according to the method as outlined in Example 1. Starting from (4-methoxy-benzenesulfonyl)-2-prop-2-ynyl-pent-4-ynoic acid (0.25 g, 0.81 mmol) and hydroxylamine hydrochloride (0.70 g, 10 mmol), 0.22 g of the product was isolated.
- 35 Yield 85%; white solid; mp 156°C; MS: 321.9 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 2.00-2.13 (m, 2H), 3.00-3.30 (m, 4H), 3.90(s, 3H), 7.01 (d, J = 9.0 Hz, 2H),

WO 99/42436

PCT/US98/17633

- 76 -

7.82 (d, J = 9.0 Hz, 2H), 8.76 (brs, 1H), 10.65 (brs, 1H); IR (KBr, cm⁻¹): 3392s, 3293s, 3271m, 2955m, 1650s, 1594s;

Anal. Calc'd for C₁₅H₁₅NO₅S: C, 56.07; H, 4.70; N, 4.36.

Found: C, 55.65; H, 4.67; N, 4.10.

5

Example 74

2-(4-Methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-dec-4-ynoic acid hydroxyamide

10 The title compound was prepared according to the procedure as outlined in Example 38. Starting from 2-(4-methoxy-benzenesulfonyl)-pyridin-3-ylpropionic acid tert-butyl ester (2.20 g, 5.8 mmol) and 1-bromo-2-octyne (1.14 g, 6 mmol), 2.60 gm of the product isolated. Yield 92%; yellowish gum; MS: 486.0 (M+H)⁺.

15 A mixture of the 2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-dec-4-ynoic acid tert-butyl ester (2.60 g, 5.35 mmol), in methylene chloride/TFA (1:1) is stirred at room temperature for about 2 hours. (Ref. example 70) The solvents are then evaporated and the 2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-dec-4-ynoic acid was purified by silica gel chromatography (~30% methanol/methylene chloride).

20 Yield: 2.0 g, 87%; White solid; mp 146°C; MS: 428.1 (M-H)⁻.

2-(4-Methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-dec-4-ynoic acid hydroxyamide was prepared according to the procedure outlined in Example 1. Starting from 2-(4-

25 methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-dec-4-ynoic acid (0.71 g, 1.62 mmol) and hydroxylamine hydrochloride (1.39 g, 20 mmol), 0.48 g of the product was isolated. Yield 67%; off-white solid; mp 65°C; MS: 445.0 (M+H)⁺; ¹H NMR (300

MHz, DMSO-d₆) δ 0.84 (t, J = 6.8 Hz, 3H), 1.10-1.40 (m, 6H), 1.85-2.00 (m, 2H),

2.79 (d, J = 17.9 Hz, 1H), 2.90 (d, J = 17.9 Hz, 1H), 3.50 (d, J = 13.7 Hz, 1H),

3.74 (d, J = 13.7 Hz, 1H), 3.89 (s, 3H), 7.19 (d, J = 9.0 Hz, 2H), 7.76 (d, J = 9.0

30 Hz, 2H), 7.85-7.89 (m, 1H), 8.37-8.40 (m, 1H), 8.70-8.80 (m, 2H), 11.0 (brs, 1H);

IR (KBr, cm⁻¹): 3157m, 3095m, 2954s, 2932s, 2858m, 1671m, 1593s;

Anal. Calc'd for C₂₃H₂₈N₂O₅S·HCl·0.9H₂O: C, 55.56; H, 6.24; N, 5.63.

Found: C, 55.84; H, 6.19; N, 5.59.

WO 99/42436

PCT/US98/17633

- 77 -

Example 75

2-(4-Methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-pent-4-ynoic acid hydroxyamide

5 2-(4-Methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-pent-4-ynoic acid tert-butyl ester was prepared according to the procedure as outlined in Example 38. Starting from 2-(4-methoxy-benzenesulfonyl)-pyridin-3-ylpropionic acid tert-butyl ester (3.77 g, 10 mmol) and propargyl bromide (1.74 g, 13 mmol), 2.50 g of the product was isolated. Yield 60%; yellowish solid; mp 132-133°C; MS: 416.0 (M+H)⁺.

10 2-(4-Methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-pent-4-ynoic acid was prepared according to the procedure as outlined in Example 70. Starting from 2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-pent-4-ynoic acid tert-butyl ester (2.0 g, 4.8 mmol), 1.2 g of the product isolated. Yield 69%; white solid; mp 119-121°C; MS:

15 358.1 (M-H)⁻.

2-(4-Methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-pent-4-ynoic acid hydroxyamide was prepared according to the method as outlined in Example 1. Starting from 2-(4-methoxy-benzenesulfonyl)-2-pyridin-3-ylmethyl-pent-4-ynoic acid (0.29 g, 0.81 mmol) and hydroxylamine hydrochloride (0.70 m, 10 mmol), 0.065 g of the product was isolated. Yield 25%; off-white solid; mp 70°C; MS: 375.0 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 1.19 (brs, 1H), 2.90-3.00 (m, 2H), 3.55 (d, J = 13.8 Hz, 1H), 3.67 (d, J = 13.8 Hz, 1H), 3.89 (s, 3H), 7.18 (d, J = 9.0 Hz, 2H), 7.75 (d, J = 9.0 Hz, 2H), 7.80-7.89 (m, 1H), 8.35-8.40 (m, 1H), 8.70-8.80 (m, 2H), 11.1 (brs, 1H); IR (KBr, cm⁻¹): 3168m, 3095s, 1670m, 1593s.

25

Example 76

2-(4-Fluoro-benzenesulfonyl)-2-pyridin-3-ylmethyl-hex-4-ynoic acid hydroxyamide

30

2-(4-Fluoro-benzenesulfanyl)-acetic acid tert-butyl ester was prepared according to the procedure as outlined in Example 1. Starting from 4-fluorothiophenol (30.0 g, 230 mmol) and tert-butyl bromoacetate (45.67 g, 230 mmol), 53.4 g of the product was isolated. Yield 100%; pale yellowish oil; MS: 243.1 (M+H)⁺.

35

2-(4-Fluoro-benzenesulfonyl)-acetic acid tert-butyl ester was prepared according to the general method as outlined in Example 9. Starting from 2-(4-fluoro-benzenesulfanyl)-

WO 99/42436

PCT/US98/17633

- 78 -

acetic acid tert-butyl ester (48.4 g, 200 mmol) and 3-chloroperoxybenzoic acid (121.3g (57%), 400 mmol), 48.0 g of the product was isolated. Yield 88%; pale yellowish oil; MS: 275.1 (M+H)⁺.

5 The title compound was prepared according to the procedure as outlined in Example 70. Starting from 2-(4-fluoro-benzenesulfonyl)-3-pyridin-3-ylpropionic acid tert-butyl ester (1.83 g, 5.0 mmol) and 1-bromo-2-butyne (0.67 g, 5.0 mmol), 2.18 g of the product was isolated. Yield 100%; yellowish gum; MS: 419.2 (M+H)⁺.

10 2-(4-Fluoro-benzenesulfonyl)-2-pyridin-3-ylmethyl-hex-4-ynoic acid was prepared according to the method as outlined in Example 38. Starting from 2-(4-fluoro-benzenesulfonyl)-2-pyridin-3-ylmethyl-hex-4-ynoic acid tert-butyl ester (2.1 g, 5.0 mmol), 1.20 g of the product was isolated. Yield 67%; off-white solid; mp 150°C; MS: 360.2 (M-H)⁻.

15 2-(4-Fluoro-benzenesulfonyl)-2-pyridin-3-ylmethyl-hex-4-ynoic acid hydroxyamide was prepared according to the method as outlined in Example 1. Starting from 2-(4-fluoro-benzenesulfonyl)-2-pyridin-3-ylmethyl-hex-4-ynoic acid (0.29 g, 0.81 mmol) and hydroxylamine hydrochloride (0.70 g, 10 mmol), 0.15 g of the product was
20 isolated. Yield 45%; white solid; mp 190°C; MS: 377.2 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 1.60 (s, 3H), 2.70-3.00 (m, 2H), 3.53 (d, J = 13.8 Hz, 1H), 3.74 (d, J = 13.8 Hz, 1H), 7.50-7.58 (m, 2H), 7.80-7.95 (m, 3H), 8.35-8.40 (m, 1H), 8.74-8.79 (m, 2H), 11.1 (brs, 1H); IR (KBr, cm⁻¹): 3154m, 3105s, 3068s, 2875m, 1696s, 1630w, 1590s;

25 Anal. Calc'd for C₁₈H₁₇FN₂O₄S·HCl·0.5H₂O: C, 51.24; H, 4.54; N, 6.64.
Found: C, 51.21; H, 4.35; N, 6.46.

Example 77

30 2-(4-Fluoro-benzenesulfonyl)-2-pyridin-3-ylmethyl-dec-4-ynoic acid hydroxyamide

The title compound was prepared according to the procedure as outlined in Example 9. Starting from 2-(4-fluoro-benzenesulfonyl)-3-pyridin-3-ylpropionic acid tert-butyl ester (1.83 g, 5.0 mmol) and 1-bromo-2-octyne (0.95 g, 5.0 mmol), 1.80 g of the product

35 was isolated. Yield 56%; yellowish gum; MS: 474.3 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 79 -

2-(4-Fluoro-benzenesulfonyl)-2-pyridin-3-ylmethyl-dec-4-ynoic acid was prepared according to the method as outlined in Example 70. Starting from 2-(4-fluoro-benzenesulfonyl)-2-pyridin-3-ylmethyl-dec-4-ynoic acid tert-butyl ester (1.80 g, 3.8 mmol), 1.40 g of the product was isolated. Yield 88%; off-white solid; mp 123-124°C; MS:

5 416.3 (M-H)⁻.

2-(4-Fluoro-benzenesulfonyl)-2-pyridin-3-ylmethyl-dec-4-ynoic acid hydroxyamide was prepared according to the method as outlined in Example 1. Starting from 2-(4-fluoro-benzenesulfonyl)-2-pyridin-3-ylmethyl-dec-4-ynoic acid (0.67 g, 1.62 mmol) and hydroxylamine hydrochloride (1.39 g, 20 mmol), 0.22 g of the product was isolated. Yield 29%; white solid; mp 180-182°C; MS: 433.2 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 0.84 (t, J = 6.8 Hz, 3H), 1.20-1.40 (m, 6H), 1.90-2.05 (m, 2H), 2.75 (d, J = 19.9 Hz, 1H), 2.94 (d, J = 19.9 Hz, 1H), 3.54 (d, J = 13.7 Hz, 1H), 3.75 (d, J = 13.7 Hz, 1H), 7.40-7.60 (m, 2H), 7.70-8.00 (m, 3H), 8.30-8.40 (m, 1H), 8.70-8.80 (m, 2H), 11.1 (brs, 1H); IR (KBr, cm⁻¹): 3154m, 3105s, 3067m, 2957s, 2933s, 2873m, 1690s, 1631m.

Anal. Calc'd for C₂₂H₂₅N₂O₄S·HCl: C, 56.34; H, 5.59; N, 5.97.

Found: C, 56.18; H, 5.54; N, 5.76.

20

Example 78

2-(4-Fluoro-benzenesulfonyl)-2-but-2-ynyl-hex-4-ynoic acid hydroxyamide

2-(4-Fluoro-benzenesulfonyl)-2-but-2-ynyl-hex-4-ynoic acid tert-butyl ester was prepared according to the procedure as outlined in Example 9. Starting from 2-(4-fluoro-benzenesulfonyl)-acetic acid tert-butyl ester (4.87 g, 20 mmol) and 1-bromo-2-butyne (5.36 g, 40 mmol), 6.0 g of the product was isolated. Yield 77%; white solid; mp 85°C; MS: 379.1 (M+H)⁺.

2-(4-Fluoro-benzenesulfonyl)-2-but-2-ynyl-hex-4-ynoic acid was prepared according to the procedure as outlined in Example 70, starting from 2-(4-fluoro-benzenesulfonyl)-2-but-2-ynyl-hex-4-ynoic acid tert-butyl ester (3.50 g, 8.47 mmol), 2.35 g of the product was isolated. Yield 79%; white solid; mp 129-131°C; MS: 642.8 (2M-H)⁻.

2-(4-Fluoro-benzenesulfonyl)-2-but-2-ynyl-hex-4-ynoic acid hydroxyamide was prepared according to the method as outlined in Example 1. . Starting from 2-(4-fluoro-

WO 99/42436

PCT/US98/17633

- 80 -

benzenesulfonyl)-2-but-2-ynyl-hex-4-ynoic acid (0.26 g, 0.81 mmol) and hydroxylamine hydrochloride (0.70 g, 10 mmol), 0.21 g of the product was isolated.

Yield 77%; white solid; mp 161-163°C; MS: 338.1(M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 1.67 (s, 6H), 2.80-3.10 (m, 4H), 7.51 (dd, J = 9.0, 9.0 Hz, 2H), 7.87

5 (m, 2H), 9.26 (brs, 1H), 10.95 (brs, 1H); IR (KBr, cm⁻¹): 3336s, 3245m, 1681s, 1589m, 1493m;

Anal. Calc'd for C₁₆H₁₆FNO₄S: C, 56.96; H, 4.78; N, 4.15.

Found: C, 56.59; H, 4.75; N, 4.04.

10 Example 79

2-(4-Methoxy-benzenesulfonyl)-5-methyl-2-(3-methyl-but-2-enyl)-hex-4-enoic acid hydroxyamide

15 Following the procedure as outlined in Example 9, 2-(4-methoxy-benzenesulfonyl)-5-methyl-2-(3-methyl-but-2-enyl)-hex-4-enoic acid ethyl ester was prepared, starting from (5.0 g, 20 mmol) 2-(4-methoxy-benzenesulfonyl)-acetic acid ethyl ester and isoprenyl bromide (6.0 g, 40 mmol). Yield 7.0 g, 88%; Colorless oil; MS: 395 (M+H)⁺.

20 Starting from 2-(4-methoxy-benzenesulfonyl)-5-methyl-2-(3-methyl-but-2-enyl)-hex-4-enoic acid ethyl ester (3.5 g, 9 mmol), 3.3g (97%) of 2-(4-methoxy-benzenesulfonyl)-5-methyl-2-(3-methyl-but-2-enyl)-hex-4-enoic acid was isolated as a colorless oil by following the procedure as outlined in Example 9. MS: 365 (M-H)⁻.

25 Starting from 2-(4-methoxy-benzenesulfonyl)-5-methyl-2-(3-methyl-but-2-enyl)-hex-4-enoic acid (2.6 g, 7.0 mmol) and following the procedure as outlined in Example 1, 1.36 g of 2-(4-methoxy-benzenesulfonyl)-5-methyl-2-(3-methyl-but-2-enyl)-hex-4-enoic acid hydroxyamide was isolated as a colorless solid. Yield: 67%; mp 93 - 96 °C;

30 MS: 383 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): δ 1.68 (s, 6H), 1.73 (s, 6H), 2.72 (m, 4H), 3.82 (s, 3H), 5.12 (m, 2H), 6.92 (d, J=8 Hz, 2H), 7.33 (bs, 1H), 7.72 (d, J=8 Hz, 2H), 9.71 (bs, 1H).

WO 99/42436

PCT/US98/17633

- 81 -

Example 80

2-(4-methoxy-phenylsulfanyl)-heptanoic acid hydroxyamide.

- 5 2-(4-Methoxy-phenylsulfanyl)-heptanoic acid ethyl ester (13.8 g, 98%) was prepared according to the general method as outlined in example 1 starting from ethyl 2-bromo-heptanoate (11 g, 47 mmol) and 4-methoxythiophenol (6g, 42.8 mmol), as a yellow oil; MS: 297.2 (M+H)⁺.
- 10 2-(4-Methoxy-phenylsulfanyl)-heptanoic acid was prepared starting with 2-(4-methoxy-phenylsulfanyl)-heptanoic acid ethyl ester (4 g, 13.5 mmol) dissolved in methanol (300 ml) and 10 N NaOH (25 ml). The resulting reaction mixture was worked up as outlined in example 1. Yield 3 g (83%). yellow oil. MS: 267.1 (M-H)⁻.
- 15 Starting from 2-(4-methoxy-phenylsulfanyl)-heptanoic acid (2.49 g, 9.32 mmol) and following the procedure as outlined in example 1, 1.83 g of 2-(4-methoxy-phenylsulfanyl)-heptanoic acid hydroxyamide was isolated as an off white solid. Mp 90-95°C; Yield 70%; MS: 284.0 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.826 (t, J= 6.9 Hz, 3H), 1.135-1.76 (m, 8H), 3.35 (m, 1H), 3.82 (s, 3H), 6.91-7.49 (m, 4H).

20

Example 81

(49A) 2R^{*}-(4-methoxy-phenyl- S^{*}- sulfinyl)-heptanoic acid hydroxyamide
and

25 (49B) 2S^{*}-(4-methoxy-phenyl- R^{*}- sulfinyl)-heptanoic acid hydroxyamide

- Starting from 2-(4-methoxy-phenylsulfanyl)-heptanoic acid hydroxyamide (1.69 g, 6 mmol) and following the procedure outlined in example 5, the two diastereomers of 2-(4-methoxy-phenylsulfinyl)-heptanoic acid hydroxyamide were separated on a silica gel
- 30 column using 75% ethyl acetate:hexanes. The less polar isomer, 2R^{*}-(4-methoxy-phenyl- S^{*}- sulfinyl)-heptanoic acid hydroxyamide was isolated as a white powder. Yield: 390 mg (22%); mp 115 °C; MS: 300.0 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): 0.828 (t, J= 6.2 Hz, 3H), 1.18-1.23 (m, 6H), 1.73-1.99 (m, 2H), 3.11-3.15 (m, 1H), 3.82 (s, 3H), 7.09-7.61 (m, 4H). The more polar isomer, 2S^{*}-(4-methoxy-
- 35 phenyl- R^{*}- sulfinyl)-heptanoic acid hydroxyamide was isolated as a gray solid. Yield: 200 mg (11%); mp 112 °C; MS: 300.0 (M+H)⁺, ¹H NMR (300 MHz, DMSO-d₆): δ 0.754 (t, J= 6.9 Hz, 3H), 1.014-1.121 (m, 6H), 1.58-1.89 (m, 2H), 3.10-3.15 (m, 1H), 3.834 (s, 3H), 7.13-7.65 (m, 4H).

WO 99/42436

PCT/US98/17633

- 82 -

Example 82

2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-morpholin-4-yl-ethoxy)-phenyl]-
propionic hydroxyamide hydrochloride.

Following the procedure as outlined in example 12, 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-morpholin-1-yl-ethoxy)-phenyl]-propionic acid ethyl ester was prepared, starting from (4.0 g, 15 mmol) of 2-(4-methoxy-benzenesulfonyl)-propionic acid ethyl ester and 4-(morpholin-1-yl-ethoxy)-benzyl chloride hydrochloride (2.9 g, 10 mmol). Yield 4.8 g, 98%; Brown oil; MS: 492 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-morpholin-1-yl-ethoxy)-phenyl]-propionic acid ethyl ester (4.0 gm, 8.1 mmol) 3.2 g (Yield: 84 %) of 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-morpholin-1-yl-ethoxy)-phenyl]-propionic acid was isolated as colorless crystals by following the procedure as outlined in example 9. Mp 171 °C; MS: 464 (M+H)⁺.

Starting from 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-morpholin-1-yl-ethoxy)-phenyl]-propionic acid (4.0 g, 8.6 mmol) and following the procedure as outlined in example 1, 2.5 g of 2-(4-methoxy-benzenesulfonyl)-2-methyl-3-[4-(2-morpholin-1-yl-ethoxy)-phenyl]-propionic hydroxyamide was isolated as colorless solid. The hydrochloride salt was prepared by reacting the free base with methanolic hydrogen chloride at 0 °C. Yield: 2.5 g, 60%; mp 98°C; MS: 479 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃): 1.36 (s, 3H), 3.8 - 12.6 (m, 16 H), 3.9 (s, 3H), 4.1 - 4.3 (m, 1H), 6.6 (d, J= 8 Hz, 2H), 6.96 (d, J= 9 Hz, 2H), 7.1 (d, 8 Hz, 2H), 7.84 (d, 9 Hz, 2H), 10.8 (bs, 1H).

Example 83

1-Benzyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic Acid hydroxyamide

To a stirred solution of 4-methoxybenzenethiol (2.8 gm, 20 mmol) and anhydrous K₂CO₃ (10 gm, excess) in dry acetone (100 ml), α-bromo ethyl acetate (3.3 gm, 20 mmol) was added in a round bottom flask and the reaction mixture was heated at reflux for 8 hours with good stirring. At the end, the reaction mixture was allowed to cool and the potassium salts were filtered off and the reaction mixture was concentrated. The residue was extracted with chloroform and washed with H₂O and 0.5 N NaOH solution. The organic layer was further washed well with water, dried over MgSO₄,

WO 99/42436

PCT/US98/17633

- 83 -

filtered and concentrated. (4-methoxy-phenylsulfonyl)-acetic acid ethyl ester was isolated as pale yellow oil. Yield: 4.4 g (100%); MS; 227 (M+H)⁺.

To a stirred solution of 60% 3-chloroperoxybenzoic acid (14.0 gm, 40 mmol) in methylene chloride (100 ml) at 0° C, (4-methoxy-phenylsulfonyl)-acetic acid ethyl ester (4.4 g, 20 mmol) in CH₂Cl₂ (15 ml) was added slowly. The reaction mixture turned cloudy and was stirred at room temperature for 6 hours. The reaction mixture was then diluted with hexanes (300 ml) and stirred for 15 minutes. The solids were filtered off and Na₂SO₃ solution was added to the organic layer which was stirred for at least 3 hours before the mixture was extracted with CHCl₃ and washed with H₂O. The organic layer was dried over MgSO₄, filtered and concentrated and the colorless (4-methoxy-phenylsulfonyl)-acetic acid ethyl ester was isolated as an oil. Yield: 100%; MS: 259.1 (M+H)⁺.

To a stirred solution of diethanol amine (10.5 g, 100 mmol), and anhydrous K₂CO₃ (30 gm, excess) in dry acetone (250 ml), benzyl bromide (17.2 gm, 100 mmol) was added in a round bottom flask and the reaction mixture was heated at reflux for 8 hours with good stirring. At the end, the reaction mixture was allowed to cool and the potassium salts were filtered off and the reaction mixture was concentrated. The residue was extracted with chloroform and washed with H₂O. The organic layer was further washed well with water, dried over MgSO₄, filtered and concentrated. Colorless oil. Yield: 19.0 g, 97%; MS: 196 (M+H).

N-Benzyl diethanolamine (9.75 g, 50 mmol) was dissolved in saturated methanolic hydrochloric acid and concentrated to dryness. The hydrochloride thus formed was dissolved in methylene chloride (300 ml) and thionyl chloride (20 g, excess) was added dropwise and stirred at room temperature for 1 hr. At the end reaction mixture was concentrated to dryness and the product bis-(2-chloro-ethyl)-benzyl amine hydrochloride was used for further transformation without any purification. Yield: 13.0 g, 97%; Mp: MS: 232 (M+H).

To a stirred solution of bis-(2-chloro-ethyl)-benzyl amine hydrochloride (6.6 g, 24.7 mmol), 18-Crown-6 (500 mg), and anhydrous K₂CO₃ (30 gm, excess) in dry acetone (250 ml), (4-methoxy-phenylsulfonyl)-acetic acid ethyl ester (6.12 gm, 24 mmol) was added in a round bottom flask and the reaction mixture was heated at reflux for 16 hours with good stirring. At the end, the reaction mixture was allowed to cool and the potassium salts were filtered off and the reaction mixture was concentrated. The

WO 99/42436

PCT/US98/17633

- 84 -

residue was extracted with chloroform and washed with H₂O. The organic layer was further washed well with water, dried over MgSO₄, filtered and concentrated. The dark brown reaction mixture was purified by silica gel column chromatography by eluting it with 30% ethylacetate: hexane and the product 4-(4-Methoxy-benzenesulfonyl)-1-benzyl-piperidine-4-carboxylic acid ethyl ester was isolated as Brown oil. Yield: 6.0 g, 60%; MS: 418 (M+H).

4-(4-Methoxy-benzenesulfonyl)-1-benzyl-piperidine-4-carboxylic acid ethyl ester (5.0 g, 11.9 mmol) was dissolved in MeOH/THF (1:1, 200 ml) and stirred at room temperature for 72 hrs. At the end reaction mixture was concentrated and the product was neutralised with con. Hcl by dissolving it in water (200 ml). After the neutralization reaction mixture was concentrated to dryness. Ice cold water (100 ml) was added to the solid and filtered. The product 4-(4-Methoxy-benzenesulfonyl)-1-benzyl-piperidine-4-carboxylic acid was dried at 50 °C and taken to next step without any purification. Colorless solid. Yield: 3.2 g, 69% ; MS: 390 (M+H).

To a stirred solution of 4-(4-Methoxy-benzenesulfonyl)-1-benzyl-piperidine-4-carboxylic acid (2.0 g, 5.1 mmol) and DMF (2 drops) in CH₂Cl₂ (100 ml) at 0°C, oxalyl chloride (1.0 gm, 8 mmol) was added in a drop-wise manner. After the addition, the reaction mixture was stirred at room temperature for 1 hour. Simultaneously, in a separate flask a mixture of hydroxylamine hydrochloride (2.0 gm, 29 mmol) and triethylamine (5 ml, excess) was stirred in THF:water (5:1, 30 ml) at 0°C for 1 hour. At the end of 1 hour, the oxalyl chloride reaction mixture was concentrated and the pale yellow residue was dissolved in 10 ml of CH₂Cl₂ and added slowly to the hydroxylamine at 0°C. The reaction mixture was stirred at room temperature for 24 hours and concentrated. The residue obtained was extracted with chloroform and washed well with water. The product obtained was purified by silica gel column chromatography and eluted with chloroform the product 4-(4-Methoxy-benzenesulfonyl)-1-benzyl-piperidine-4-carboxylic acid hydroxyamide was isolated as a colorless solid. mp 90-95 °C; Yield, 1.2 g, 48%; MS: 405 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 2.29 (m, 3H), 2.76-2.79 (m, 2H), 3.43 (m, 4H), 4.30 (s, 2H), 7.14-7.17 (d, 2H), 7.50-7.73 (m, 5H), 9.37 (s, 1H), 10.53 (s, 1H), 11.18 (s, 1H).

WO 99/42436

PCT/US98/17633

- 85 -

Example 84

4-(4-methoxy-benzenesulfonyl)-1-(3-methoxy-benzyl)-piperidine-4-carboxylic acid hydroxyamide

- 5 2-[(2-Hydroxy-ethyl)-(3-methoxy-benzyl)-amino]-ethanol was prepared according to the general method as outlined in example 83. Starting from diethanolamine (3.1 g, 29.5 mmol) and 3-methoxybenzyl chloride (5 g, 31.9 mmol). Yield 9.28 g, (99 %); yellow oil; MS: 226 (M+H).
- 10 3-Methoxybenzyl-bis-(2-chloro-ethyl)-amine was prepared according to the general method as outlined in example 83. Starting from 3-Methoxy-benzyl diethanolamine (4.4 g, 20 mmol). Yield 4.5 g (93 %); yellow solid mp 86 -88 °C; MS: 263. (M+H)⁺.
- 15 4-(4-Methoxy-benzenesulfonyl)-1-(3-methoxy-benzyl)piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl)-acetic acid ethyl ester (5.0 g, 22 mmol) and bis- (2-chloro ethyl)-(3-methoxy-benzyl)-amine (8.0 g, 23.5 mmol). Yield 2.4 g (24%); low melting solid; MS: 447.9 (M+H)⁺.
- 20 4-(4-Methoxy-benzenesulfonyl)1-(3-methoxy-benzyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-Methoxy-benzenesulfonyl)-1-(3-methoxy-benzyl)-piperidine-4-carboxylic acid ethyl ester (2.4g, 5.36 mmol) dissolve in methanol (30 mL) , 10 N sodium hydroxide (10 mL), tetrahydrohydrofuran (20 mL). The resulting
- 25 reaction mixture was worked up as outlined in example 83. Yield 710 mg (32%). white solid mp 199 °C , MS: 419.9 (M+H)⁺.
- Starting from 4-(4-methoxy-benzenesulfonyl)-1-(3-methoxy-benzyl)-piperidine-4-carboxylic acid (830 mg, 1.98 mmol) and following the procedure as outlined in
- 30 example 83, 190 mg of 4-(4-methoxy-benzenesulfonyl)-1-(3-methoxy-benzyl)-piperidine-4-carboxylic acid hydroxamide was isolated as a white solid. mp 130 °C; Yield 20.4%; MS: 435.0 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 2.24-2.32 (m, 2H), 2.51(d, 2H), 2.73-2.83 (m, 2H), 3.37 (d, 2H), 3.76 (s, 3H), 3.88 (s, 3H), 4.32 (s, 2H), 7.01-7.77 (m,8H), 9.38 (s, 1H), 10.1 (s, 1H).
- 35

WO 99/42436

PCT/US98/17633

- 86 -

Example 85

1-(3,4-dichlorobenzyl) -4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid
hydroxamide

5 2-[(2-Hydroxy-ethyl)-(3,4-dichloro-benzyl)-amino]-ethanol was prepared according to the general method as outlined in example 83. Starting from diethanolamine (4.84 g, 46 mmol) and 3,4-dichlorobenzyl chloride (9.0 g, 46 mmol). Yield 13.8 g (99 %); colorless oil; MS: 264.3 (M+H)⁺.

10 3,4-Dichlorobenzyl-bis-(2-chloro-ethyl)-amine was prepared according to the general method as outlined in example 83. Starting from 3,4-dichlorobenzyl diethanolamine (10.7 g, 41 mmol). Yield 99%; yellow solid mp 218-220 °C; MS: 301.8 (M+H)⁺.

15 1-(3,4-Dichloro-benzyl)-4-(methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl)-acetic acid ethyl ester (2.9 g, 11 mmol) and 3,4-dichlorobenzyl-bis(2-chloroethyl)-amine (3.4 g, 11 mmol). Yield 5.9g (60 %); brown oil; MS: 494.5 (M+H)⁺.

20 1-(3,4-Dichloro-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-(3,4-dichloro-benzyl)-4-(methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (5.0 g, 10 mmol) dissolved in methanol (50 mL), 10 N sodium hydroxide (15 mL) and tetrahydrofuran (75 mL). The resulting
25 reaction mixture was worked up as outlined in example 83. Yield 2.94 g (62 %), MS: 458.3 (M+H)⁺.

Starting from 1-(3,4-dichlorobenzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid (2.67g, 5.8 mmol) and following the procedure as outlined in example
30 83, .2 g of 1-(3,4-dichlorobenzyl) -4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxamide was isolated as a white solid. mp 192-195 °C; Yield 10%;
MS 472.9 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 2.20-2.28 (m, 2H), 2.76-2.79 (m, 2H), 3.43-3.44 (m, 4H), 4.30 (s, 2H), 7.14-7.17 (d, J=.030, 2H), 7.50-7.73 (d, J=.027, 1H), 7.65-7.68 (d, J=.029, 2H), 7.72-7.75 (d, J=.027, 2H), 7.87 (s, 1H),
35 9.37 (s, 1H), 10.53 (s, 1H), 11.18 (s, 1H).

WO 99/42436

PCT/US98/17633

- 87 -

Example 86

4-(4-methoxy-benzenesulfonyl)-1-(4-methylbenzyl)-piperidine-4-carboxylic acid
hydroxamide

5

2-[(2-Hydroxy-ethyl)-(4-methyl-benzyl)- amino]-ethanol was prepared according to the general method as outlined in example 83. Starting from diethanolamine (4.8 g, 46 mmol) and 4-methylbenzyl chloride (8.5 g, 46 mmol). Yield 9.8 g (99 %); MS: 209.9 (M+H)⁺.

10

4-Methylbenzyl-bis-(2-chloro-ethyl)-amine was prepared according to the general method as outlined in example 83. Starting from 4-methyl-benzyl diethanolamine (6 g, 20 mmol). Yield 5.2 g (84 %); yellow solid mp 145-147 °C; MS: 245.9 (M+H)⁺.

15

4-(4-Methoxy-benzenesulfonyl)-1-(4-methyl-benzyl)piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl)-acetic acid ethyl ester (7.0 g, 27 mmol) and 4-methyl-bis-(2-chloro-ethyl)-amine (5.0 g, 17 mmol). Yield 4.64 g (63 %); low melting solid; MS: 431.9 (M+H)⁺.

20

4-(4-Methoxy-benzenesulfonyl)-1-(4-methyl-benzyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (4.3g, 9.9 mmol) dissolve in methanol (30 mL) , 10 N sodium hydroxide (10 mL), tetrahydrofuran (20 mL). The resulting reaction mixture was worked up as outlined in example 83. Yield 1.6 g (40 %). white solid mp 207-208 °C , MS: 404.3 (M+H)⁺.

25

Starting from 4-(4-methoxy-benzenesulfonyl)-1-(4-methylbenzyl)-piperidine-4-carboxylic acid (1.59g, 3.9 mmol) and following the procedure as outlined in example 83, .505 g of 4-(4-methoxy-benzenesulfonyl)-1-(4-methylbenzyl)-piperidine-4-carboxylic acid hydroxamide was isolated as a white solid. mp 176-177 °C; Yield 32%; MS: 419.0 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 2.24-2.32 (m, 2H), 2.51(t, 3H), 2.73-2.80 (m, 2H), 3.35-3.50 (m, 4H), 3.87 (s, 3H), 4.24 (s, 2H), 7.13-7.17 (d, J=.039, 2H), 7.23-7.60 (d, J=.036, 2H), 7.38-7.41 (d, J=.025, 2H), 7.65-7.68 (d, J=.039, 2H).

35

WO 99/42436

PCT/US98/17633

- 88 -

Example 87

4-(4-methoxy-benzene-sulfonyl)-1-napthalene-2-yl-methylpiperidine-4-carboxylic acid
hydroxamide

- 5 2-[(2-Hydroxy-ethyl)-(2-naphthyl-2-ylmethyl)-amino]-ethanol was prepared according to the general method as outlined in example 83. Starting from diethanolamine (6.18 g, 59 mmol) and 2-(bromomethyl)naphthalene (10 g, 45 mmol). Yield 12.7 g (96 %); yellow solid mp 162-164 °C; MS: 246.0 (M+H)⁺.
- 10 2-Naphthyl-2-ylmethyl-bis-(2-chloro-ethyl)-amine was prepared according to the general method as outlined in example 83. Starting from 2-naphthyl-ylmethyl-diethanol amine (10 g, 36 mmol). Yield 9.1 g (79 %); brown solid mp 124-126 °C; MS: 281.9 (M+H)⁺.
- 15 4-(4-Methoxy-benzenesulfonyl)-napthalene-ylmethyl-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl)-acetic acid ethyl ester (8.4 g, 32 mmol) and 1-napthalene-ylmethyl-bis-(2-chloro-ethyl)-amine ((8.6 g, 27 mmol). Yield 6.5 g
- 20 (52%); low melting solid; MS: 440.0 (M+H)⁺.
- 4-(4-Methoxy-benzenesulfonyl)-1-napthalene-ylmethyl-piperidine-4-carboxylic acid was prepared starting from 4-(4-methoxy-benzenesulfonyl)-napthalene-ylmethyl-piperidine-4-carboxylic acid ethyl ester (6.3g, 13 mmol) dissolved in methanol (30
- 25 mL), 10 N sodium hydroxide (30 mL) and tetrahydrofuran (30 mL). The resulting reaction mixture was worked up as outlined in example 83. Yield 2.3 g (36 %). yellow solid mp 226-228 °C, MS: 440.0 (M+H)⁺.
- Starting from 4-(4-methoxy-benzenesulfonyl)-1-napthalene-2-yl-methylpiperidine-4-carboxylic acid (2.18g, 5.0 mmol) and following the procedure as outlined in example
- 30 83, 753 g of 4-(4-methoxy-benzene-sulfonyl)-1-napthalene-2-yl-methylpiperidine-4-carboxylic acid hydroxamide was isolated as a off white solid. mp 168-170 °C; Yield 31%; MS 455.0 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 2.29-2.33 (m, 2H), 2.86-2.89 (m, 2H), 3.42-3.46 (m, 4H), 3.85 (s, 3H), 4.46 (s, 2H), 7.13-7.16 (d, J=.030, 2H), 7.56-7.64 (m, 3H), 7.65-7.68 (d, J=.030, 2H), 7.98-8.00 (m, 3H), 8.21 (s,
- 35 1H), 10.70 (s, 1H), 11.20 (s, 1H).

WO 99/42436

PCT/US98/17633

- 89 -

Example 88

1-Biphenyl-4-ylmethyl-4-(4-methoxy-benzenesulfonyl)piperidine-4-carboxylic acid hydroxamide

5 2-[(2-Hydroxy-ethyl)-(1-biphenyl-4-ylmethyl)]-amino]-ethanol was prepared according to the general method as outlined in example 83. Starting from diethanol amine (5.2 g, 49 mmol) and 4-(chloromethyl)biphenyl (10 g, 49 mmol). Yield 9.98 g (66 %); white solid mp 160-162 °C; MS: 271.9 (M+H)⁺. This was converted to the dichloride as
10 outlined in example 83

1-Biphenyl-4-ylmethyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl)-acetic acid ethyl ester (2.85 g, 11 mmol)
15 and 1-biphenyl-4-ylmethyl-bis-(2-chloro-ethyl)-amine (3.4 g, 11 mmol). Yield 2.1 g, (39 %); beige solid, mp 176-178 °C, MS: 494.1 (M+H)⁺.

1-Biphenyl-4-ylmethyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-biphenyl-4-ylmethyl-(4-methoxy-benzenesulfonyl)-
20 piperidine-4-carboxylic acid ethyl ester (5.7g, 12 mmol) dissolved in ethanol (20 mL), tetrahydrofuran (20 mL) and 10 N sodium hydroxide (10 mL). The resulting reaction mixture was worked up as outlined in example 83. Yield 2.1g (39%) MS: 465.8 (M+H)⁺.

25 Starting from 1-biphenyl-4-ylmethyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid (1.0g, 2.2 mmol and following the procedure as outlined in example 83, .132g of 1-biphenyl-4-ylmethyl-4-(4-methoxy-benzenesulfonyl)piperidine-4-carboxylic acid hydroxamide was isolated as a tan solid. mp 168 °C; Yield 20%; MS:
440.9 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 2.30-2.35 (m, 2H), 2.83-2.87 (m,
30 2H), 3.35-3.5 (m, 4H), 3.87 (s, 3H), 7.15-7.721 (d, J=.059 Hz, 2H), 7.49-7.65 (m, 5H), 7.68-7.74 (d, J=.06 Hz, 2H), 9.3 (s, 1H), 10.3 (s, 1H), 11.15 (s, 1H).

WO 99/42436

PCT/US98/17633

- 90 -

Example 89

4-(4-methoxy-benzene-sulfonyl)-1-(3-methyl-but-2-enyl)piperidine-4-carboxylic acid hydroxamide

5 2-[(2-Hydroxy-ethyl)-1-(3-methyl-but-2-enyl)-amino]-ethanol was prepared according to the general method as outlined in example 83. Starting from diethanol amine (4.1 g, 39 mmol) and 4-bromo-2-methyl-butene (6.0 g, 40 mmol). Yield (98 %); brown oil; MS: 173.8 (M+H)⁺.

10 1-(3-methyl-but-2-enyl)-bis-(2-chloro-ethyl)-amine was prepared according to the general method as outlined in example 83. Starting from 2-[(2-hydroxy-ethyl)-1-(3-methyl-but-2-enyl)-amino]-ethanol (10.4g, 50 mmol). Yield 10.5g (99%); brown solid; MS: 210.3 (M+H)

15 4-(4-Methoxy-benzenesulfonyl)-1-(3-methyl-but-2-enyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 1. Starting from 4-(methoxy-benzenesulfonyl)-acetic acid ethyl ester (11.32 g, 44 mmol) and 3-methyl-but-2-enyl)-bis-(2-chloroethyl)-amine (10.4 g, 50 mmol). Yield 6.2 g (36 %); brown oil; MS: 395.6 (M+H)⁺.

20 4-(4-Methoxy-benzenesulfonyl)-1-(3-methyl-but-2-enyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-methoxy-benzenesulfonyl)-1-(3-methyl-but-2-enyl)-piperidine-4-carboxylic acid ethyl ester (6.2g, 16 mmol) dissolved in ethanol (15 mL), 10 N sodium hydroxide (10 mL) and tetrahydrofuran (75 mL). The resulting reaction mixture was worked up as outlined in example 83. Yield 1.2 g (21 %). brown solid mp 196-197 °C, MS: 367.9 (M+H)⁺.

30 Starting from 4-(4-methoxy-benzenesulfonyl)-1-(3-methyl-but-2-enyl)-piperidine-4-carboxylic acid (1.0g, 3.0 mmol) and following the procedure as outlined in example 83, 110 mg of 4-(4-methoxy-benzene-sulfonyl)-1-(3-methyl-but-2-enyl)piperidine-4-carboxylic acid hydroxamide was isolated as a yellow solid. mp 142-145 °C; Yield 12%; MS: 382.9 (M+H)⁺ ; ¹H NMR (300 MHz, DMSO-d₆): δ 1.67 (s, 3H), 1.79 (s, 3H), 2.18-2.23 (m, 2H), 2.66-2.73 (m, 2 H), 3.37-3.46 (m, 2H), 3.67-3.69 (m, 2H), 5.19-5.24 (m, 1H), 7.15-7.18 (d, J=.03, 2H), 7.67-7.70 (d, J=.030, 2H), 9.34 (s, 1H), 9.88 (s, 1H), 11.15 (s, 1H).

WO 99/42436

PCT/US98/17633

- 91 -

Example 90

1-(4-Bromo-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid
hydroxyamide

- 5 2-[(4-Bromobenzyl)-(2-hydroxy-ethyl)-amino]-ethanol was prepared according to the general method as outlined in example 83. Starting from diethanolamine (22.5 g, 150 mmol), and 4-bromobenzyl bromide (25 g, 100 mmol). Yield 33.66g, (99%); yellow oil; MS: 273.8 (M+H)⁺.
- 10 (4-Bromo-benzyl)-bis-(2-chloro-ethyl)-amine was prepared according to the general method as outlined in example 83. Starting from 2-[(4-bromobenzyl)-(2-hydroxy-ethyl)-amino]-ethanol (33.28 g, 122 mmol). Yield 47 g, (99%); brown solid; mp 125°C; MS: 309.8 (M+H)⁺.
- 15 1-(4-Bromo-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl) acetic acid ethyl ester (8.6 g, 33.5 mmol) and (4-bromo-benzyl)-bis-(2-chloro-ethyl)-amine (13.3 g, 38.6 mmol). Yield 17 g (44%);
- 20 brown oil; MS: 497.8 (M+H)⁺.
- 1-(4-Bromo-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-(4-bromo-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (16.5 g, 33.3 mmol) dissolved in THF:methanol 3:1 and
- 25 10 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 6.18 g (40%); tan solid; mp 215 °C; MS: 469.7 (M+H)⁺.
- Starting from 1-(4-Bromo-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid (1.95 g, 4.2 mmol) and following the procedure as outlined in example
- 30 83, 1.29 g of 1-(4-bromo-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as an off white solid. Yield 60%; mp 180°C; MS: 484.7 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 2.18-2.29 (m, 2H), 2.46 (d, 2H), 2.74-2.89 (m, 2H), 3.39 (d, 2H), 3.87 (s, 3H), 4.28 (s, 2H), 7.18 (d, J = 17 Hz, 2H), 7.49 (d, J = 8.1 Hz, 2H), 7.65- 7.68 (m, 4H), 9.37 (s, 1H), 10.5 (s,
- 35 1H).

WO 99/42436

PCT/US98/17633

- 92 -

Example 91

4-(4-methoxy-benzenesulfonyl)-1-(3-phenyl-propyl)-piperidine-4-carboxylic acid
hydroxyamide

- 5 2-[(2-Hydroxy-ethyl)-(3-phenyl-propyl)-amino]-ethanol was prepared according to the general method as outlined in example 83. Starting from diethanolamine (15.8 g, 151 mmol), and 1-bromo-3-phenylpropane (20 g, 101 mmol). Yield 21.31 g, (95%); yellow oil; MS: 223.9 (M+H)⁺.
- 10 Bis-(2-Chloro-ethyl)-(3-phenyl-propyl)-amine was prepared according to the general method as outlined in example 83. Starting from 2-[(2-hydroxy-ethyl)-(3-phenyl-propyl)-amino]-ethanol (20.32 g, 90.7 mmol). Yield 24.9 g (92%); brown oil; MS: 259.8 (M+H)⁺.
- 15 4-(4-Methoxy-benzenesulfonyl)-1-(3-phenyl-propyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl) acetic acid ethyl ester (12 g, 46.5 mmol) and bis-(2-chloro-ethyl)-(3-phenyl-propyl)-amine (24.8 g, 93.8 mmol). Yield 11.24 g
- 20 (54%); brown oil; MS: 446 (M+H)⁺.
- 4-(4-Methoxy-benzenesulfonyl)-1-(3-phenyl-propyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-Methoxy-benzenesulfonyl)-1-(3-phenyl-propyl)-piperidine-4-carboxylic acid ethyl ester (10.74 g, 24.13 mmol) dissolved in THF:methanol 3:1
- 25 and 10 N NaOH (40 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 4.67 g (47%); off white powder; mp 203 °C; MS: 418.2 (M+H)⁺.
- Starting from 4-(4-methoxy-benzenesulfonyl)-1-(3-phenyl-propyl)-piperidine-4-carboxylic acid (4.37 g, 10.4 mmol) and following the procedure as outlined in
- 30 example 83, 1.64 g of 4-(4-methoxy-benzenesulfonyl)-1-(3-phenyl-propyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as an off white solid. Yield 37%; mp 143°C; MS: 432.9 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 1.92-1.97 (m, 2H), 2.18-2.29 (m, 2H), 2.47 (d, 2H), 2.58 (t, J = 7.7 Hz, 2H), 2.6-2.73 (m, 2H), 3.0-3.06 (m, 2H), 3.60 (d, J = 12.3 Hz, 2H), 3.87 (s, 2H), 7.15-7.30 (m, 7 H), 7.68,
- 35 (d, J = 9 Hz, 2H), 9.3 (s, 1H), 10.1 (s, 1H).

WO 99/42436

PCT/US98/17633

- 93 -

Example 92

1-Tert-butyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid
hydroxyamide

5

tert-Butyl-bis-(2-chloro-ethyl)-amine was prepared according to the general method as outlined in example 83. Starting from 1-tert-butyl-diethanolamine (6 g, 37.2 mmol). Yield 11.15 g, (99%); white solid; MS: 197.8 (M+H)⁺.

10

1-tert-Butyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl) acetic acid ethyl ester (10 g, 38.76 mmol) and tert-butyl-bis-(2-chloro-ethyl)-amine (5.25 g, 22.53 mmol). Yield 5.37 g, (62%); brown oil; MS: 384 (M+H)⁺.

15

1-tert-Butyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-tert-butyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (5.37 g 14 mmol) dissolved in methanol (300 ml) and 10 N NaOH (23 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 1.52 g (30.6%); white powder; mp 204 °C; MS: 356 (M+H)⁺.

20

Starting from 1-tert-butyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid (320 mg, 0.9 mmol) and following the procedure as outlined in example 83, 190 mg of 1-tert-butyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as a green solid. Yield 52%; mp 40 °C; MS: 371.1 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 1.29 (s, 9H), 1.54 (m, 2H), 1.66 (m, 2H), 2.39 (m, 2H), 2.98 (m, 2H), 3.88 (s, 3H), 7.18 (d, 2H), 7.67 (d, 2H).

25

Example 93

30

1-Butyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide

Butyl-bis-(2-chloro-ethyl)-amine was prepared according to the general method as outlined in example 83. Starting from N-butyl-diethanolamine (6 g, 37.2 mmol). Yield 11.3 g, (99%); white powder; mp 165 °C; MS: 197.9 (M+H)⁺.

35

1-Butyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl) acetic acid ethyl ester (5 g, 19.38 mmol) and butyl-bis-(2-

WO 99/42436

PCT/US98/17633

- 94 -

chloro-ethyl)-amine (4.52 g, 19.38 mmol). Yield 6.86 g, (93%); brown oil; MS: 384 (M+H)⁺.

5 1-Butyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-butyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (6.42 g 16.8 mmol) dissolved in methanol (200 ml) and 10 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 1.6 g (27%); white powder; mp 206 °C; MS: 356.4 (M+H)⁺.

10 Starting from 1-butyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid (1.51 g, 4.3 mmol) and following the procedure as outlined in example 83, 200 mg of 1-butyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as an off white solid. Yield 9.3%; mp 75 °C; MS: 371.1 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.87 (t, J = 7.2 Hz, 3H), 1.27 (m, 2H), 1.59 (m, 2H), 2.27 (m, 2H), 2.45 (m, 2H), 2.50 (m, 2H), 2.65 (m, 2H), 2.97 (m, 2H) 3.88 (s, 3H), 7.18 (d, 2H), 7.69 (d, 2H).

Example 94

20 1-Cyclooctyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide

Cyclooctyl-bis-(2-chloro-ethyl)-amine was prepared according to the general method as outlined in example 83. Starting from N-cyclooctyldiethanolamine (6 g, 28 mmol).
25 Yield 10 g, (99%); off white solid; mp 158 °C; MS: 251.9 (M+H)⁺.

1-Cyclooctyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl) acetic acid ethyl ester (5 g, 19.4 mmol) and cyclooctyl-bis-(2-chloro-ethyl)-amine (5.57 g, 19.4 mmol). Yield 8.2 g, (96%); brown oil; MS: 438 (M+H)⁺.

1-Cyclooctyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-cyclooctyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (8 g, 18.3 mmol) dissolved in methanol (200 ml) and 10 N NaOH (25 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 2.36 g (32%); white powder; mp 180 °C; MS: 410 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 95 -

Starting from 1-Cyclooctyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid (2.26 g, 5.53 mmol) and following the procedure as outlined in example 83, 570 mg of 1-cyclooctyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as a white powder. Yield 22%; mp >200 °C; MS: 425

- 5 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 1.42-1.66 (m, 14H), 1.83 (m, 2H), 2.33 (m, 2H), 2.67 (m, 2H), 3.30-3.51 (m, 3H) 3.88 (s, 3H) 7.17 (d, 2H), 7.66 (d, 2H).

Example 95

- 10 1-Ethyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide

- 1-Ethyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl) acetic acid ethyl ester (3 g, 11.6 mmol) and ethyl-bis-(2-chloro-ethyl)-amine (2.39g, 11.6 mmol). Yield 3.09 g, (75%); low melting brown solid; MS: 356 (M+H)⁺.

- 20 1-Ethyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-ethyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (2.42 g, 6.8 mmol) dissolved in methanol (100 ml) and 10 N NaOH (15 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 1.29 g (58%); white solid; mp 209 °C; MS: 328 (M+H)⁺.

- 25 Starting from 1-ethyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid (1.23 g, 3.76 mmol) and following the procedure as outlined in example 83, 1.02 g of 1-ethyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as an off white powder. Yield 80%; mp 85 °C; MS: 343 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.926 (t, J = 7.1 Hz, 3H), 1.68-1.89 (m, 4H), 2.05-2.24 (m, 4H), 2.73 (q, 2H), 3.85 (s, 3H), 7.07 (d, 2H), 7.64 (d, 2H).

30

Example 96

1-Isopropyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide

35

1-Isopropyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl) acetic acid ethyl ester (5.7 g, 22.2 mmol) and isopropyl-

WO 99/42436

PCT/US98/17633

- 96 -

bis-(2-chloro-ethyl)-amine (4.9 g, 22.2 mmol). Yield 5.64 g, (68%); low melting brown solid; MS: 370 (M+H)⁺.

1-Isopropyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-isopropyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (5.6 g, 15.2 mmol) dissolved in methanol (75 ml) and 10 N NaOH (25 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 2.18 g (42%); white powder; mp 204 °C; MS: 341.9 (M+H)⁺.

Starting from 1-isopropyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid (2.13 g, 6.25 mmol) and following the procedure as outlined in example 83, 590 mg of 1-isopropyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as a white powder. Yield 2.4%; mp 75 °C; MS: 357 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 1.21 (d, J = 6.6 Hz, 6H), 2.33-3.53 (m, 9H), 3.88 (s, 3H), 7.16 (d, 2H), 7.66 (d, 2H).

Example 97

1-Methyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide
1-Methyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl) acetic acid ethyl ester (3 g, 11.6 mmol) and methyl-bis-(2-chloro-ethyl)-amine (2.2g, 11.6 mmol). Yield 3.09 g, (75%); low melting brown solid; MS: 342 (M+H)⁺.

1-Methyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-methyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (8.7 g, 25.6 mmol) dissolved in methanol (300 ml) and 10 N NaOH (35 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 3.23 g (41%); white solid; mp 204 °C; MS: 313.9 (M+H)⁺.

Starting from 1-methyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid (2.0 g, 6.38 mmol) and following the procedure as outlined in example 83, 1.10 g of 1-methyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as a yellow powder. Yield 53%; mp 89 °C; MS: 329 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 1.67- 1.76 (m, 2H), 1.85-1.96 (m, 2H), 2.05 (s, 3H), 2.17

WO 99/42436

PCT/US98/17633

- 97 -

(d, J = 11.4 Hz, 2H), 2.57 (d, J = 10.4 Hz, 2H) 3.83 (s, 3H), 7.02 (d, 2H), 7.62 (d, 2H).

Example 98

5

1-Benzyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide

1-Benzyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from
10 from 4-(butoxy-benzenesulfonyl) acetic acid ethyl ester (6 g, 20 mmol) and bis-(2-chloro-ethyl)-benzylamine (10 g, 30 mmol). Yield 5.15 g (56%); yellow oil; MS: 460 (M+H)⁺.

1-Benzyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared
15 starting from 1-benzyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (5.1 g, 11.1 mmol) dissolved in THF:methanol 3:1 and 10 N NaOH (10 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 2.66 g (56%); off white solid; mp 210 °C; MS: 432 (M+H)⁺.

20 Starting from 1-benzyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid (2.61 g, 6.06 mmol) and following the procedure as outlined in example 83, 860 mg of 1-benzyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as an off white powder. Yield 32%; mp 144 °C; MS: 446.9 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.94 (t, J = 7.3 Hz, 3H), 1.44 (q, J = 7.5 Hz, 2H),
25 1.70 (q, 2H), 2.28-2.32 (m, 2H), 2.50 (d, 2H), 2.74-2.83 (m, 2H), 3.35 (d, 2H), 4.08 (t, J = 6.3 Hz, 2H), 4.34 (s, 2H), 7.13 (d, J = 8.7, 2H), 7.45 (s, 3H), 7.54 (s, 2H), 7.74 (d, J = 8.7, 2H), 9.35 (s, 1H), 10.7 (s, 1H).

Example 99

30

1-(4-Fluoro-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide

1-(4-Fluoro-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl
35 ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl) acetic acid ethyl ester (18.8 g, 72.8 mmol) and (4-fluoro-benzyl)-bis-(2-chloro-ethyl)-amine (20.8 g, 73 mmol). Yield 25 g (79%); brown oil; MS: 436.9 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 98 -

1-(4-Fluoro-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-(4-fluoro-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (17.4 g, 40 mmol) dissolved in THF:methanol 3:1 and 10 N NaOH (40 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 10.8 g (66%); colorless solid; mp 154 °C; MS: 408 (M+H)⁺.

Starting from 1-(4-Fluoro-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid (8.14 g, 20 mmol) and following the procedure as outlined in example 83, 4.3 g of 1-(4-fluoro-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as an off white solid. Yield 51%; mp 176-178 °C; MS: 484.7 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 2.12-2.20 (m, 2H), 2.64-2.79 (m, 2H), 3.32-3.45 (m, 4H), 3.87 (s, 3H), 4.31 (s, 2H), 7.14-7.19 (d, J = 17 Hz, 2H), 7.27-7.33 (d, J = 8.1 Hz, 2H), 7.50-7.54 (d, 2H), 7.65-7.68 (d, 2H), 9.38 (s, 1H), 9.75 (s, 1H).

Example 100

1-(4-Fluoro-benzyl)-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide

1-(4-Fluoro-benzyl)-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(butoxy-benzenesulfonyl) acetic acid ethyl ester (6 g, 20 mmol) and (4-fluoro-benzyl)-bis-(2-chloro-ethyl)-amine (5.73 g, 20 mmol). Yield 8.2 g (86%); yellow oil; MS: 478 (M+H)⁺.

1-(4-Fluoro-benzyl)-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-(4-Fluoro-benzyl)-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (4.77 g, 10 mmol) dissolved in THF:methanol 3:1 and 10 N NaOH (10 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 3.5 g (79%); off white solid; mp 114 °C; MS: 450 (M+H)⁺.

Starting from 1-(4-Fluoro-benzyl)-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid (2.24 g, 5.0 mmol) and following the procedure as outlined in example 83, 200 mg of 1-(4-Fluoro-benzyl)-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as an off white powder. Yield 9%; mp 112 °C; MS: 465.9 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.94 (t, J = 7.3 Hz, 3H), 1.35-1.50 (m, 2H), 1.68-1.77 (m, 2H), 2.20-2.28 (m, 2H), 2.66-2.77 (m, 2H),

WO 99/42436

PCT/US98/17633

- 99 -

3.77-3.78 (m, 4H), 4.06-4.10 (m, 2H), 4.19 (s, 2H), 7.14-7.19 (d, J = 8.7, 2H), 7.27-7.33 (d, 2H), 7.50-7.54 (d, 2H), 7.65-7.68 (d, 2H), 9.34 (s, 1H), 10.55 (s, 1H).

5

Example 101

4-(4-methoxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid hydroxyamide

10 2-[(2-Hydroxy-ethyl)-(4-methoxy-benzyl)-amino]-ethanol was prepared according to the general method as outlined in example 83. Starting from diethanolamine (12.0 g, 114 mmol) and 4-methoxybenzyl chloride (14.2 g, 100 mmol). Yield 17.5 g, (77 %); yellow oil; MS: 226 (M+H).

15 4-Methoxybenzyl-bis-(2-chloro-ethyl)-amine was prepared according to the general method as outlined in example 83. Starting from 4-Methoxy-benzyl diethanolamine (10 g, 44 mmol). Yield 10 g (75 %); yellow solid mp 55 °C; MS: 263.1 (M+H)⁺.

20 4-(4-Methoxy-benzenesulfonyl)-1-(4-methoxy-benzyl)piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl)-acetic acid ethyl ester (5.0 g, 20 mmol) and bis-(2-chloro ethyl)-(4-methoxy-benzyl)-amine (7.0 g, 22 mmol). Yield 5.0 g (56 %); low melting solid; MS: 448.5 (M+H)⁺.

25 4-(4-Methoxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-Methoxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid ethyl ester (4.2g, 10 mmol) dissolve in methanol (30 mL) , 10 N sodium hydroxide (10 mL), tetrahydrofuran (20 mL). The resulting reaction mixture was worked up as outlined in example 83. Yield 3.0 g (71 %). white solid mp 30 190 °C , MS: 420.4 (M+H)⁺.

Starting from 4-(4-methoxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid (2.0 g, 4.7 mmol) and following the procedure as outlined in example 83, 1.2 g of 4-(4-methoxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-

35 carboxylic acid hydroxamide was isolated as a white solid. mp 175 °C (HCl); Yield:

1.2 g, 59 %; MS: 433.0 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 1.8 (m, 4H), 2.3(m, 2H), 2.73 (m, 2H), 3.37 (d, 2H), 3.76 (s, 3H), 3.88 (s,3H), 6.87 (d, 2H), 7.11 (d, 2H), 7.21 (d, 2H), 7.65 (d, 2H), 9.2 (bs, 1H), 10.9 (bs, 1H).

WO 99/42436

PCT/US98/17633

- 100 -

Example 102

4-(4-methoxy-benzenesulfonyl)-1-[2-(4-methoxyphenyl)-ethyl]-piperidine-4-carboxylic
acid hydroxyamide

5

2-[(2-Hydroxy-ethyl)-[2-(4-methoxy-phenyl)-ethyl]-amino]-ethanol was prepared according to the general method as outlined in example 83. Starting from diethanolamine (10.0 g, excess), and 1-(2-chloroethyl)-4-methoxybenzene (8.5 g, 50 mmol).

Yield 11 g, (92%); yellow oil; MS: 240 (M+H)⁺.

10

The corresponding dichloride, bis-(2-chloro-ethyl)-(4-methoxyphenyl-2-ethyl)-amine was prepared according to the general method as outlined in example 83. Starting from 2-[(2-hydroxy-ethyl)-[2-(4-methoxy-phenyl)-ethyl]-amino]-ethanol (10 g, 41.8 mmol). Yield 11 g (95%); brown oil; MS: 277.2 (M+H)⁺.

15

4-(4-methoxy-benzenesulfonyl)-1-[2-(4-methoxyphenyl)-ethyl]-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl) acetic acid ethyl ester (5.0 g, 20 mmol) and bis-(2-chloro-ethyl)-(4-methoxyphenyl-2-ethyl)-amine (6.4 g, 20 mmol). Yield 6.0 g (65%); brown oil; MS: 462.5 (M+H)⁺.

20

4-(4-methoxy-benzenesulfonyl)-1-[2-(4-methoxyphenyl)-ethyl]-piperidine-4-carboxylic acid was prepared starting from 4-(4-methoxy-benzenesulfonyl)-1-[2-(4-methoxy-phenyl)-ethyl]-piperidine-4-carboxylic acid ethyl ester (5.0 g, 10.8 mmol) dissolved in THF:methanol 3:1 and 10 N NaOH (40 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 4.0 g (85%); off white powder; mp 205°C; MS: 434.5 (M+H)⁺.

25

Starting from 4-(4-methoxy-benzenesulfonyl)-1-[2-(4-methoxyphenyl)-ethyl]-piperidine-4-carboxylic acid (1.5 g, 3.46 mmol) and following the procedure as outlined in example 83, 900 mg of 4-(4-methoxy-benzenesulfonyl)-1-[2-(4-methoxyphenyl)-ethyl]-piperidine-4-carboxylic acid hydroxyamide was isolated as an off white solid. Yield 58%; mp 206 °C (HCl); MS: 449.5 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 2.3 (m, 2H), 2.5 (m, 3H), 2.8 (m, 2H), 2.95 (m, 2H), 3.25 (m, 2H), 3.4 (m, 4H), 3.60 (d, J = 12.3 Hz, 2H), 3.77 (s, 3H), 3.99 (s, 3H), 6.9 (d, 2H), 7.1 - 7.25, (q, 4H), 7.7 (d, 2H), 9.3 (s, 1H), 10.6 (s, 1H).

30

35

WO 99/42436

PCT/US98/17633

- 101 -

Example 103

4-(4-methoxy-benzenesulfonyl)-1-(2-phenyl-ethyl)-piperidine-4-carboxylic acid
hydroxyamide

5 2-[(2-Hydroxy-ethyl)-(2-phenyl-ethyl)-amino]-ethanol was prepared according to the
general method as outlined in example 1. Starting from diethanolamine (6.0 g, 57).
and 2-bromo-ethylbenzene (9.0 g, 48.3 mmol). Yield 9 g, (90%); yellow oil; MS: 210
(M+H)⁺.

10 Bis-(2-Chloro-ethyl)-(2-phenyl-ethyl)-amine was prepared according to the general
method as outlined in example 83. Starting from 2-[(2-Hydroxy-ethyl)-(2-phenyl-
ethyl)-amino]-ethanol (8.5 g, 40.6 mmol). Yield 11 g (95%); brown oil; MS: 247.1
(M+H)⁺.

15 4-(4-methoxy-benzenesulfonyl)-1-(2-phenyl-ethyl)-piperidine-4-carboxylic acid
ethyl ester was prepared according to the general method as outlined in example 83.
Starting from 4-(methoxy-benzenesulfonyl) acetic acid ethyl ester (5.0 g, 20
mmol) and bis-(2-chloro-ethyl)-(2-phenyl-ethyl)-amine (5.6 g, 20 mmol). Yield 5.5 g
20 (63%); brown oil; MS: 432.5 (M+H)⁺.

4-(4-methoxy-benzenesulfonyl)-1-(2-phenyl-ethyl)-piperidine-4-carboxylic acid was
prepared starting from 4-(4-methoxy-benzenesulfonyl)-1-(2-phenyl-ethyl)-piperidine-4-
carboxylic acid ethyl ester (3.0 g, 6.9 mmol) dissolved in THF:methanol 3:1 and 10 N
25 NaOH (40 ml). The resulting reaction mixture was worked up as outlined in example
83. Yield 2.0 g (72%); off white powder; mp 208 °C; MS: 404.5 (M+H)⁺.

Starting from 4-(4-methoxy-benzenesulfonyl)-1-(2-phenyl-ethyl)-piperidine-4-
carboxylic acid (1.5 g, 3.7 mmol) and following the procedure as outlined in example
30 83, 900 mg of 4-(4-methoxy-benzenesulfonyl)-1-(2-phenyl-ethyl)-piperidine-4-
carboxylic acid hydroxyamide was isolated as an off white solid. Yield 58%; mp
205°C (HCl); MS: 419.4 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 2.3 (m, 2H),
2.5 (m, 3H), 2.8 (m, 2H), 2.95 (m, 2H), 3.25 (m, 2H), 3.4 (m, 4H), 3.9 (s,
3H), 7.22 - 7.8 (m, 9H), 10.6 (s, 1H), 11.2 (bs, 1H).

35

WO 99/42436

PCT/US98/17633

- 102 -

Example 104

4-(4-n-Butoxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid
hydroxyamide

5 4-(4-n-Butoxy-benzenesulfonyl)-1-(4-methoxy-benzyl)piperidine-4-carboxylic acid
ethyl ester was prepared according to the general method as outlined in example 83.
Starting from 4-(n-Butoxy-benzenesulfonyl)-acetic acid ethyl ester (2.5 g, 10 mmol)
and bis- (2-chloro ethyl)-(4-methoxy-benzyl)-amine (3.0 g, 10 mmol). Yield 3.5 g (71
10 %); low melting solid; MS: 490.5 (M+H)⁺.

4-(4-n-Butoxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid
was prepared starting from 4-(4-Butoxy-benzenesulfonyl)-1-(4-methoxy-
benzyl)piperidine-4-carboxylic acid ethyl ester (3.0g, 6.1 mmol) dissolve in methanol
15 (30 mL), 10 N sodium hydroxide (10 mL), tetrahydrohydrofuran (20 mL). The
resulting reaction mixture was worked up as outlined in example 83. Yield 1.5 g
(53%). white solid mp 207 °C, MS: 462.5 (M+H)⁺.

Starting from 4-(4-n-Butoxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-
20 carboxylic acid (1.0 g, 2.1 mmol) and following the procedure as outlined in example
83, 1.2 g of 4-(4-Butoxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-
carboxylic acid hydroxamide was isolated as a white solid. mp 173 °C (HCl); Yield:
800 mg, 77 %; MS: 477.5 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.9 (t, 3H),
1.4 (m, 2H), 1.7 (m, 2H), 2.3 (m, 2H), 2.5 (m, 2H), 2.7 (m, 2H), 3.3 (m, 2H),
25 3.5(m, 2H), 4.1 (t, 2H), 4.3 (m, 2H), 6.97 (d, 2H), 7.14 (d, 2H), 7.48 (d, 2H), 7.7
(d, 2H), 9.4 (bs, 1H), 10.9 (bs, 1H).

Example 105

30 4-(4-Methoxy-benzenesulfonyl)-1-(3-phenoxy-propyl)-piperidine-4-carboxylic acid
hydroxyamide

2-[(2-Hydroxy-ethyl)-(3-phenoxy-propyl)-amino]-ethanol was prepared according to
the general method as outlined in example 83. Starting from diethanolamine (15.8 g,
35 151 mmol). and 3-Phenoxypropyl bromide (21.5 g, 100 mmol). Yield 21.31 g,
(95%); yellow oil; MS: 238.1 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 103 -

Bis-(2-Chloro-ethyl)-(3-phenoxy-propyl)-amine was prepared according to the general method as outlined in example 83. Starting from 2-[(2-hydroxy-ethyl)-(3-phenoxy-propyl)-amino]-ethanol (20.0 g, 84 mmol). Yield 24.0 g (91%); brown oil; MS: 277.8 (M+H)⁺.

5

4-(4-Methoxy-benzenesulfonyl)-1-(3-phenoxy-propyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl) acetic acid ethyl ester (5.2 g, 20 mmol) and bis-(2-chloro-ethyl)-(3-phenoxy-propyl)-amine (7.0 g, 22 mmol). Yield 6.5 g (70%); brown oil; MS: 462.5 (M+H)⁺.

10

4-(4-Methoxy-benzenesulfonyl)-1-(3-phenoxy-propyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-Methoxy-benzenesulfonyl)-1-(3-phenoxy-propyl)-piperidine-4-carboxylic acid ethyl ester (4.2 g, 9.1 mmol) dissolved in THF:Methanol 3:1 and 10 N NaOH (40 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 3.0 g (75%); off white powder; mp 195 °C; MS: 434.5 (M+H)⁺.

15

Starting from 4-(4-methoxy-benzenesulfonyl)-1-(3-phenoxy-propyl)-piperidine-4-carboxylic acid (2.5 g, 5.77 mmol) and following the procedure as outlined in example 83, 1.2 g of 4-(4-methoxy-benzenesulfonyl)-1-(3-phenoxy-propyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as an off white solid. Yield 46%; mp 101 °C; MS: 448.5 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 2.18 (m, 2H), 2.3 (m, 2H), 2.58 (m, 2H), 2.6-2.73 (m, 2H), 3.0-3.06 (m, 2H), 3.60 (m 2H), 3.87 (s, 3H), 4.01 (t, 2H), 6.9 - 7.7 (m, 9H), 9.33 (bs, 1H), 10.28 (bs, 1H).

25

Example 106

4-(4-n-Butoxy-benzenesulfonyl)-1-(3-phenoxy-propyl)-piperidine-4-carboxylic acid hydroxyamide

30

4-(4-n-Butoxy-benzenesulfonyl)-1-(3-phenoxy-propyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(butoxy-benzenesulfonyl) acetic acid ethyl ester (3.0 g, 10 mmol) and bis-(2-chloro-ethyl)-(3-phenoxy-propyl)-amine (3.0 g, 11 mmol). Yield 4.5 g (89%); brown oil; MS: 504.6 (M+H)⁺.

35

WO 99/42436

PCT/US98/17633

- 104 -

4-(4-n-Butoxy-benzenesulfonyl)-1-(3-phenoxy-propyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-n-Butoxy-benzenesulfonyl)-1-(3-phenoxy-propyl)-piperidine-4-carboxylic acid ethyl ester (4.0 g, 7.9 mmol) dissolved in THF:Methanol 3:1 and 10 N NaOH (40 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 3.0 g (79%); off white powder; mp 191 °C; MS: 476.5 (M+H)⁺.

Starting from 4-(4-n-butoxy-benzenesulfonyl)-1-(3-phenoxy-propyl)-piperidine-4-carboxylic acid (700 mg, 1.4 mmol) and following the procedure as outlined in example 83, 300 mg of 4-(4-n-butoxy-benzenesulfonyl)-1-(3-phenoxy-propyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as an off white solid. Yield 43%; mp 84 °C; MS: 491.5 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.9 (t, 3H), 1.5 (m, 2H), 1.8 (m, 2H), 2.18 (m, 2H), 2.3 (m, 2H), 2.58 (m, 2H), 2.6-2.73 (m, 2H), 3.2 (m, 2H), 3.40 (m 6H), 3.97 (t, 2H), 4.1 (t, 2H), 6.9 - 7.7 (m, 9H), 10.7 (bs, 1H), 11.28 (bs, 1H).

Example 107

4-(4-methoxy-benzenesulfonyl)-1-(2-phenoxy-ethyl)-piperidine-4-carboxylic acid hydroxyamide

2-[(2-Hydroxy-ethyl)-(2-phenoxy-ethyl)-amino]-ethanol was prepared according to the general method as outlined in example 83. Starting from diethanolamine (15.0 g, 150) and 2-chloro-phenetol (15.6 g, 100 mmol). Yield 18 g, (80%); Colorless oil; MS: 226 (M+H)⁺.

Bis-(2-Chloro-ethyl)-(2-phenoxy-ethyl)-amine was prepared according to the general method as outlined in example 83. Starting from 2-[(2-Hydroxy-ethyl)-(2-phenoxy-ethyl)-amino]-ethanol (20.0 g, 88.8 mmol). Yield 25 g (94%); brown oil; MS: 263.1 (M+H)⁺.

4-(4-methoxy-benzenesulfonyl)-1-(2-phenoxy-ethyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl) acetic acid ethyl ester (5.0 g, 20 mmol) and bis-(2-chloro-ethyl)-(2-phenoxy-ethyl)-amine (6.0 g, 20 mmol). Yield 5.8 g (64%); brown oil; MS: 448.5 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 105 -

4-(4-methoxy-benzenesulfonyl)-1-(2-phenoxy-ethyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-methoxy-benzenesulfonyl)-1-(2-phenyl-ethoxy)-piperidine-4-carboxylic acid ethyl ester (5.0 g, 11.1 mmol) dissolved in THF:methanol 3:1 and 10 N NaOH (40 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 3.0 g (63%); off white powder; mp 235 °C; MS: 420.5 (M+H)⁺.

Starting from 4-(4-methoxy-benzenesulfonyl)-1-(2-phenoxy-ethyl)-piperidine-4-carboxylic acid (2.5 g, 5.9 mmol) and following the procedure as outlined in example 83, 1.3 g of 4-(4-methoxy-benzenesulfonyl)-1-(2-phenoxy-ethyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as an off white solid. Yield 50%; mp 168-172 °C (HCl); MS: 435.4 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 2.3 (m, 2H), 2.5 (m, 2H), 2.9 (m, 2H), 3.4 (m, 4H), 3.5 (m, 2H), 3.7 (m, 2H), 3.9 (s, 3H), 4.4 (m, 2H), 6.9 - 7.8 (m, 9H), 9.3 (s, 1H), 10.2 (bs, 1H), 11.3 (s, 1H).

15

Example 108

4-(4-n-Butoxy-benzenesulfonyl)-1-(2-phenoxy-ethyl)-piperidine-4-carboxylic acid hydroxyamide

4-(4-Butoxy-benzenesulfonyl)-1-(2-phenoxy-ethyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl) acetic acid ethyl ester (2.5 g, 10 mmol) and bis-(2-chloro-ethyl)-(2-phenoxy-ethyl)-amine (2.98 g, 10 mmol). Yield 3.0 g (69%); brown oil; MS: 490.6 (M+H)⁺.

25

4-(4-n-Butoxy-benzenesulfonyl)-1-(2-phenoxy-ethyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-n-butoxy-benzenesulfonyl)-1-(2-phenyl-ethoxy)-piperidine-4-carboxylic acid ethyl ester (2.5 g, 5.76 mmol) dissolved in THF:methanol 3:1 and 10 N NaOH (40 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 1.5 g (56%); off white powder; mp 204 °C; MS: 462.5 (M+H)⁺.

30

Starting from 4-(4-n-butoxy-benzenesulfonyl)-1-(2-phenoxy-ethyl)-piperidine-4-carboxylic acid (1.0 g, 2.16 mmol) and following the procedure as outlined in example 83, 600 mg of 4-(4-butoxy-benzenesulfonyl)-1-(2-phenoxy-ethyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as an off white solid. Yield 58%; mp 112°C (HCl); MS: 477.4 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.942 (t, 3H),

35

WO 99/42436

PCT/US98/17633

- 106 -

1.4 (m, 2H), 1.7 (m, 2H), 2.3 (m, 2H), 2.5 (m, 4H), 2.8 (m, 2H), 2.9-3.4 (m, 4H), 3.3 (m, 4H), 4.2 (t, 2H), 4.4 (m, 2H), 6.9 - 7.7 (m, 9H), 9.4 (s, 1H), 10.5 (bs, 1H), 11.3 (s, 1H).

5

Example 109

4-(4-Methoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid hydroxyamide

- 10 Bis-(2-chloro-ethyl)-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-amine was prepared according to the general method as outlined in example 83. Starting from diethanolamine (15.0 g, 150), and 4-(2-piperidin-1-yl-ethoxy)-benzyl chloride (5.9 g, 20 mmol). Yield 5.5 g, (85%); Brown semi-solid; MS: 323 (M+H)⁺.
- 15 Bis-(2-chloro-ethyl)-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-amine was prepared according to the general method as outlined in example 83. Starting from 2-[(2-Hydroxy-ethyl)-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-amine (3.22 g, 10 mmol). Yield 4.0 g (92%); brown semi-solid; MS: 361.1 (M+H)⁺.
- 20 4-(4-Methoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 83. Starting from 4-(methoxy-benzenesulfonyl) acetic acid ethyl ester (5.0 g, 20 mmol) and Bis-(2-chloro-ethyl)-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-amine (8.6 g, 20 mmol). Yield 6.0 g (55%); brown oil; MS: 545.7 (M+H)⁺.
- 25 4-(4-Methoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid was prepared starting from 4-(4-Methoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid ethyl ester (5.4 g, 10 mmol) dissolved in THF:methanol 3:1 and 10 N NaOH (40 ml). The resulting reaction
- 30 mixture was worked up as outlined in example 83. Yield 4.0 g (77%); off white powder; mp 174 °C; MS: 517.6 (M+H)⁺.

Starting from 4-(4-Methoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid (3.5 g, 6.78 mmol) and following the procedure as

35 outlined in example 83, 1.8 g of 4-(4-Methoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid hydroxy amide was isolated as a pale yellow solid. Yield 49%; mp 114 °C (HCl); MS: 532 (M+H)⁺; ¹H NMR (300 MHz,

WO 99/42436

PCT/US98/17633

- 107 -

DMSO-d₆: δ 1.4-1.6 (m, 4H), 1.9 (m, 2H), 2.3 (m, 2H), 2.8 (m, 2H), 3.4 (m, 4H), 3.9 (s, 3H), 4.2 (m, 1H), 6.9 - 7.8 (m, 8H), 9.1 (s, 1H), 10.8 (bs, 1H).

Example 110

5

N-Hydroxy-2-(4-methoxy-benzenesulfonyl)-propionamide

Step A: Coupling of 2-bromo-propionic acid to hydroxylamine resin.

4-*O*-Methylhydroxylamine-phenoxymethyl-copoly(styrene-1%-divinylbenzene)-resin¹

10 (2 g, 1.1 meq/g) was placed in a peptide synthesis vessel (Chemglass Inc. Part Number CG-1866) and suspended in DMF (20 mL). 2-Bromopropionic acid (0.6 mL, 3.0 eq.) 1-hydroxybenzotriazole hydrate (HOBt, 1.8 g, 6.0 eq.) and 1,3-diisopropyl-carbodiimide (DIC, 1.4 mL, 4.0 eq.) were added. The reaction was shaken on an orbital shaker at room temperature for 2 - 16 hours. The reaction was filtered and
15 washed with DMF (3 x 20 mL). A sample of resin was removed and subjected to the Kaiser test. If the test showed the presence of free amine (resin turned blue) the coupling described above was repeated, otherwise the resin was washed with DCM (3 x 20 mL), MeOH (2 x 20 mL), and DCM (2 x 20 mL). (A wash consisted of addition of the solvent and agitation either by nitrogen bubbling or shaking on the orbital shaker
20 for 1-5 minutes, then filtration under vacuum). The resin was dried *in vacuo* at room temperature.

A sample of resin (5-20 mg) was subjected to cleavage with DCM (0.5 mL) and TFA (0.5 mL) for 1 hour at room temperature. The reaction was filtered and the resin washed with DCM (1 x 1 mL). The filtrate and the washing were combined and
25 concentrated *in vacuo* on a Savant SpeedVac Plus. Methanol (1 mL) was added and the mixture concentrated. The product was then characterized by ¹H NMR, (DMSO d-6) δ 4.54 (q, 1H), 1.83 (d, 3H).

Step B: Displacement of bromide with 4-methoxybenzenethiol.

30 The *N*-Hydroxy-2-bromo-propionamide resin prepared in Step A (0.35 g, 1.1 meq/g) was placed in a 20 mL scintillation vial and suspended in THF (2 mL). 4-Methoxybenzenethiol (0.23 mL, 5.0 eq.), sodium iodide (288 mg, 5.0 eq.) and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 0.17 mL, 3.0 eq.) were added. The reaction was shaken at room temperature for 12 - 16 hours. The reaction mixture was poured
35 into a polypropylene syringe barrel fitted with a polypropylene frit, filtered and washed with DMF (2 x 2 mL), DMF:water 9:1 (2 x 2 mL), DMF (2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

WO 99/42436

PCT/US98/17633

- 108 -

Step C: Oxidation of sulfide to sulfoxide.

5 *N*-Hydroxy-2-(4-methoxy-benzenesulfanyl)-propionamide resin prepared in Step B (175 mg, 1.1 meq/g) was suspended in DCM (3.0 mL) and 70% *tert*-butylhydroperoxide (1.0 mL) and benzenesulfonic acid (50 mg) were added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

10

Step D: Oxidation of sulfide to sulfone.

15 *N*-Hydroxy-2-(4-methoxy-benzenesulfanyl)-propionamide resin prepared in Step B (175 mg, 1.1 meq/g) was suspended in DCM (3.0 mL) and mCPBA (180 mg) was added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

20 Step E: Cleavage of *N*-Hydroxy-2-(4-methoxy-benzenesulfonyl)-propionamide from resin.

The *N*-Hydroxy-2-(4-methoxy-benzenesulfonyl)-propionamide resin prepared in Step D (73 mg, 1.2 meq/g) was suspended in DCM (1.0 mL) and TFA (1.0 mL) was added. The reaction was shaken for 1 hour at room temperature. The reaction was filtered and the resin washed with DCM (2 x 1 mL). The filtrate and the washing were combined and concentrated to dryness on a Savant SpeedVac Plus. Methanol (1 mL) was added and the mixture concentrated.

25

84% @ 215 nm; ¹H NMR (DMSO d-6) δ 10.75 (brs, 1 H), 7.95 (brs, 1 H), 7.71 (dd, 2 H), 7.16 (dd, 2 H), 3.87 (s, 3 H), 3.83 (q, 1 H), 1.26 (d, 3 H).

30

The hydroxamic acids of Examples 111-113 are synthesized using appropriate starting materials and following the steps in example 110.

WO 99/42436

PCT/US98/17633

- 109 -

Example 111

N-Hydroxy-2-(4-methoxy-benzenesulfanyl)-propionamide. 72% @ 215 nm

5 *N*-Hydroxy-2-(4-methoxy-benzenesulfinyl)-propionamide. 76% @ 215 nm; ¹H NMR (DMSO d-6) δ 10.90 & 10.60 (brs, 1 H), 7.95 (brs, 1 H) 7.61 & 7.52 (dd, 2 H), 7.15 & 7.10 (dd, 2 H), 3.83 & 3.82 (s, 3 H), 3.42 & 3.28 (q 1H), 1.23 & 0.97 (d, 3 H).

Example 112

10 *N*-Hydroxy-2-(3-methyl-butane-1-sulfanyl)-propionamide. 74% @ 215 nm.

N-Hydroxy-2-(3-methyl-butane-1-sulfinyl)-propionamide. ¹H NMR (DMSO d-6) δ 10.8 (brs 1 H), 7.95 (brs, 1 H), 3.45 & 3.31 (q, 1 H), 2.71 -2.50 (m, 2 H), 1.71-
15 1.46 (m, 3 H), 1.33 & 1.25 (d, 3 H), 0.94-0.82 (m, 6 H)

Example 113

20 *N*-Hydroxy-2-(3-methyl-butane-1-sulfonyl)-propionamide. 84% @ 215 nm.

Example 114

25 *N*-hydroxy-3-methyl-2-(naphthalen-2-ylsulfanyl)-butyramide

Step A: Coupling of 2-bromo-3-methyl-butyric acid to hydroxylamine resin.
4-*O*-Methylhydroxylamine-phenoxyethyl-copoly(styrene-1%-divinylbenzene)-resin¹
(5 g, 1.1 meq/g) was placed in a peptide synthesis vessel and suspended in DMF (40 mL). 2-Bromo-3-methyl-butyric acid (9.96 g, 10.0 eq.) and DIC (9.04 mL, 10.5 eq.)
30 were added. The reaction was shaken on an orbital shaker at room temperature for 2 - 16 hours. The reaction was filtered and washed with DMF (3 x 20 mL). A sample of resin was removed and subjected to the Kaiser test. If the test showed the presence of free amine (resin turned blue) the coupling described above was repeated, otherwise the resin was washed with DCM (3 x 20 mL), MeOH (2 x 20 mL), and DCM (2 x 20 mL).
35 The resin was dried *in vacuo* at room temperature.

Step B: Displacement of bromide with 2-naphthalenethiol.

The 2-bromo hydroxamate resin prepared in Step A (0.15 g, 1.1 meq/g) was placed in a 20 mL scintillation vial and suspended in THF (2 mL). 2-Naphthalenethiol (138 mg,

WO 99/42436

PCT/US98/17633

- 110 -

5.0 eq.), sodium iodide (129 mg, 5.0 eq.) and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 0.078 mL, 3.0 eq.) were added. The reaction was shaken at room temperature for 12 - 16 hours. The reaction mixture was poured into a polypropylene syringe barrel fitted with a polypropylene frit, filtered and washed with DMF (2 x 2 mL), DMF:water 9:1 (2 x 2 mL), DMF (2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step C: Oxidation of sulfide to sulfoxide.

2-(2-Naphthalenesulfanyl)-*N*-hydroxypropionamide resin prepared in Step B (175 mg, 1.1 meq/g) was suspended in DCM (3.0 mL) and 70% *tert*-butylhydroperoxide (1.0 mL) benzenesulfonic acid (50 mg) were added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step D: Oxidation of sulfide to sulfone.

2-(2-Naphthalenesulfanyl)-*N*-hydroxypropionamide resin prepared in Step B (175 mg, 1.1 meq/g) was suspended in DCM (3.0 mL) and mCPBA (180 mg) was added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step E: Cleavage of *N*-Hydroxy-3-methyl-2-(naphthalen-2-ylsulfanyl)-butyramide from resin.

The 2-(2-Naphthalenesulfanyl)-*N*-hydroxypropionamide resin prepared in Step B (73 mg, 1.2 meq/g) was suspended in DCM (1.0 mL) and TFA (1.0 mL) was added. The reaction was shaken for 1 hour at room temperature. The reaction was filtered and the resin washed with DCM (2 x 1 mL). The filtrate and the washing were combined and concentrated to dryness on a Savant SpeedVac Plus. Methanol (1 mL) was added and the mixture concentrated.

83% @ 215 nm; LCMS (API-electrospray) m/z 276 (M+H)⁺; ¹H NMR (DMSO *d*-6) δ 10.7 (brs, 1 H), 7.91 (brs, 1 H), 7.91-7.81 (m, 4 H), 7.55-7.45 (m, 3 H), 3.41 (d, 1 H), 2.09-1.97 (m, 1 H), 1.05 (d, 3 H), 0.97 (d, 3 H).

WO 99/42436

PCT/US98/17633

- 111 -

The hydroxamic acids of Examples 115-118 are synthesized using appropriate starting materials and following the steps in example 114:

Example 115

5 *N*-Hydroxy-3-methyl-2-(naphthalen-2-ylsulfinyl)-butyramide. 67% @ 215 nm.

Example 116

10 *N*-Hydroxy-3-methyl-2-(naphthalen-2-ylsulfonyl)-butyramide. 97% @ 215 nm; LCMS (API-electrospray) m/z 308 (M+H)⁺.

Example 117

15 *N*-Hydroxy-3-methyl-2-phenethylsulfinyl-butylamide. 93% @ 215 nm; LCMS (API-electrospray) m/z 254 (M+H)⁺.

Example 118

20 *N*-Hydroxy-3-methyl-2-phenethylsulfonyl-butylamide. 97% @ 215 nm; LCMS (API-electrospray) m/z 286 (M+H)⁺.

Example 119

25 (1-Hydroxycarbamoyl-propane-1-sulfanyl)-acetic acid methyl ester

Step A: Coupling of 2-bromobutyric acid to hydroxylamine resin.

4-*O*-Methylhydroxylamine-phenoxy-methyl-copoly(styrene-1%-divinylbenzene)-resin¹ (5 g, 1.1 meq/g) was placed in a peptide synthesis vessel and suspended in DMF (40 mL). 2-Bromobutyric acid (3.0 g, 3.0 eq.) HOBt (4.86 g, 6.0 eq.) and DIC (3.75 mL, 4.0 eq.) were added. The reaction was shaken on an orbital shaker at room temperature for 2 - 16 hours. The reaction was filtered and washed with DMF (3 x 20 mL). A sample of resin was removed and subjected to the Kaiser test. If the test showed the presence of free amine (resin turned blue) the coupling described above was repeated, otherwise the resin was washed with DCM (3 x 20 mL), MeOH (2 x 20 mL), and DCM (2 x 20 mL). The resin was dried *in vacuo* at room temperature.

Step B: Displacement of bromide with methyl thioglycolate.

40 The 2-bromo hydroxymate resin prepared in Step A (0.45 g, 1.1 meq/g) was placed in a 20 mL scintillation vial and suspended in THF (2 mL). Methyl thioglycolate (286 mg,

WO 99/42436

PCT/US98/17633

- 112 -

5.0 eq.), sodium iodide (404 mg, 5.0 eq.) and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 0.24 mL, 3.0 eq.) were added. The reaction was shaken at room temperature for 12 - 16 hours. The reaction mixture was poured into a polypropylene syringe barrel fitted with a polypropylene frit, filtered and washed with DMF (2 x 2 mL), DMF:water 9:1 (2 x 2 mL), DMF (2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step C: Oxidation of sulfide to sulfoxide.

(1-Hydroxycarbamoyl-propane-1-sulfanyl)-acetic acid methyl ester resin prepared in Step B (150 mg, 1.1 meq/g) was suspended in DCM (3.0 mL) and 70% *tert*-butylhydroperoxide (1.0 mL) benzenesulfonic acid (50 mg) were added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step D: Oxidation of sulfide to sulfone.

(1-Hydroxycarbamoyl-propane-1-sulfanyl)-acetic acid methyl ester resin prepared in Step B (150 mg, 1.1 meq/g) was suspended in DCM (3.0 mL) and mCPBA (180 mg) was added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step E: Cleavage of (1-Hydroxycarbamoyl-propane-1-sulfanyl)-acetic acid methyl ester from resin

The (1-Hydroxycarbamoyl-propane-1-sulfanyl)-acetic acid methyl ester resin prepared in Step B (150 mg, 1.2 meq/g) was suspended in DCM (1.0 mL) and TFA (1.0 mL) was added. The reaction was shaken for 1 hour at room temperature. The reaction was filtered and the resin washed with DCM (2 x 1 mL). The filtrate and the washing were combined and concentrated to dryness on a Savant SpeedVac Plus. Methanol (1 mL) was added and the mixture concentrated. LCMS (API-electrospray) *m/z* 228 (M+Na)⁺.

The hydroxamic acids of Examples 120-124 are synthesized using appropriate starting materials and following the steps in example 119.

WO 99/42436

PCT/US98/17633

- 113 -

Example 120

(1-Hydroxycarbamoyl-propane-1-sulfonyl)-acetic acid hydroxyamide.
LCMS (API-electrospray) m/z 224 (M+H)⁺.

5

Example 121

(1-Hydroxycarbamoyl-propane-1-sulfinyl)-acetic acid hydroxy amide.
100% @ 220 nm; LCMS (API-electrospray) m/z 240 (M+H)⁺.

10

Example 122

(1-Hydroxycarbamoyl-propane-1-sulfanyl)-propionic acid hydroxyamide.
¹H NMR (DMSO d-6) δ 10.7 (brs, 1 H), 4.03 (t, 2 H), 2.95 (q, 1 H), 2.75-2.70 (m, 1
H), 2.60-2.54 (m, 1 H), 1.74-1.66 (m, 2 H), 1.58-1.50 (m, 4 H), 1.32 (sextet, 2 H),
0.88 (t, 3 H), 0.85 (t, 3 H); LCMS (API-electrospray) m/z 264 (M+H)⁺.

15

Example 123

(1-Hydroxycarbamoyl-propane-1-sulfinyl)-propionic acid hydroxyamide.
83% @ 220 nm; LCMS (API-electrospray) m/z 280 (M+H)⁺.

20

Example 124

(1-Hydroxycarbamoyl-propane-1-sulfonyl)-propionic acid hydroxyamide.
100% @ 220 nm;

25

Example 125

2-(4-Hydroxybenzenesulfanyl)-N-hydroxy-3-phenyl-propionamide

30

Step A: Coupling of 2-bromo-3-phenyl-propionic acid to hydroxylamine resin.
4-O-Methylhydroxylamine-phenoxy methyl-copoly(styrene-1%-divinylbenzene)-resin¹
(5 g, 1.2 meq/g) was placed in a peptide synthesis vessel and suspended in DMF (40
mL). 2-Bromo-3-phenyl-propionic acid (3.5 g, 3.0 eq.) HOBt (4.4 g, 6.0 eq.) and
DIC (3.4 mL, 4.0 eq.) were added. The reaction was shaken on an orbital shaker at
room temperature for 2 - 16 hours. The reaction was filtered and washed with DMF (3
x 20 mL). A sample of resin was removed and subjected to the Kaiser test. If the test
showed the presence of free amine (resin turned blue) the coupling described above

35

WO 99/42436

PCT/US98/17633

- 114 -

was repeated, otherwise the resin was washed with DCM (3 x 20 mL), MeOH (2 x 20 mL), and DCM (2 x 20 mL). The resin was dried *in vacuo* at room temperature.

Step B: Displacement of bromide with 4-hydroxythiophenol.

- 5 The 2-bromo hydroxymate resin prepared in Step A (0.33 g, 1.2 meq/g) was placed in a 20 mL scintillation vial and suspended in THF (2 mL). 4-Hydroxythiophenol (250 mg, 5.0 eq.), sodium iodide (297 mg, 5.0 eq.) and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 0.18 mL, 3.0 eq.) were added. The reaction was shaken at room temperature for 12 - 16 hours. The reaction mixture was poured into a polypropylene
10 syringe barrel fitted with a polypropylene frit, filtered and washed with DMF (2 x 2 mL), DMF:water 9:1 (2 x 2 mL), DMF (2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step C: Oxidation of sulfide to sulfoxide.

- 15 2-(4-Hydroxybenzenesulfanyl)-*N*-hydroxy-3-phenyl-propionamide resin prepared in Step B (110 mg, 1.1 meq/g) was suspended in DCM (3.0 mL) and 70% *tert*-butylhydroperoxide (0.73 mL) benzenesulfonic acid (36 mg) were added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x
20 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step D: Oxidation of sulfide to sulfone.

- 2-(4-Hydroxybenzenesulfanyl)-*N*-hydroxy-3-phenyl-propionamide resin prepared in Step B (110 mg, 1.1 meq/g) was suspended in DCM (3.0 mL) and mCPBA (132 mg)
25 was added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

- 30 Step E: Cleavage of 2-(4-Hydroxybenzenesulfanyl)-*N*-hydroxy-3-phenyl-propionamide from resin.

- The 2-(4-Hydroxybenzenesulfanyl)-*N*-hydroxy-3-phenyl-propionamide resin prepared in Step B (110 mg, 1.2 meq/g) was suspended in DCM (1.0 mL) and TFA (1.0 mL) was added. The reaction was shaken for 1 hour at room temperature. The reaction was
35 filtered and the resin washed with DCM (2 x 1 mL). The filtrate and the washing were combined and concentrated to dryness on a Savant SpeedVac Plus. Methanol (1 mL)

WO 99/42436

PCT/US98/17633

- 115 -

was added and the mixture concentrated. 84% @ 215 nm; ¹H NMR (DMSO d-6) δ 10.41 (brs, 1 H), 7.95 (brs (1 H), 7.30-7.15 (m, 5 H), 7.10 (dd, 2 H), 6.75 (dd, 2 H), 3.53 (q, 1 H), 3.05 (dd, 1 H), 2.79 (dd, 1 H).

- 5 The hydroxamic acids of Examples 126-130 are synthesized using appropriate starting materials and following the steps in example 125.

Example 126

- 10 2-(4-Hydroxybenzenesulfinyl)-*N*-hydroxy-3-phenyl-propionamide. 73% @ 215 nm;

Example 127

- 2-(4-Hydroxybenzenesulfonyl)-*N*-hydroxy-3-phenyl-propionamide. 77% @ 215 nm;
15 ¹H NMR (DMSO d-6) δ 10.50 (brs, 1 H), 7.95 (brs, 1 H), 7.68-7.57 (m, 2 H), 7.28-7.17 (m, 3 H), 7.08-7.98 (m, 2 H), 6.95-6.87 (m, 2 H), 3.96 (t, 1 H), 3.02 (d, 2 H).

Example 128

- 20 2-(4-Acetylamino-benzenesulfanyl)-*N*-hydroxy-3-phenyl-propionamide. 86% @ 215 nm; ¹H NMR (DMSO d-6) δ 10.50 (brs, 1 H), 10.03 (brs, 1 H), 8.13 (brs, 1 H), 7.56-7.12 (m, 9 H), 3.67 (q, 1 H), 3.08 (dd, 1 H), 2.84 (dd, 1 H), 2.04 (s, 3 H)

Example 129

- 25 2-(4-Acetylamino-benzenesulfinyl)-*N*-hydroxy-3-phenyl-propionamide.
73% @ 215 nm.

Example 130

- 30 2-(4-Acetylamino-benzenesulfonyl)-*N*-hydroxy-3-phenyl-propionamide.
95% @ 215 nm;

Example 131

- 35 4-Hydroxycarbamoyl-4-(4-methanesulfanyl-phenylsulfanyl)-butyric acid methyl ester

Step A: Coupling of 2-bromo-5-methyl glutaric acid to hydroxylamine resin.

4-*O*-Methylhydroxylamine-phenoxy methyl-copoly(styrene-1%-divinylbenzene)-resin¹

- 40 (4.5 g, 1.2 meq/g) was placed in a peptide synthesis vessel and suspended in DMF (40

WO 99/42436

PCT/US98/17633

- 116 -

- mL). S-2-Bromo-5-methyl glutarate (3.87 g, 3.0 eq.) HOBt (4.4 g, 6.0 eq.) and DIC (3.4 mL, 4.0 eq.) were added. The reaction was shaken on an orbital shaker at room temperature for 2 - 16 hours. The reaction was filtered and washed with DMF (3 x 20 mL). A sample of resin was removed and subjected to the Kaiser test. If the test showed the presence of free amine (resin turned blue) the coupling described above was repeated, otherwise the resin was washed with DCM (3 x 20 mL), MeOH (2 x 20 mL), and DCM (2 x 20 mL). The resin was dried *in vacuo* at room temperature.

Step B: Displacement of bromide with 4-hydroxythiophenol.

- The 2-bromo hydroxymate resin prepared in Step A (0.22 g, 1.2 meq/g) was placed in a 20 mL scintillation vial and suspended in THF (2 mL). 4-(Methylthio)thiophenol (206 mg, 5.0 eq.), sodium iodide (197 mg, 5.0 eq.) and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 0.12 mL, 3.0 eq.) were added. The reaction was shaken at room temperature for 12 - 16 hours. The reaction mixture was poured into a polypropylene syringe barrel fitted with a polypropylene frit, filtered and washed with DMF (2 x 2 mL), DMF:water 9:1 (2 x 2 mL), DMF (2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step C: Oxidation of sulfide to sulfoxide.

- 4-Hydroxycarbamoyl-4-(4-methanesulfanyl-phenylsulfanyl)-butyric acid methyl ester resin prepared in Step B (73 mg, 1.1 meq/g) was suspended in DCM (1.5 mL) and 70% *tert*-butylhydroperoxide (0.49 mL) benzenesulfonic acid (24 mg) were added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step D: Oxidation of sulfide to sulfone.

- 4-Hydroxycarbamoyl-4-(4-methanesulfanyl-phenylsulfanyl)-butyric acid methyl ester resin prepared in Step B (73 mg, 1.1 meq/g) was suspended in DCM (1.5 mL) and mCPBA (87 mg) was added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step E: Cleavage of 4-Hydroxycarbamoyl-4-(4-methanesulfanyl-phenylsulfanyl)-butyric acid methyl ester from resin.

WO 99/42436

PCT/US98/17633

- 117 -

The 4-Hydroxycarbamoyl-4-(4-methanesulfanyl-phenylsulfanyl)-butyric acid methyl ester resin prepared in Step B (73 mg, 1.2 meq/g) was suspended in DCM (1.0 mL) and TFA (1.0 mL) was added. The reaction was shaken for 1 hour at room temperature. The reaction was filtered and the resin washed with DCM (2 x 1 mL).

- 5 The filtrate and the washing were combined and concentrated to dryness on a Savant SpeedVac Plus. Methanol (1 mL) was added and the mixture concentrated. 77% @ 215 nm; LCMS (API-electrospray) m/z 316 (M+H)⁺.

- 10 The hydroxamic acids of Examples 132-139 are synthesized using appropriate starting materials and following the steps in example 131.

Example 132

- 15 4-Hydroxycarbamoyl-4-(4-methanesulfinyl-phenylsulfinyl)-butyric acid hydroxyamide. 79% @ 215 nm; LCMS (API-electrospray) m/z 348 (M+H)⁺.

Example 133

- 20 4-Hydroxycarbamoyl-4-(4-methanesulfonyl-phenylsulfonyl)-butyric acid hydroxyamide. 78% @ 215 nm; LCMS (API-electrospray) m/z 380 (M+H)⁺.

Example 134

- 25 4-Hydroxycarbamoyl-4-(4-bromo-benzenesulfanyl)-butyric acid hydroxyamide. 93% @ 215 nm.

Example 135

- 30 4-Hydroxycarbamoyl-4-(4-bromo-benzenesulfinyl)-butyric acid hydroxyamide. 80% @ 215 nm.

Example 136

- 35 4-Hydroxycarbamoyl-4-(4-bromo-benzenesulfonyl)-butyric acid hydroxyamide. 77% @ 215 nm.

Example 137

- 40 4-Hydroxycarbamoyl-4-(2-trifluoromethyl-benzenesulfanyl)-butyric acid hydroxyamide. 93% @ 215 nm.

WO 99/42436

PCT/US98/17633

- 118 -

Example 138

4-Hydroxycarbamoyl-4-(2-trifluoromethyl-benzenesulfinyl)-butyric acid hydroxyamide. 72% @ 215 nm.

5

Example 139

4-Hydroxycarbamoyl-4-(2-trifluoromethyl-benzenesulfonyl)-butyric acid hydroxyamide. 90% @ 215 nm.

10

Example 140

2-(3-methoxy-benzenesulfanyl)decanoic acid hydroxamide

15 Step A: Coupling of 2-bromo-decanoic acid to hydroxylamine resin.

4-O-Methylhydroxylamine-phenoxyethyl-copoly(styrene-1%-divinylbenzene)-resin¹ (4.5 g, 1.2 meq/g) was placed in a peptide synthesis vessel and suspended in DMF (40 mL). 2-Bromo-decanoic acid (4.07 g, 3.0 eq.) HOBt (4.4 g, 6.0 eq.) and DIC (3.4 mL, 4.0 eq.) were added. The reaction was shaken on an orbital shaker at room temperature for 2 - 16 hours. The reaction was filtered and washed with DMF (3 x 20 mL). A sample of resin was removed and subjected to the Kaiser test. If the test showed the presence of free amine (resin turned blue) the coupling described above was repeated, otherwise the resin was washed with DCM (3 x 20 mL), MeOH (2 x 20 mL), and DCM (2 x 20 mL). The resin was dried *in vacuo* at room temperature.

25

Step B: Displacement of bromide with 3-methoxy-benzenethiol.

The 2-bromo hydroxamate resin prepared in Step A (0.22 g, 1.2 meq/g) was placed in a 20 mL scintillation vial and suspended in THF (2 mL). 3-Methoxy-benzenethiol (185 mg, 5.0 eq.), sodium iodide (197 mg, 5.0 eq.) and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 0.12 mL, 3.0 eq.) were added. The reaction was shaken at room temperature for 12 - 16 hours. The reaction mixture was poured into a polypropylene syringe barrel fitted with a polypropylene frit, filtered and washed with DMF (2 x 2 mL), DMF:water 9:1 (2 x 2 mL), DMF (2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

35

Step C: Oxidation of sulfide to sulfoxide.

2-(3-Methoxy-benzenesulfanyl)decanoic acid hydroxamide resin prepared in Step B (73 mg, 1.1 meq/g) was suspended in DCM (1.5 mL) and 70% *tert*-butylhydroperoxide (0.49 mL) benzenesulfonic acid (24 mg) were added. The reaction mixture was shaken

WO 99/42436

PCT/US98/17633

- 119 -

on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

5 Step D: Oxidation of sulfide to sulfone.

2-(3-Methoxy-benzenesulfanyl)decanoic acid hydroxamide resin prepared in Step B (73 mg, 1.1 meq/g) was suspended in DCM (1.5 mL) and mCPBA (87 mg) was added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL),
10 MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step E: Cleavage of 2-(3-methoxy-benzenesulfanyl)decanoic acid hydroxamide from resin.

15 The 2-(3-methoxy-benzenesulfanyl)decanoic acid hydroxamide resin prepared in Step B (73 mg, 1.2 meq/g) was suspended in DCM (1.0 mL) and TFA (1.0 mL) was added. The reaction was shaken for 1 hour at room temperature. The reaction was filtered and the resin washed with DCM (2 x 1 mL). The filtrate and the washing were combined and concentrated to dryness on a Savant SpeedVac Plus. Methanol (1 mL) was added
20 and the mixture concentrated. 89% @ 215 nm.

The hydroxamic acids of Examples 141-145 are synthesized using appropriate starting materials and following the steps in example 140.

25 Example 141

2-(3-Methoxy-benzenesulfanyl)decanoic acid hydroxamide. 96% @ 215 nm.

Example 142

30 2-(3-Methoxy-benzenesulfonyl)decanoic acid hydroxamide. 96% @ 215 nm.

Example 143

35 2-(4-methanesulfanyl-benzenesulfanyl)decanoic acid hydroxamide. 85% @ 215 nm; LCMS (API-electrospray) m/z 342 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 120 -

Example 144

2-(4-methanesulfinyl-benzenesulfinyl)decanoic acid hydroxamide. 86% @ 215 nm;
LCMS (API-electrospray) m/z 374 (M+H)⁺.

5

Example 145

2-(4-methanesulfonyl-benzenesulfonyl)decanoic acid hydroxamide. 92% @ 215 nm.

10

Example 146

3-benzyloxy-*N*-hydroxy-2-(4-methanesulfanyl-benzenesulfanyl)-propionamide

Step A: Coupling of 2-bromo-3-benzyloxy propionic acid to hydroxylamine resin.

- 15 4-*O*-Methylhydroxylamine-phenoxymethyl-copoly(styrene-1%-divinylbenzene)-resin¹
(4.5 g, 1.2 meq/g) was placed in a peptide synthesis vessel and suspended in DMF (40 mL). S-2-Bromo-3-benzyloxy-propionic acid (4.2 g, 3.0 eq.) HOBt (4.4 g, 6.0 eq.) and DIC (3.4 mL, 4.0 eq.) were added. The reaction was shaken on an orbital shaker at room temperature for 2 - 16 hours. The reaction was filtered and washed with DMF
20 (3 x 20 mL). A sample of resin was removed and subjected to the Kaiser test. If the test showed the presence of free amine (resin turned blue) the coupling described above was repeated, otherwise the resin was washed with DCM (3 x 20 mL), MeOH (2 x 20 mL), and DCM (2 x 20 mL). The resin was dried *in vacuo* at room temperature.

- 25 Step B: Displacement of bromide with 4-(methylthio)thiophenol.

The 2-bromo hydroxymate resin prepared in Step A (0.22 g, 1.2 meq/g) was placed in a 20 mL scintillation vial and suspended in THF (2 mL). 4-(Methylthio)thiophenol (206 mg, 5.0 eq.), sodium iodide (197 mg, 5.0 eq.) and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 0.12 mL, 3.0 eq.) were added. The reaction was shaken at room
30 temperature for 12 - 16 hours. The reaction mixture was poured into a polypropylene syringe barrel fitted with a polypropylene frit, filtered and washed with DMF (2 x 2 mL), DMF:water 9:1 (2 x 2 mL), DMF (2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

- 35 Step C: Oxidation of sulfide to sulfoxide.

3-Benzyloxy-*N*-hydroxy-2-(4-methanesulfanyl-benzenesulfanyl)-propionamide resin prepared in Step B (73 mg, 1.1 meq/g) was suspended in DCM (1.5 mL) and 70% *tert*-butylhydroperoxide (0.49 mL) benzenesulfonic acid (24 mg) were added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The

WO 99/42436

PCT/US98/17633

- 121 -

reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step D: Oxidation of sulfide to sulfone.

- 5 3-Benzyloxy-*N*-hydroxy-2-(4-methanesulfanyl-benzenesulfanyl)-propionamide resin prepared in Step B (73 mg, 1.1 meq/g) was suspended in DCM (1.5 mL) and mCPBA (87 mg) was added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried
10 *in vacuo* at room temperature.

Step E: Cleavage of 3-benzyloxy-*N*-hydroxy-2-(4-methanesulfanyl-benzenesulfanyl)-propionamide from resin.

- The 3-benzyloxy-*N*-hydroxy-2-(4-methanesulfanyl-benzenesulfanyl)-propionamide
15 resin prepared in Step B (73 mg, 1.2 meq/g) was suspended in DCM (1.0 mL) and TFA (1.0 mL) was added. The reaction was shaken for 1 hour at room temperature. The reaction was filtered and the resin washed with DCM (2 x 1 mL). The filtrate and the washing were combined and concentrated to dryness on a Savant SpeedVac Plus. Methanol (1 mL) was added and the mixture concentrated. 76% @ 215 nm; LCMS
20 (API-electrospray) m/z 350 (M+H)⁺.

The hydroxamic acids of Examples 147-151 are synthesized using appropriate starting materials and following the steps in example 146.

25 Example 147

3-Benzyloxy-*N*-hydroxy-2-(4-methanesulfinyl-benzenesulfinyl)-propionamide.
70% @ 215 nm; LCMS (API-electrospray) m/z 382 (M+H)⁺.

30 Example 148

3-Benzyloxy-*N*-hydroxy-2-(4-methanesulfonyl-benzenesulfonyl)-propionamide.
63% @ 215 nm; LCMS (API-electrospray) m/z 414 (M+H)⁺.

35 Example 149

3-Benzyloxy-*N*-hydroxy-2-(2-chloro-benzylsulfanyl)-propionamide. 90% @ 215 nm.

WO 99/42436

PCT/US98/17633

- 122 -

Example 150

3-Benzyloxy-N-hydroxy-2-(2-chloro-benzylsulfinyl)-propionamide. 70% @ 215 nm.

5

Example 151

3-Benzyloxy-N-hydroxy-2-(2-chloro-benzylsulfonyl)-propionamide. 72% @ 215 nm.

10

Example 152

2-(2-bromo-benzenesulfanyl)-N-hydroxy-3-(3H-imidazol-4-yl)-propionamide

Step A: Coupling of 2-bromo-3-(3H-imidazol-4-yl)-propionic acid to hydroxylamine resin.

15

4-O-Methylhydroxylamine-phoxymethyl-copoly(styrene-1%-divinylbenzene)-resin¹ (4.5 g, 1.2 meq/g) was placed in a peptide synthesis vessel and suspended in DMF (40 mL). S-2-Bromo-3-(3H-imidazol-4-yl)-propionic acid (3.55 g, 3.0 eq.) HOBt (4.4 g, 6.0 eq.) and DIC (3.4 mL, 4.0 eq.) were added. The reaction was shaken on an orbital shaker at room temperature for 2 - 16 hours. The reaction was filtered and washed with DMF (3 x 20 mL). A sample of resin was removed and subjected to the Kaiser test. If the test showed the presence of free amine (resin turned blue) the coupling described above was repeated, otherwise the resin was washed with DCM (3 x 20 mL), MeOH (2 x 20 mL), and DCM (2 x 20 mL). The resin was dried *in vacuo* at room temperature.

25

Step B: Displacement of bromide with 2-bromothiophenol.

The 2-bromo hydroxymate resin prepared in Step A (0.22 g, 1.2 meq/g) was placed in a 20 mL scintillation vial and suspended in THF (2 mL). 2-Bromothiophenol (249 mg, 5.0 eq.), sodium iodide (197 mg, 5.0 eq.) and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 0.12 mL, 3.0 eq.) were added. The reaction was shaken at room temperature for 12 - 16 hours. The reaction mixture was poured into a polypropylene syringe barrel fitted with a polypropylene frit, filtered and washed with DMF (2 x 2 mL), DMF:water 9:1 (2 x 2 mL), DMF (2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

35

Step C: Oxidation of sulfide to sulfoxide.

2-(2-Bromo-benzenesulfanyl)-N-hydroxy-3-(3H-imidazol-4-yl)-propionamide resin prepared in Step B (73 mg, 1.1 meq/g) was suspended in DCM (1.5 mL) and 70% *tert*-butylhydroperoxide (0.49 mL) benzenesulfonic acid (24 mg) were added. The reaction

WO 99/42436

PCT/US98/17633

- 123 -

mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

5 Step D: Oxidation of sulfide to sulfone.

2-(2-Bromo-benzenesulfanyl)-N-hydroxy-3-(3H-imidazol-4-yl)-propionamide resin prepared in Step B (73 mg, 1.1 meq/g) was suspended in DCM (1.5 mL) and mCPBA (87 mg) was added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step E: Cleavage of 2-(2-bromo-benzenesulfanyl)-N-hydroxy-3-(3H-imidazol-4-yl)-propionamide from resin.

15 The 2-(2-bromo-benzenesulfanyl)-N-hydroxy-3-(3H-imidazol-4-yl)-propionamide resin prepared in Step B (73 mg, 1.2 meq/g) was suspended in DCM (1.0 mL) and TFA (1.0 mL) was added. The reaction was shaken for 1 hour at room temperature. The reaction was filtered and the resin washed with DCM (2 x 1 mL). The filtrate and the washing were combined and concentrated to dryness on a Savant SpeedVac Plus.

20 Methanol (1 mL) was added and the mixture concentrated. 86% @ 215 nm.

The hydroxamic acids of Examples 153-154 are synthesized using appropriate starting materials and following the steps in example 152.

25 Example 153

2-(4-bromo-benzenesulfinyl)-N-hydroxy-3-(3H-imidazol-4-yl)-propionamide.
69% @ 215 nm

30 Example 154

2-(4-chloro-benzenesulfonyl)-N-hydroxy-3-(3H-imidazol-4-yl)-propionamide.

35 Example 155

2-(3-fluorophenylsulfanyl)-5-guanidino-pentanoic acid hydroxyamide

Step A: Coupling of 2-bromo-5-guanidino-pentanoic acid to hydroxylamine resin.

4-O-Methylhydroxylamine-phoxymethyl-copoly(styrene-1%-divinylbenzene)-resin¹
40 (4.5 g, 1.2 meq/g) was placed in a peptide synthesis vessel and suspended in DMF (40

WO 99/42436

PCT/US98/17633

- 124 -

- mL). S-2-Bromo-5-guanidino-pentanoic acid (3.85 g, 3.0 eq.) HOBt (4.4 g, 6.0 eq.) and DIC (3.4 mL, 4.0 eq.) were added. The reaction was shaken on an orbital shaker at room temperature for 2 - 16 hours. The reaction was filtered and washed with DMF (3 x 20 mL). A sample of resin was removed and subjected to the Kaiser test. If the test showed the presence of free amine (resin turned blue) the coupling described above was repeated, otherwise the resin was washed with DCM (3 x 20 mL), MeOH (2 x 20 mL), and DCM (2 x 20 mL). The resin was dried *in vacuo* at room temperature.

Step B: Displacement of bromide with 3-fluorothiophenol.

- The 2-bromo hydroxymate resin prepared in Step A (0.22 g, 1.2 meq/g) was placed in a 20 mL scintillation vial and suspended in THF (2 mL). 3-Fluorothiophenol (169 mg, 5.0 eq.), sodium iodide (197 mg, 5.0 eq.) and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 0.12 mL, 3.0 eq.) were added. The reaction was shaken at room temperature for 12 - 16 hours. The reaction mixture was poured into a polypropylene syringe barrel fitted with a polypropylene frit, filtered and washed with DMF (2 x 2 mL), DMF:water 9:1 (2 x 2 mL), DMF (2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step C: Oxidation of sulfide to sulfoxide.

- 2-(3-Fluorophenylsulfanyl)-5-guanidino-pentanoic acid hydroxyamide resin prepared in Step B (73 mg, 1.1 meq/g) was suspended in DCM (1.5 mL) and 70% *tert*-butylhydroperoxide (0.49 mL) benzenesulfonic acid (24 mg) were added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step D: Oxidation of sulfide to sulfone.

- 2-(3-Fluorophenylsulfanyl)-5-guanidino-pentanoic acid hydroxyamide resin prepared in Step B (73 mg, 1.1 meq/g) was suspended in DCM (1.5 mL) and mCPBA (87 mg) was added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

- Step E: Cleavage of 2-(3-fluorophenylsulfanyl)-5-guanidino-pentanoic acid hydroxyamide from resin.

WO 99/42436

PCT/US98/17633

- 125 -

The 2-(3-fluorophenylsulfanyl)-5-guanidino-pentanoic acid hydroxyamide resin prepared in Step B (73 mg, 1.2 meq/g) was suspended in DCM (1.0 mL) and TFA (1.0 mL) was added. The reaction was shaken for 1 hour at room temperature. The reaction was filtered and the resin washed with DCM (2 x 1 mL). The filtrate and the washing were combined and concentrated to dryness on a Savant SpeedVac Plus. Methanol (1 mL) was added and the mixture concentrated. 93% @ 215 nm.

The hydroxamic acids of Examples 156-159 are synthesized using appropriate starting materials and following the steps in example 155:

10

Example 156

2-(3-Fluorophenylsulfinyl)-5-guanidino-pentanoic acid hydroxyamide. 80% @ 220 nm; LCMS (API-electrospray) m/z 317 (M+H)⁺.

15

Example 157

2-(2-Bromosulfinyl)-5-guanidino-pentanoic acid hydroxyamide. 92% @ 220 nm; ¹H NMR (DMSO d-6) δ 10.90 (brs, 2 H), 10.41 (brs, 1H), 7.95 (brs, 1 H), 7.66-7.14 (m, 5 H), 3.72 (q, 1 H), 3.13 (q, 2 H), 1.90-1.66 (m, 2 H), 1.58-1.43 (2 H).

20

Example 158

2-(2-Bromosulfinyl)-5-guanidino-pentanoic acid hydroxyamide. 79% @ 220 nm; LCMS (API-electrospray) m/z 379 (M+H)⁺.

25

Example 159

2-(2-Bromosulfonyl)-5-guanidino-pentanoic acid hydroxyamide. ¹H NMR (DMSO d-6) δ 8.03-7.45 (m, 5 H), 4.52 (q, 1 H), 3.16 (q, 2 H), 2.07-1.90 (m, 2 H), 1.66-1.59 (2 H).

30

Example 160

2-(2,5-dichlorobenzenesulfanyl)-octanoic acid hydroxyamide

35

Step A: Coupling of 2-bromo-octanoic acid to hydroxylamine resin.

4-O-Methylhydroxylamine-phenoxy-methyl-copoly(styrene-1%-divinylbenzene)-resin¹ (10.0 g, 1.2 meq/g) was placed in a peptide synthesis vessel and suspended in DMF

WO 99/42436

PCT/US98/17633

- 126 -

- (80 mL). 2-Bromo-octanoic acid (8.4 g, 3.0 eq.) HOBt (8.8 g, 6.0 eq.) and DIC (7.2 mL, 4.0 eq.) were added. The reaction was shaken on an orbital shaker at room temperature for 2 - 16 hours. The reaction was filtered and washed with DMF (3 x 20 mL). A sample of resin was removed and subjected to the Kaiser test. If the test
- 5 showed the presence of free amine (resin turned blue) the coupling described above was repeated, otherwise the resin was washed with DCM (3 x 20 mL), MeOH (2 x 20 mL), and DCM (2 x 20 mL). The resin was dried *in vacuo* at room temperature.

Step B: Displacement of bromide with 2,5-dichlorothiophenol.

- 10 The 2-bromo hydroxymate resin prepared in Step A (0.45 g, 1.2 meq/g) was placed in a 20 mL scintillation vial and suspended in THF (6 mL). 2,5-Dichlorothiophenol (483 mg, 5.0 eq.), sodium iodide (404 mg, 5.0 eq.) and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 0.24 mL, 3.0 eq.) were added. The reaction was shaken at room temperature for 12 - 16 hours. The reaction mixture was poured into a polypropylene
- 15 syringe barrel fitted with a polypropylene frit, filtered and washed with DMF (2 x 2 mL), DMF:water 9:1 (2 x 2 mL), DMF (2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step C: Oxidation of sulfide to sulfoxide.

- 20 2-(2,5-Dichlorobenzenesulfanyl)-octanoic acid hydroxyamide resin prepared in Step B (150 mg, 1.1 meq/g) was suspended in DCM (3.0 mL) and 70% *tert*-butylhydroperoxide (1.0 mL) benzenesulfonic acid (50 mg) were added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x
- 25 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step D: Oxidation of sulfide to sulfone.

- 2-(2,5-Dichlorobenzenesulfanyl)-octanoic acid hydroxyamide resin prepared in Step B (150 mg, 1.1 meq/g) was suspended in DCM (3.0 mL) and mCPBA (180 mg) was
- 30 added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

- 35 Step E: Cleavage of 2-(2,5-dichlorobenzenesulfanyl)-octanoic acid hydroxyamide from resin.

WO 99/42436

PCT/US98/17633

- 127 -

- The 2-(2,5-dichlorobenzenesulfanyl)-octanoic acid hydroxyamide resin prepared in Step B (73 mg, 1.2 meq/g) was suspended in DCM (1.0 mL) and TFA (1.0 mL) was added. The reaction was shaken for 1 hour at room temperature. The reaction was filtered and the resin washed with DCM (2 x 1 mL). The filtrate and the washing were
5 combined and concentrated to dryness on a Savant SpeedVac Plus. Methanol (1 mL) was added and the mixture concentrated. 92% @ 215 nm; ¹H NMR (DMSO d-6) δ 10.96 (brs, 1 H), 9.26 (brs, 1 H), 7.93-7.76 (m, 3 H), 4.07 (q, 1 H), 2.04-1.85 (m, 1 H), 1.78-1.64 (m, 1 H), 1.32-1.09 (m, 8 H), 0.81 (t, 3 H).
- 10 The hydroxamic acids of Examples 161-167 are synthesized using appropriate starting materials and following the steps in example 160.

Example 161

- 15 2-(2,5-Dichlorobenzenesulfonyl)-octanoic acid hydroxyamide. 96% @ 215 nm.

Example 162

- 2-(3-Methoxybenzenesulfanyl)-octanoic acid hydroxyamide 86% @ 220 nm; LCMS
20 (API-electrospray) m/z 298 (M+H)⁺.

Example 163

- 2-(3-Methoxybenzenesulfinyl)-octanoic acid hydroxyamide 96% @ 220 nm.
25

Example 164

- 2-(3-Methoxybenzenesulfonyl)-octanoic acid hydroxyamide 83% @ 220 nm.

Example 165

- 30 2-(3,4-Dimethoxybenzenesulfanyl)-octanoic acid hydroxyamide 87% @ 215 nm; LCMS (API-electrospray) m/z 328 (M+H)⁺.

Example 166

- 35 2-(3,4-Dimethoxybenzenesulfinyl)-octanoic acid hydroxyamide 90% @ 215 nm.

Example 167

- 40 2-(3,4-Dimethoxybenzenesulfonyl)-octanoic acid hydroxyamide 87% @ 215 nm.

WO 99/42436

PCT/US98/17633

- 128 -

The hydroxamic acid compounds of Examples 168-198 are synthesized using appropriate starting materials and following the steps in example 160. The crude products are dissolved in DMSO:methanol (1:1, 2 mL) and purified by reverse phase HPLC under the conditions described below:

Column: ODS-A, 20 mm x 50 mm, 5 μ m particle size (YMC, Inc. Wilmington, North Carolina)

Solvent Gradient	Time	Water	Acetonitrile
10	0.0	95	5
	25 min.	5	95

Flow Rate: 15 mL/min.

Example 168

15 2-(2-Benzimidazol-2-ylsulfanyl)-octanoic acid hydroxyamide 81% @ 215 nm; LCMS (API-electrospray) m/z 308 (M+H)⁺.

Example 169

20 2-(2-Benzooxazol-2-ylsulfanyl)-octanoic acid hydroxyamide 72% @ 215 nm; LCMS (API-electrospray) m/z 309 (M+H)⁺.

Example 170

25 2-(2-Benzothiazol-2-ylsulfanyl)-octanoic acid hydroxyamide 72% @ 215 nm; LCMS (API-electrospray) m/z 325 (M+H)⁺.

Example 171

30 2-(2-Pyridine-2-sulfanyl)-octanoic acid hydroxyamide 76% @ 215 nm; LCMS (API-electrospray) m/z 269 (M+H)⁺.

Example 172

35 2-(4-Phenyl-thiazole-2-sulfanyl)-octanoic acid hydroxyamide 97% @ 215 nm; LCMS (API-electrospray) m/z 336 (M+H)⁺.

Example 173

40

WO 99/42436

PCT/US98/17633

- 129 -

2-(2-Pyridin-2-yl-ethylsulfanyl)-octanoic acid hydroxyamide 84% @ 215 nm; LCMS (API-electrospray) m/z 297 (M+H)⁺.

Example 174

5

2-(2-Phenyl-5H-tetrazol-5-ylsulfanyl)-octanoic acid hydroxyamide 67% @ 215 nm; LCMS (API-electrospray) m/z 338 (M+H)⁺.

Example 175

10

2-(2-Pyrazin-2-yl-ethylsulfanyl)-octanoic acid hydroxyamide 98% @ 215 nm; LCMS (API-electrospray) m/z 298 (M+H)⁺.

Example 176

15

2-(1-Methyl-1H-tetrazol-5-ylsulfanyl)-octanoic acid hydroxyamide 66% @ 215 nm; LCMS (API-electrospray) m/z 274 (M+H)⁺.

Example 177

20

2-(2-Benzimidazol-2-ylsulfinyl)-octanoic acid hydroxyamide 81% @ 215 nm.

Example 178

25

2-(2-Pyridine-2-sulfinyl)-octanoic acid hydroxyamide 76% @ 215 nm;.

Example 179

30

2-(4-Phenyl-thiazole-2-sulfinyl)-octanoic acid hydroxyamide 78% @ 215 nm.

Example 180

35

2-(2-Pyrazin-2-yl-ethylsulfinyl)-octanoic acid hydroxyamide 96% @ 215 nm; LCMS (API-electrospray) m/z 314 (M+H)⁺.

Example 181

40

2-(3-Oxy-1H-benzimidazole-2-sulfonyl)-octanoic acid hydroxyamide 63% @ 215 nm; LCMS (API-electrospray) m/z 356 (M+H)⁺.

Example 182

2-(4-Phenyl-thiazole-2-sulfonyl)-octanoic acid hydroxyamide 70% @ 215 nm; LCMS (API-electrospray) m/z 383 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 130 -

Example 183

2-[2-(1-Oxy-pyridin-2-yl)-ethanesulfonyl]-octanoic acid hydroxyamide 77% @ 215
5 nm; LCMS (API-electrospray) m/z 345 (M+H)⁺.

Example 184

3-(1-Hydroxycarbamoyl-heptylsulfanyl)-benzoic acid hydroxyamide. 100% @ 220
10 nm; LCMS (API-electrospray) m/z 312 (M+H)⁺.

Example 185

3-[4-(1-Hydroxycarbamoyl-heptylsulfanyl)-phenyl]-propionic acid hydroxyamide.
15 90% @ 220 nm; LCMS (API-electrospray) m/z 340 (M+H)⁺.

Example 186

2-(Thiazol-2-ylsulfanyl)-octanoic acid hydroxyamide. 75% @ 215 nm; LCMS (API-
20 electrospray) m/z 275 (M+H)⁺.

Example 187

2-(2,5-Dioxo-imidazolidin-4-ylmethylsulfanyl)-octanoic acid hydroxyamide. 98% @
25 215 nm; LCMS (API-electrospray) m/z 304 (M+H)⁺.

Example 188

3-(1-Hydroxycarbamoyl-heptylsulfinyl)-benzoic acid hydroxyamide. 84% @ 220 nm;
30 LCMS (API-electrospray) m/z 328 (M+H)⁺.

Example 189

3-[4-(1-Hydroxycarbamoyl-heptylsulfinyl)-phenyl]-propionic acid hydroxyamide. 78%
35 @ 220 nm; LCMS (API-electrospray) m/z 356 (M+H)⁺.

Example 190

2-(Quinoline-8-sulfinyl)-octanoic acid hydroxyamide. 87% @ 220 nm; LCMS (API-
40 electrospray) m/z 335 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 131 -

Example 191

2-(Naphthalen-2-ylcarbamoylmethanesulfinyl)-octanoic acid hydroxyamide. 83% @ 220 nm; LCMS (API-electrospray) m/z 391 (M+H)⁺.

5

Example 192

3-(1-Hydroxycarbamoyl-heptylsulfonyl)-benzoic acid hydroxyamide. 72% @ 215 nm.

10

Example 193

3-[4-(1-Hydroxycarbamoyl-heptylsulfonyl)-phenyl]-propionic acid hydroxyamide. 67% @ 215 nm.

15

Example 194

2-(1H-Imidazole-2-sulfonyl)-octanoic acid hydroxyamide. 95% @ 215 nm; LCMS (API-electrospray) m/z 290 (M+H)⁺.

20

Example 195

2-(Thiazol-2-ylsulfonyl)-octanoic acid hydroxyamide. 91% @ 215 nm; LCMS (API-electrospray) m/z 307 (M+H)⁺.

25

Example 196

2-(Quinoline-8-sulfonyl)-octanoic acid hydroxyamide. 94% @ 220 nm; LCMS (API-electrospray) m/z 351 (M+H)⁺.

30

Example 197

2-(Naphthalen-2-ylcarbamoylmethanesulfonyl)-octanoic acid hydroxyamide. 79% @ 220 nm; LCMS (API-electrospray) m/z 407 (M+H)⁺.

35

Example 198

2-(2,5-Dioxo-imidazolidin-4-ylmethylsulfonyl)-octanoic acid hydroxyamide. 97% @ 215 nm.

WO 99/42436

PCT/US98/17633

- 132 -

Example 199

Step A: Displacement of bromide with 4-fluorothiophenol.

5 The 2-bromo hydroxymate resin prepared in Example 160, Step A (9.4 g, 1.2 meq/g) was placed in a peptide synthesis vessel and suspended in THF (50 mL). 4-Fluorothiophenol (6.6 g, 5.0 eq.), sodium iodide (7.7 g, 5.0 eq.) and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 4.6 mL, 3.0 eq.) were added. The reaction was shaken at room temperature for 12 - 16 hours, then filtered and washed with DMF (2 x 30 mL), DMF:water 9:1 (2 x 30 mL), DMF (30 mL), MeOH (2 x 20 mL), and DCM (2 x 20 mL). The resin was dried *in vacuo* at room temperature.

Step B: Coupling of 2-(4-fluorobenzenesulfanyl)-octanoic acid hydroxyamide resin with benzyl alcohol.

15 2-(4-Fluorobenzenesulfanyl)-octanoic acid hydroxyamide resin prepared in Step A (330 mg, 1.1 meq/g) was suspended in DMF (2.0 mL) and benzyl alcohol (731 mg, 15 eq.) and sodium hydride (237 mg, 15 eq.) were added. The reaction was heated to 80°C for 15 hours while shaking on an orbital shaker. After cooling to room temperature the mixture was filtered and washed with DMF (2 x 2 mL), DMF:water 9:1 (2 x 3 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step C: Oxidation of sulfide to sulfoxide.

25 2-(4-Benzyloxy-phenylsulfanyl)-octanoic acid hydroxyamide resin prepared in Step B (110 mg, 1.1 meq/g) was suspended in DCM (2.2 mL) and 70% *tert*-butylhydroperoxide (0.73 mL) benzenesulfonic acid (36 mg) were added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step D: Oxidation of sulfide to sulfone.

30 2-(4-Benzyloxy-phenylsulfanyl)-octanoic acid hydroxyamide resin prepared in Step B (110 mg, 1.1 meq/g) was suspended in DCM (2.2 mL) and mCPBA (132 mg) was added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 2 mL), DMF (2 x 2 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

WO 99/42436

PCT/US98/17633

- 133 -

Step E: Cleavage of 2-(4-benzyloxy-benzenesulfanyl)-octanoic acid hydroxyamide from resin.

The 2-(4-benzyloxy-phenylsulfanyl)-octanoic acid hydroxyamide resin prepared in Step B (110 mg, 1.2 meq/g) was suspended in DCM (1.0 mL) and TFA (1.0 mL) was added. The reaction was shaken for 1 hour at room temperature. The reaction was filtered and the resin washed with DCM (2 x 1 mL). The filtrate and the washing were combined and concentrated to dryness on a Savant SpeedVac Plus. Methanol (1 mL) was added and the mixture concentrated. The crude product was dissolved in DMSO:methanol (1:1, 2 mL) and purified by reverse phase HPLC under the conditions described below:

Column: ODS-A, 20mm x 50 mm, 5 μ m particle size (YMC, Inc. Wilmington, North Carolina)

Solvent Gradient	Time	Water	Acetonitrile
	0.0	95	5
	25 min.	5	95

Flow Rate: 15 mL/min.

2-(4-Benzyloxy-phenylsulfanyl)-octanoic acid hydroxyamide 100% @ 215 nm; LCMS (API-electrospray) m/z 374 (M+H)⁺.

The hydroxamic acid compounds of Examples 200-220 are synthesized using appropriate starting materials and following the steps in example 199:

Example 200

2-(4-Butoxy-benzenesulfanyl)-octanoic acid hydroxyamide 100% @ 215 nm; LCMS (API-electrospray) m/z 374 (M+H)⁺.

Example 201

2-[4-(2-Piperazine-1-yl-ethoxy)-benzenesulfanyl]-octanoic acid hydroxyamide 98% @ 215 nm; LCMS (API-electrospray) m/z 340 (M+H)⁺.

Example 202

2-[4-(5-Hydroxy-pentyloxy)-phenylsulfanyl]-octanoic acid hydroxyamide 65% @ 215 nm.; LCMS (API-electrospray) m/z 370 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 134 -

Example 203

2-[4-(3-Pyridin-2-yl-propoxy)-benzenesulfanyl]-octanoic acid hydroxyamide
95% @ 215 nm; LCMS (API-electrospray) m/z 403 (M+H)⁺.

5

Example 204

2-(4-Benzyloxy-phenylsulfinyl)-octanoic acid hydroxyamide 100% @ 215 nm.

10

Example 205

2-(4-Butoxy-benzenesulfinyl)-octanoic acid hydroxyamide 98% @ 215 nm.

15

Example 206

2-[4-(2-Piperazine-1-yl-ethoxy)-benzenesulfinyl]-octanoic acid hydroxyamide
98% @ 215 nm.

20

Example 207

2-[4-(3-Pyridin-2-yl-propoxy)-benzenesulfinyl]-octanoic acid hydroxyamide
99% @ 215 nm.

25

Example 208

2-(4-Benzyloxy-phenylsulfonyl)-octanoic acid hydroxyamide 100% @ 215 nm.

Example 209

30 2-(4-Butoxy-benzenesulfonyl)-octanoic acid hydroxyamide 100% @ 215 nm.

Example 210

35 2-[4-(2-Piperazine-1-yl-ethoxy)-benzenesulfonyl]-octanoic acid hydroxyamide
97% @ 215 nm.

Example 211

40 2-[4-(3-Pyridin-2-yl-propoxy)-benzenesulfonyl]-octanoic acid hydroxyamide
100% @ 215 nm.

Example 212

2-[4-(1-Methyl-pyrrolidin-3-yloxy)-benzenesulfanyl]-octanoic acid hydroxyamide
91% @ 215 nm; LCMS (API-electrospray) m/z 367 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 135 -

Example 213

2-[4-(1-Ethyl-propoxy)-benzenesulfanyl]-octanoic acid hydroxyamide
5 100% @ 215 nm; LCMS (API-electrospray) m/z 354 (M+H)⁺.

Example 214

2-[4-(Tetrahydro-pyran-4-yloxy)-benzenesulfanyl]-octanoic acid hydroxyamide
10 97% @ 215 nm; LCMS (API-electrospray) m/z 368 (M+H)⁺.

Example 215

2-[4-(1-Methyl-pyrrolidin-3-yloxy)-benzenesulfinyl]-octanoic acid hydroxyamide
15 96% @ 215 nm.

Example 216

2-[4-(1-Ethyl-propoxy)-benzenesulfinyl]-octanoic acid hydroxyamide 97% @ 215 nm.
20

Example 217

2-[4-(Tetrahydro-pyran-4-yloxy)-benzenesulfinyl]-octanoic acid hydroxyamide
97% @ 215 nm.
25

Example 218

2-[4-(1-Methyl-pyrrolidin-3-yloxy)-benzenesulfonyl]-octanoic acid hydroxyamide
96% @ 215 nm.
30

Example 219

2-[4-(1-Ethyl-propoxy)-benzenesulfonyl]-octanoic acid hydroxyamide
100% @ 215 nm.
35

Example 220

2-[4-(Tetrahydro-pyran-4-yloxy)-benzenesulfonyl]-octanoic acid hydroxyamide
100% @ 215 nm.

WO 99/42436

PCT/US98/17633

- 136 -

Example 221

Step A: Displacement of bromide with 4-bromothiophenol.

- The 2-bromo-octanoic acid hydroxamate resin prepared in Example 160, Step A (5.0 g, 1.1 meq/g) was placed in a peptide synthesis vessel and suspended in THF (60 mL). 4-Bromothiophenol (5.2 g, 5.0 eq.), sodium iodide (4.1 g, 5.0 eq.) and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 2.5 mL, 3.0 eq.) were added. The reaction was shaken at room temperature for 12 - 16 hours, then filtered and washed with DMF (2 x 30 mL), DMF:water 9:1 (2 x 30 mL), DMF (30 mL), MeOH (2 x 30 mL), and DCM (2 x 30 mL). The resin was dried *in vacuo* at room temperature.

Step B: Oxidation of sulfide to sulfoxide.

- 2-(4-Bromobenzenesulfanyl)-octanoic acid hydroxyamide resin prepared in Step A (4.4 g, 1.1 meq/g) was suspended in DCM (60 mL) and 70% *tert*-butylhydroperoxide (30 mL) benzenesulfonic acid (1.5 g) were added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 30 mL), DMF (2 x 30 mL), MeOH (2 x 30 mL), and DCM (2 x 30 mL). The resin was dried *in vacuo* at room temperature.

Step C: Oxidation of sulfide to sulfone.

- 2-(4-Bromobenzenesulfanyl)-octanoic acid hydroxyamide resin prepared in Step B (4.4 g, 1.1 meq/g) was suspended in DCM (60 mL) and mCPBA (5.2 g) was added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 30 mL), DMF (2 x 30 mL), MeOH (2 x 30 mL), and DCM (2 x 30 mL). The resin was dried *in vacuo* at room temperature.

Step D: Coupling of 2-(4-bromobenzenesulfinyl)-octanoic acid hydroxyamide resin with 4-chlorobenzeneboronic acid.

- 2-(4-Bromobenzenesulfinyl)-octanoic acid hydroxyamide resin prepared in Step B (150 mg, 1.1 meq/g) was suspended in DME (2.0 mL) and nitrogen gas bubbled through the suspension for 1-2 minutes. 4-Chlorobenzeneboronic acid (51.6 mg, 2 eq.), tetrakis(triphenylphosphine) palladium(0) (19.07 mg, 0.1 eq.) and sodium carbonate (2 M solution, 0.825 mL, 10 eq.) were added. The reaction was heated to 80°C for 8 hours while shaking on an orbital shaker. After cooling to room temperature the mixture was filtered and washed with DME (2 x 2 mL), DMF:water 9:1 (2 x 3 mL),

WO 99/42436

PCT/US98/17633

- 137 -

MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

5 Step E: Cleavage of 2-(4'-chloro-biphenyl-4-sulfinyl)-octanoic acid hydroxyamide from resin.

The 2-(4'-chloro-biphenyl-4-sulfinyl)-octanoic acid hydroxyamide resin prepared in Step D (150 mg, 1.1 meq/g) was suspended in DCM (1.0 mL) and TFA (1.0 mL) was added. The reaction was shaken for 1 hour at room temperature. The reaction was filtered and the resin washed with DCM (2 x 1 mL). The filtrate and the washing were
10 combined and concentrated to dryness on a Savant SpeedVac Plus. Methanol (1 mL) was added and the mixture concentrated. The crude product was dissolved in DMSO:methanol (1:1, 2 mL) and purified by reverse phase HPLC under the conditions described below:

15 Column: ODS-A, 20mm x 50 mm, 5 μ m particle size (YMC, Inc. Wilmington, North Carolina)

Solvent Gradient	Time	Water	Acetonitrile
	0.0	95	5
	25 min.	5	95

20 Flow Rate: 15 mL/min.

2-(4'-Chloro-biphenyl-4-sulfinyl)-octanoic acid hydroxyamide 96% @ 215 nm; LCMS (API-electrospray) m/z 394 (M+H)⁺.

25 The hydroxamic acid compounds of Exmples 222-224 are synthesized using appropriate starting materials and following the steps in example 221:

Example 222

2-[4-(5-Chloro-thiophen-2-yl)-benzenesulfinyl]-octanoic acid hydroxyamide 100% @
30 215 nm; LCMS (API-electrospray) m/z 400 (M+H)⁺.

Example 223

2-(4'-Chloro-biphenyl-4-sulfonyl)-octanoic acid hydroxyamide 94% @ 215 nm;
35 LCMS (API-electrospray) m/z 410 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 138 -

Example 224

2-[4-(5-Chloro-thiophen-2-yl)-benzenesulfonyl]-octanoic acid hydroxyamide 85% @ 215 nm; LCMS (API-electrospray) m/z 416 (M+H)⁺.

5

Example 225

Step A: Coupling of 2-(4-bromobenzenesulfonyl)-octanoic acid hydroxyamide resin with *N*-(3-aminopropyl)-morpholine.

- 10 2-(4-Bromobenzenesulfonyl)-octanoic acid hydroxyamide resin prepared in Example 199, Step A (100 mg, 1.1 meq/g) was suspended in dioxane (2.0 mL) and nitrogen gas bubbled through the suspension for 1-2 minutes. *N*-(3-Aminopropyl)-morpholine (346 mg, 20 eq.), tris(dibenzylideneacetone)-dipalladium(0) (22 mg, 0.2 eq.), (S)-(-)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl((S)-BINAP, 60 mg, 0.8 eq.) and sodium *tert*-butoxide (207 mg, 18 eq.) were added. The reaction was heated to 80°C for 8 hours
- 15 while shaking on an orbital shaker. After cooling to room temperature the mixture was filtered and washed with DMF (2 x 2 mL), DMF:water 9:1 (2 x 3 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

- 20 Step B: Cleavage of 2-[4-(3-morpholin-4-yl-propylamino)-phenylsulfonyl]-octanoic acid hydroxyamide from resin.

The 2-[4-(3-morpholin-4-yl-propylamino)-phenylsulfonyl]-octanoic acid hydroxyamide resin prepared in Step A (100 mg, 1.1 meq/g) was suspended in DCM (1.0 mL) and TFA (1.0 mL) was added. The reaction was shaken for 1 hour at room temperature.

- 25 The reaction was filtered and the resin washed with DCM (2 x 1 mL). The filtrate and the washing were combined and concentrated to dryness on a Savant SpeedVac Plus. Methanol (1 mL) was added and the mixture concentrated. The crude product was dissolved in DMSO:methanol (1:1, 2 mL) and purified by reverse phase HPLC under the conditions described below:

30

Column: ODS-A, 20mm x 50 mm, 5 µm particle size (YMC, Inc. Wilmington, North Carolina)

Solvent Gradient	Time	Water	Acetonitrile
	0.0	95	5
35	25 min.	5	95

Flow Rate: 15 mL/min.

2-[4-(3-morpholin-4-yl-propylamino)-phenylsulfonyl]-octanoic acid hydroxyamide 88% @ 215 nm; LCMS (API-electrospray) m/z 410 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 139 -

The hydroxamic acid compounds of Examples 226-231 are synthesized using appropriate starting materials and following the steps in this example:

5

Example 226

2-[4-(Biphenyl-4-ylamino)-phenylsulfanyl]-octanoic acid hydroxyamide 95% @ 215 nm; LCMS (API-electrospray) m/z 435 (M+H)⁺.

10

Example 227

2-[4-(Pyridin-4-ylamino)-phenylsulfanyl]-octanoic acid hydroxyamide 97% @ 215 nm; LCMS (API-electrospray) m/z 360 (M+H)⁺.

15

Example 228

2-(4-Cyclopentylamino-phenylsulfanyl)-octanoic acid hydroxyamide 77% @ 215 nm; LCMS (API-electrospray) m/z 351 (M+H)⁺.

20

Example 229

2-(4-Methylamino-phenylsulfanyl)-octanoic acid hydroxyamide 99% @ 215 nm; LCMS (API-electrospray) m/z 297 (M+H)⁺.

25

Example 230

2-(4-Piperidin-1-yl-phenylsulfanyl)-octanoic acid hydroxyamide 72% @ 215 nm; LCMS (API-electrospray) m/z 351 (M+H)⁺.

30

Example 231

2-(4-Piperazin-1-yl-phenylsulfanyl)-octanoic acid hydroxyamide 74% @ 215 nm; LCMS (API-electrospray) m/z 352 (M+H)⁺.

35

Example 232

Step A: Displacement of bromide with 4-hydroxythiophenol.

The 2-bromo-octanoic acid hydroxamate resin prepared in Example 160, Step A (15.0 g, 1.1 meq/g) was placed in a peptide synthesis vessel and suspended in THF (120 mL). 4-Hydroxythiophenol (11.3 g, 5.0 eq.), sodium iodide (13.5 g, 5.0 eq.) and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 8.1 mL, 3.0 eq.) were added. The reaction

WO 99/42436

PCT/US98/17633

- 140 -

was shaken at room temperature for 12 - 16 hours, then filtered and washed with DMF (2 x 60 mL), DMF:water 9:1 (2 x 60 mL), DMF (60 mL), MeOH (2 x 60 mL), and DCM (2 x 60 mL). The resin was dried *in vacuo* at room temperature.

- 5 Step B: Coupling of 2-(4-hydroxybenzenesulfanyl)-octanoic acid hydroxyamide resin with benzene sulfonyl chloride.
- 2-(4-Hydroxybenzenesulfanyl)-octanoic acid hydroxyamide resin prepared in Step A (240 mg, 1.2 meq/g) was suspended in DCM (3.0 mL). Benzene sulfonyl chloride (225 mg, 5 eq.), and triethylamine (0.06 mL, 2 eq.) were added. The reaction was
- 10 shaken on an orbital shaker at room temperature for 8 hours, then filtered and washed with DME (2 x 2 mL), DMF:water 9:1 (2 x 3 mL), MeOH (2 x 2 mL), and DCM (2 x 2 mL). The resin was dried *in vacuo* at room temperature.

Step C: Oxidation of sulfide to sulfoxide.

- 15 Benzenesulfonic acid 4-(1-hydroxycarbamoyl-heptylsulfanyl)-phenyl ester resin prepared in Step B (80 mg, 1.2 meq/g) was suspended in DCM (3 mL) and 70% *tert*-butylhydroperoxide (1 mL) benzenesulfonic acid (23 mg) were added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 3 mL), DMF (2 x 3 mL), MeOH (2 x
- 20 3 mL), and DCM (2 x 3 mL). The resin was dried *in vacuo* at room temperature.

Step D: Oxidation of sulfide to sulfone.

- Benzenesulfonic acid 4-(1-hydroxycarbamoyl-heptylsulfanyl)-phenyl ester resin prepared in Step B (80 mg, 1.2 meq/g) was suspended in DCM (3 mL) and mCPBA
- 25 (84 mg) was added. The reaction mixture was shaken on an orbital shaker at room temperature for 12 - 24 hours. The reaction was filtered and washed with DCM (2 x 3 mL), DMF (2 x 3 mL), MeOH (2 x 3 mL), and DCM (2 x 3 mL). The resin was dried *in vacuo* at room temperature.

- 30 Step E: Cleavage of benzenesulfonic acid 4-(1-hydroxycarbamoyl-heptylsulfanyl)-phenyl ester resin.
- The benzenesulfonic acid 4-(1-hydroxycarbamoyl-heptylsulfanyl)-phenyl ester resin prepared in Step B (80 mg, 1.2 meq/g) was suspended in DCM (1.0 mL) and TFA (1.0 mL) was added. The reaction was shaken for 1 hour at room temperature. The
- 35 reaction was filtered and the resin washed with DCM (2 x 1 mL). The filtrate and the washing were combined and concentrated to dryness on a Savant SpeedVac Plus. Methanol (1 mL) was added and the mixture concentrated. The crude product was

WO 99/42436

PCT/US98/17633

- 141 -

dissolved in DMSO:methanol (1:1, 2 mL) and purified by reverse phase HPLC under the conditions described below:

Column: ODS-A, 20mm x 50 mm, 5 µm particle size (YMC, Inc. Wilmington, North

5 Carolina)

Solvent Gradient	Time	Water	Acetonitrile
	0.0	95	5
	25 min.	5	95

Flow Rate: 15 mL/min.

10 Benzenesulfonic acid 4-(1-hydroxycarbamoyl-heptylsulfanyl)-phenyl ester 91% @ 215 nm; LCMS (API-electrospray) m/z 424 (M+H)⁺.

The hydroxamic acid compounds of Examples 233-240 are synthesized using appropriate starting materials and following the steps in example 232:

15

Example 233

2,5-Dichloro-thiophene-3-sulfonic acid 4-(1-hydroxycarbamoyl-heptylsulfanyl)-hydroxyamide 98% @ 215 nm; LCMS (API-electrospray) m/z 498 (M+H)⁺.

20

Example 234

Ethanesulfonic acid 4(1-hydroxycarbamoyl-heptylsulfanyl)-hydroxyamide. 72% @ 215 nm; LCMS (API-electrospray) m/z 376 (M+H)⁺.

25

Example 235

5-Chloro-1,3-dimethyl-1H-pyrazole-4-sulfonic acid 4-(1-hydroxycarbamoyl-heptylsulfinyl)-hydroxyamide 99% @ 215 nm; LCMS (API-electrospray) m/z 492 (M+H)⁺.

30

Example 236

2,5-Dichloro-thiophene-3-sulfonic acid 4-(1-hydroxycarbamoyl-heptylsulfinyl)-hydroxyamide 96% @ 215 nm; LCMS (API-electrospray) m/z 514 (M+H)⁺.

35

Example 237

5-Pyridin-2-yl-thiophene-2-sulfonic acid 4(1-hydroxycarbamoyl-heptylsulfinyl)-hydroxy amide 96% @ 215 nm; LCMS (API-electrospray) m/z 523 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 142 -

Example 238

2-Nitro-benzenesulfonic acid 4-(1-hydroxycarbamoyl-heptylsulfonyl)-hydroxyamide
5 97% @ 215 nm; LCMS (API-electrospray) m/z 501 (M+H)⁺.

Example 239

3-Bromo-2-chloro-thiophene-2-sulfonic acid 4-(1-hydroxycarbamoyl-heptylsulfonyl)-
10 hydroxyamide 97% @ 215 nm; LCMS (API-electrospray) m/z 576 (M+H)⁺.

Example 240

Benzo[1,2,5]thiadiazole-4-sulfonic acid 4-(1-hydroxycarbamoyl-heptylsulfonyl)-
15 hydroxyamide 83% @ 215 nm; LCMS (API-electrospray) m/z 514 (M+H)⁺.

Example 241

20 1-Benzyl-4-(4-benzyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide

To a stirred solution of 4-methoxybenzenethiol (2.8 gm, 20 mmol) and anhydrous
K₂CO₃ (10 gm, excess) in dry acetone (100 ml), α-bromo ethyl acetate (3.3 gm, 20
mmol) was added in a round bottom flask and the reaction mixture was heated at reflux
for 8 hours with good stirring. At the end, the reaction mixture was allowed to cool and
25 the potassium salts were filtered off and the reaction mixture was concentrated. The
residue was extracted with chloroform and washed with H₂O and 0.5 N NaOH
solution. The organic layer was further washed well with water, dried over MgSO₄,
filtered and concentrated. (4-methoxy-phenylsulfonyl)-acetic acid ethyl ester was
isolated as pale yellow oil. Yield: 4.4 g (100%); MS; 227 (M+H)⁺

30 To stirred solution of (4-methoxy-phenylsulfonyl)-acetic acid ethyl ester (4.4 g, 20
mmol), anhydrous K₂CO₃ (10 gm, excess) in dry acetone (100 ml) benzyl bromide
(3.5 g, 20 mmol) was added and refluxed for 4 hrs. At the end, reaction mixture was
filtered, concentrated and the residue was extracted with chloroform. It was washed
35 well with water, dried and concentrated. The crude product obtained was converted to
(4-benzyloxy-phenylsulfonyl)-acetic acid ethyl ester by oxidizing with m-chloro
perbenzoic acid as described in the example 83. Low melting solid. Yield: 6.6 g, 97%;
MS: 335 (M+1)

WO 99/42436

PCT/US98/17633

- 143 -

To a stirred solution of bis-(2-chloro-ethyl)-benzyl amine hydrochloride (6.6 g, 24.7 mmol), 18-Crown-6 (500 mg), and anhydrous K₂CO₃ (30 gm, excess) in dry acetone (250 ml), (4-benzyloxy-phenylsulfonyl)-acetic acid ethyl ester (8.01 gm, 24 mmol) was added in a round bottom flask and the reaction mixture was heated at reflux for 16 hours with good stirring. At the end, the reaction mixture was allowed to cool and the potassium salts were filtered off and the reaction mixture was concentrated. The residue was extracted with chloroform and washed with H₂O. The organic layer was further washed well with water, dried over MgSO₄, filtered and concentrated. The dark brown reaction mixture was purified by silica gel column chromatography by eluting it with 30% ethylacetate: hexane and the product 4-(4-Benzyloxy-benzenesulfonyl)-1-benzyl-piperidine-4-carboxylic acid ethyl ester was isolated as Brown oil. Yield: 6.5 g, 55%; MS: 494 (M+H)

4-(4-Benzyloxy-benzenesulfonyl)-1-benzyl-piperidine-4-carboxylic acid ethyl ester (5.0 g, 10.1 mmol) was dissolved in MeOH/THF (1:1, 200 ml) and stirred at room temperature for 72 hrs. At the end reaction mixture was concentrated and the product was neutralised with con. HCl by dissolving it in water (200 ml). After the neutralization reaction mixture was concentrated to dryness. Ice cold water (100 ml) was added to the solid and filtered. The product 4-(4-Benzyloxy-benzenesulfonyl)-1-benzyl-piperidine-4-carboxylic acid was dried at 50 °C and taken to next step without any purification. Colorless solid. MP: 66-68; Yield: 4.3 g, 91%; MS: 466 (M+H) Starting from 4-(4-Benzyloxy-benzenesulfonyl)-1-benzyl-piperidine-4-carboxylic acid (4.65 g, 10.0 mmol) and following the procedure outlined in example 83, 1.1 g of 4-(4-Benzyloxy-benzenesulfonyl)-1-benzyl-piperidine-4-carboxylic acid hydroxyamide was isolated as a colorless solid. Yield 21%; mp 89 °C; MS: 481.1 1H NMR (300 MHz, DMSO-d₆): δ 2.27 (m, 3H), 2.76-2.79 (m, 2H), 3.43 (m, 4H), 4.30 (s, 2H), 7.14-7.17 (d, 2H), 7.50-7.73 (m, 5H), 9.37 (s, 1H), 10.53 (s, 1H), 11.18 (s, 1H).

Example 242

4-(4-Butoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid hydroxyamide

From 2-[(2-Hydroxy-ethyl)-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-amine was prepared according to the general method outlined in example 83 starting from diethanolamine (15.0 g, 150), and 4-(2-piperidin-1-yl-ethoxy)-benzyl chloride (5.9 g, 20 mmol). Yield 5.5 g, (85%); Brown semi-solid; MS: 323 (M+H)⁺

WO 99/42436

PCT/US98/17633

- 144 -

Bis-(2-chloro-ethyl)-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-amine was prepared according to the general method outlined in example 83 starting from 2-[(2-Hydroxy-ethyl)- [4-(2-piperidin-1-yl-ethoxy)-benzyl]-amine (3.22 g, 10 mmol). Yield 4.0 g (92%); brown semi-solid; MS: 361.1 (M+H)⁺

5

4-(4-Butoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting from 4-(butoxy-benzenesulfonyl) acetic acid ethyl ester (6.0 g, 20 mmol) and Bis-(2-chloro-ethyl)-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-amine (8.6 g, 20 mmol). Yield 8.0 g (68%); brown oil; MS: 587.7 (M+H)⁺

10

4-(4-Butoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid was prepared starting from 4-(4-Butoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid ethyl ester (5.8 g, 10 mmol) dissolved in THF:methanol 3:1 and 10 N NaOH (40 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 4.8 g (86%); Spongy brown solid; mp 98 °C; MS: 559.6 (M+H)⁺

15

Starting from 4-(4-butoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid (5.5 g, 10 mmol) and following the procedure outlined in example 83, 2.4 g of 4-(4-butoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid hydroxy amide was isolated as a pale yellow solid. Yield 41%; mp 155 °C (HCl); MS: 574 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.9 (t, 3H), 1.1-1.8 (m, 6H), 1.9 (m, 4H), 2.3 (m, 4H), 2.8 (m, 6H), 3.2-3.6 (m, 8H), 4.2 (m, 2H), 6.9 - 7.8 (m, 8H), 9.1 (s, 1H), 10.8 (bs, 1H).

20

25

Example 243

4-(4-Butoxy-benzenesulfonyl)-1-[3-(2-morpholinyl-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid hydroxyamide

30

Bis-(2-hydroxy-ethyl)-[3-(2-morpholin-1-yl-ethoxy)-benzyl]-amine was prepared according to the general method outlined in example 83 starting from diethanolamine (15.0 g, 150) and 3-(2-morpholin-1-yl-ethoxy)-benzyl chloride (5.9 g, 20 mmol). Yield 6.2 g, (95%); Brown semi-solid; MS: 325 (M+H)⁺

35

WO 99/42436

PCT/US98/17633

- 145 -

Bis-(2-chloro-ethyl)-[3-(2-morpholin-1-yl-ethoxy)-benzyl]-amine was prepared according to the general method outlined in example 83 starting from Bis-(2-Hydroxy-ethyl)-[3-(2-morpholin-1-yl-ethoxy)-benzyl]-amine (3.24 g, 10 mmol). Yield 4.0 g (92%); brown semi-solid; MS: 363.1 (M+H)⁺

5

4-(4-Butoxy-benzenesulfonyl)-1-[3-(2-morpholin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting from 4-(butoxy-benzenesulfonyl) acetic acid ethyl ester (6.0 g, 20 mmol) and Bis-(2-chloro-ethyl)-[3-(2-morpholin-1-yl-ethoxy)-benzyl]-amine (8.6 g, 20 mmol). Yield 8.5 g (72%); brown oil; MS: 589.7 (M+H)⁺

10

4-(4-Butoxy-benzenesulfonyl)-1-[3-(2-morpholin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid was prepared starting from 4-(4-Butoxy-benzenesulfonyl)-1-[3-(2-morpholin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid ethyl ester (5.8 g, 10 mmol) dissolved in THF:methanol 3:1 and 10 N NaOH (40 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 4.8 g (85%); Spongy brown solid; MS: 561.6 (M+H)⁺

15

Starting from 4-(4-butoxy-benzenesulfonyl)-1-[3-(2-morpholin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid (5.6 g, 10 mmol) and following the procedure outlined in example 83, 4.02 g of 4-(4-butoxy-benzenesulfonyl)-1-[3-(2-morpholin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid hydroxy amide was isolated as a pale yellow solid. Yield 62%; mp 123 °C (HCl); MS: 576 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.9 (t, 3H), 1.4 (m, 2H), 1.8 (t, 2H), 2.3-4.7 (m, 24H), 7.0- 7.8 (m, 8H), 9.1 (s, 1H), 10.8 (bs, 1H).

25

Example 244

1-Methyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide

30

1-Methyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting from 4-(butoxy-benzenesulfonyl) acetic acid ethyl ester (3 g, 10 mmol) and methyl-bis-(2-chloro-ethyl)-amine (2.2g, 11.6 mmol). Yield 4.0 g, (98%); low melting brown solid; MS: 384 (M+H)⁺

35

WO 99/42436

PCT/US98/17633

- 146 -

1-Methyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-methyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (7.6 g, 20 mmol) dissolved in methanol (300 ml) and 10 N NaOH (35 ml). The resulting reaction mixture was worked up outlined in example 83. Yield 6.0 g (84%);
5 white solid; mp 195 °C; MS: 356.4 (M+H)⁺

Starting from 1-methyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid (4.0 g, 11.2 mmol) and following the procedure outlined in example 83, 3.9 g of 1-methyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated
10 as a yellow powder. Yield 85%; mp 118 °C; MS: 371 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.9 (t, 3H), 1.45 (q, 2H), 1.8 (q, 2H), 2.1 (s, 3H), 2.3 (d, J = 11.4 Hz, 2H), 2.5-3.7 (m, 8H), 4.1 (t, 2H), 7.16 (d, 2H), 7.67 (d, 2H)

Example 245

15 1-Ethyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide

1-Ethyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting from 4-(butoxy-benzenesulfonyl) acetic acid ethyl ester (3 g, 10 mmol) and ethyl-bis-(2-chloro-ethyl)-amine (2.2g, 10.6 mmol). Yield 3.5 g, (88%); low melting brown solid;
20 MS: 398 (M+H)⁺

1-Ethyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-ethyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (7.94 g, 20 mmol) dissolved in methanol (300 ml) and 10 N NaOH (35 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 6.5 g (88%); white solid; mp 162 °C; MS: 370 (M+H)⁺

25 Starting from 1-ethyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid (3.7 g, 10 mmol) and following the procedure outlined in example 83, 3.2 g of 1-ethyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as a yellow powder. Yield 76%; mp 98 °C; MS: 385 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.9 (t, 3H), 1.2 (t, 3H), 1.46 (q, 2H), 1.9 (q, 2H), 2.3 (d, J = 11.4 Hz, 2H), 2.5-3.6 (m, 10H), 4.1 (t, 2H), 7.16 (d, 2H), 7.67 (d, 2H)
35

WO 99/42436

PCT/US98/17633

- 147 -

Example 246

1-n-Butyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide

5 1-n-Butyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting from 4-(butoxy-benzenesulfonyl) acetic acid ethyl ester (3 g, 10 mmol) and n-butyl-bis-(2-chloro-ethyl)-amine (2.0g, 10.1 mmol). Yield 3.8 g, (89%); low melting brown solid; MS: 426 (M+H)⁺

10

1-n-Butyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-n-Butyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (8.5 g, 20 mmol) dissolved in methanol (300 ml) and 10 N NaOH (35 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 7.5 g
15 (88%); white solid; mp 182 °C; MS: 398 (M+H)⁺

Starting from 1-n-butyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid (3.9 g, 10 mmol) and following the procedure outlined in example 83, 1.8 g of 1-n-butyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was
20 isolated as a yellow powder. Yield 40%; mp 121 °C; MS: 413 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.9-1.0 (m, 6H), 1.2-1.8 (m, 8H), 2.2-2.8 (m, 8H), 3.0-3.6 (m, 4H), 4.2 (t, 2H), 7.16 (d, 2H), 7.67 (d, 2H), 9.3 (bs, 1H), 10.3 (bs, 1H), 11.1 (bs, 1H).

25

Example 247

[4-(4-Chloro-phenoxy)-phenylsulfanyl]-acetic acid ethyl ester

A mixture of 4-bromo chlorobenzene (1.92 g, 10 mmol), (4-Hydroxy-phenylsulfanyl)-
30 acetic acid ethyl ester (2.12 g, 10 mmol), sodium hydride (460 mg, 10 mmol) and copper(II) chloride (500 mg) was refluxed in anhydrous pyridine (50 ml) for 12 hrs. The reaction mixture was carefully quenched with ice cold water and acidified with concentrated HCl. The product was extracted with chloroform, washed well with water; dried and concentrated. The product was purified by silica gel column chromatography
35 by eluting with 30% ethyl acetate : hexane. Yield 2.5 g (77%); Colorless low melting solid; MS: 323 (M+H)⁺. Alternatively the title compound may be prepared from 4-(4-chloro-phenoxy)-benzenethiol and bromo ethyl acetate as described in example 83.

WO 99/42436

PCT/US98/17633

- 148 -

Example 248

[4-(4-Chloro-phenoxy)-benzenesulfonyl]-acetic acid ethyl ester

- 5 [4-(4-Chloro-phenoxy)-benzenesulfonyl]-acetic acid ethyl ester was prepared according to the general method outlined in example 83 starting from [4-(4-chloro-phenoxy)-phenylsulfanyl]-acetic acid ethyl ester (3.23g, 10 mmol) and oxone (10 g). Yield 3.5g (99%); oil; MS: 356EI (M+H)⁺

10

Example 249

4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-1-methyl-piperidine-4-carboxylic acid hydroxyamide

- 15 4-[4-(4-Chloro-phenoxy)-benzenesulfonyl]-1-methyl-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting from [4-(4-chloro-phenoxy)-benzenesulfonyl]-acetic acid ethyl ester (2.0g, 5.6mmol) and mechlorethanamine hydrochloride (Aldrich), (1.9g, 10mmol). Yield 2g (81%); brown oil; MS:438 (M+H)⁺

20

4-[4-(4-Chloro-phenoxy)-benzenesulfonyl]-1-methyl-piperidine-4-carboxylic acid was prepared starting from 4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-1-methyl-piperidine-4-carboxylic acid ethyl ester (4.3 g, 10 mmol) dissolved in THF:methanol (3:1 150 ml) and 10 N NaOH (100 ml). The resulting reaction mixture was worked up as outlined

25 in example 83. Yield 3.5g (86 %); white solid; mp 185°C; MS:410(M+H)⁺

- Starting from 4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-1-methyl-piperidine-4-carboxylic acid (1.0 g, 2.4 mmol) and following the procedure outlined in example 83, 460mg of 4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-1-methyl-piperidine-4-carboxylic acid hydroxyamide was isolated as a HCl salt, a white powder. Yield 41%; mp 52 °C;
- 30 MS: 426 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 1.3 (s, 3H), 2.2 - 2.9 (m, 6H), 3.5 (bd, 2H), 7.2 -7.9 (m, 8H), 8.1 (s, 1H), 11.0 (bs, 1H).

WO 99/42436

PCT/US98/17633

- 149 -

Example 250

4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-1-ethyl-piperidine-4-carboxylic acid
hydroxyamide

5

4-[4-(4-Chloro-phenoxy)-benzenesulfonyl]-1-ethyl-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting from [4-(4-chloro-phenoxy)-benzenesulfonyl]-acetic acid ethyl ester (4 g, 11.3 mmol) and ethyl-bis-(2-chloro-ethyl)-amine (2.32 g, 16.9 mmol). Yield 3.36 g (66%); brown oil; MS: 452.0 (M+H)⁺

10

4-[4-(4-Chloro-phenoxy)-benzenesulfonyl]-1-ethyl-piperidine-4-carboxylic acid was prepared starting from 4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-1-ethyl-piperidine-4-carboxylic acid ethyl ester (3.02 g, 6.7 mmol) dissolved in THF:methanol (3:1 150 ml) and 10 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 1.8 g (65%); white solid; mp 184°C; MS: 423.9 (M+H)⁺

15

Starting from 4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-1-ethyl-piperidine-4-carboxylic acid (1.75 g, 4.14 mmol) and following the procedure outlined in example 83, 650 mg of 4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-1-ethyl-piperidine-4-carboxylic acid hydroxyamide was isolated as a HCl salt, a white powder. Yield 33%; mp 158 °C; MS: 438.9 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 1.78 (t, J=7.23 Hz, 3H), 2.23-2.27 (m, 2H), 2.51-2.73 (m, 4H), 3.04 (m, 2H), 3.81 (d, J=24 Hz, 2H), 7.16-7.27 (m, 4H), 7.50-7.57 (m, 2H), 7.76 (d, J = 7 Hz, 2H), 9.34 (s, 1H), 9.85 (s, 1H).

25

Example 251

1-Butyl-4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid
hydroxyamide

30

1-Butyl-4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting from [4-(4-chloro-phenoxy)-benzenesulfonyl]-acetic acid ethyl ester (6 g, 18.3 mmol) and butyl-bis-(2-chloro-ethyl)-amine (5.2 g, 22 mmol). Yield 3.3 g (38%); yellow oil; MS: 480 (M+H)⁺

35

WO 99/42436

PCT/US98/17633

- 150 -

1-Butyl-4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid was prepared starting from 1-butyl-4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid ethyl ester (3.3 g, 6.9 mmol) dissolved in THF:methanol (3:1 150 ml) and 10 N NaOH (25 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 2.08 g (67%); white solid; mp 201°C; MS: 451.9 (M+H)⁺

Starting from 1-butyl-4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid (2 g, 4.43 mmol) and following the procedure outlined in example 83, 630 mg of 1-butyl-4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide was isolated as a HCl salt, a white solid. Yield 31%; mp 212 °C; MS: 466.9 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.87 (t, J=7.3 Hz, 3H), 1.32 (m, 2H), 1.60 (m, 2H), 2.21 (m, 2H), 2.50 (m, 2H), 2.70 (q, 2H), 3.00 (m, 2H), 3.57 (d, 2H), 7.16-7.26 (m, 4H), 7.49-7.56 (m, 2H), 7.77 (d, J = 9 Hz, 2H), 9.34 (s, 1H), 10.13 (s, 1H).

Example 252

1-Benzyl-4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide

1-Benzyl-4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting from [4-(4-chloro-phenoxy)-benzenesulfonyl]-acetic acid ethyl ester (6 g, 16.9 mmol) and bis-(2-chloro-ethyl)-benzyl amine (6.44 g, 24 mmol). Yield 2.21 g (25%); yellow oil; MS: 513.9 (M+H)⁺

1-Benzyl-4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid was prepared starting from 1-benzyl-4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid ethyl ester (2.11 g, 4.1 mmol) dissolved in THF:methanol (3:1 150 ml) and 10 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 1.11 g (56%); white solid; mp 201 °C; MS: 485.9 (M+H)⁺

Starting from 1-benzyl-4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid (1 g, 2.06 mmol) and following the procedure outlined in example 83, 430 mg of 1-benzyl-4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide was isolated as a HCl salt, an off white solid. Yield 39%; mp

WO 99/42436

PCT/US98/17633

- 151 -

90.4°C; MS: 500.9 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 2.18-2.30 (m, 2H), 2.73-2.81 (m, 4H), 3.36 (d, 2H), 4.28 (d, J = 4.5 Hz, 2H), 7.15-7.25 (m, 4H), 7.43-7.48 (m, 3H), 7.51-7.56 (m, 4H), 7.74 (d, J = 9 Hz, 2H), 9.53 (s, 1H), 10.47 (s, 1H).

5

Example 253

[4-(3-methyl-butoxy)-phenylsulfanyl]-acetic acid ethyl ester

- 10 To stirred solution of (4-Hydroxy-phenylsulfanyl)-acetic acid ethyl ester (2.12 g, 10 mmol), K₂CO₃ (anhydrous, 10 g) and 1-bromo-3-methyl butane (3 g, excess) was added in boiling acetone. The reaction mixture was refluxed for 24 hrs and cooled to room temperature. The reaction mixture was filtered and concentrated. The residue obtained was extracted with chloroform; washed well with water and concentrated.
- 15 The crude product obtained was taken to next step with out purification. Yield 2.7 g (94%); (M+H)⁺ 283.

Example 254

20 [4-(3-methyl-butoxy)-phenylsulfonyl]-acetic acid ethyl ester

- [4-(3-methyl-butoxy)-phenylsulfonyl]-acetic acid ethyl ester was prepared according to the general method outlined in example 83 starting from [4-(3-methyl-butoxy)-phenylsulfonyl]-acetic acid ethyl ester (2.8 g, 10 mmol) and oxone (10 g). Yield 3.0g (99%);
- 25 oil; MS: 314EI (M+H)⁺

Example 255

30 1-Benzyl-4-[4-(3-methyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide

- 1-Benzyl-4-[4-(3-methyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting from [4-(3-methyl-butoxy)-phenylsulfonyl]-acetic acid ethyl ester (6.2 g, 20 mmol) and
- 35 bis-(2-chloro-ethyl)-benzyl amine (6.44 g, 24 mmol). Yield 8 g (84%); yellow oil; MS: 474 (M+H)⁺

1-Benzyl-4-[4-(3-methyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid was prepared starting from 1-Benzyl-4-[4-(3-methyl-butoxy)-benzenesulfonyl]-piperidine-

WO 99/42436

PCT/US98/17633

- 152 -

4-carboxylic ethyl ester (4.7 g, 10 mmol) dissolved in THF:methanol (3:1 150 ml) and 10 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 3 g (67%); white solid; mp 182 °C; MS: 446 (M+H)⁺

- 5 Starting from 1-Benzyl-4-[4-(3-methyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid (2.2 g, 5 mmol) and following the procedure outlined in example 83, 1.82 g of 1-Benzyl-4-[4-(3-methyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide was isolated as a HCl salt, an off white solid. Yield 73%; mp 106 °C; MS: 498 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.8 (d, 6H), 1.5 (m, 1H),
10 1.6-2.0 (m, 6H), 2.73-2.81 (m, 4H), 3.5 (d, 2H), 4.28 (d, J= 4.5 Hz, 2H), 7.15-7.25 (m, 4H), 7.43-7.48 (m, 3H), 7.51-7.56 (m, 4H), 7.74 (d, J = 9 Hz, 2H), 9.53 (s, 1H), 10.47 (s, 1H).

Example 256

- 15 1-Butyl-4-[4-(3-methyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide

- 1-Butyl-4-[4-(3-methyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid ethyl
20 ester was prepared according to the general method outlined in example 83 starting from [4-(3-methyl-butoxy)-phenylsulfonyl]-acetic acid ethyl ester (6.2 g, 20 mmol) and butyl-bis-(2-chloro-ethyl)-amine (5.2 g, 22 mmol). Yield 7 g (79%); yellow oil; MS: 440 (M+H)⁺

- 25 1-Butyl-4-[4-(3-methyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid was prepared starting from 1-Butyl-4-[4-(3-methyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic ethyl ester (4.4 g, 10 mmol) dissolved in THF:methanol (3:1 150 ml) and 10 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 3.2 g (77%); white solid; mp 188 °C; MS: 412 (M+H)⁺

- 30 Starting from 1-Butyl-4-[4-(3-methyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid (2.0 g, 5 mmol) and following the procedure outlined in example 83, 1.6 g of 1-Butyl-4-[4-(3-methyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide was isolated as a HCl salt, an off white solid. Yield 69%; mp 201 °C;
35 MS: 464 (M+H)⁺.

WO 99/42436

PCT/US98/17633

- 153 -

Example 257

[4-(2-Ethyl-butoxy)-phenylsulfanyl]-acetic acid ethyl ester

- 5 To stirred solution of (4-Hydroxy-phenylsulfanyl)-acetic acid ethyl ester (2.12 g, 10 mmol), K_2CO_3 (anhydrous, 10 g) and 1-bromo-2-ethyl butane (3 g, excess) was added in boiling acetone. The reaction mixture was refluxed for 24 hrs and cooled to room temperature. The reaction mixture was filtered and concentrated. The residue obtained was extracted with chloroform; washed well with water and concentrated. The crude
- 10 product obtained was taken to next step without purification. Yield 2.8 g (94%); $(M+H)^+$ 297.

Example 258

- 15 [4-(2-Ethyl-butoxy)-phenylsulfonyl]-acetic acid ethyl ester

- [4-(2-Ethyl-butoxy)-phenylsulfonyl]-acetic acid ethyl ester was prepared according to the general method outlined in example 83 starting from [4-(2-ethyl-butoxy)-phenyl-sulfanyl]-acetic acid ethyl ester (2.96 g, 10 mmol) and oxone (10 g). Yield 3.1g (
- 20 99%); oil; MS: 329EI $(M+H)^+$

Example 259

- 25 1-Benzyl-4-[4-(2-ethyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide

- 1-Benzyl-4-[4-(2-ethyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting from [4-(2-ethyl-butoxy)-phenylsulfonyl]-acetic acid ethyl ester (6.4 g, 20 mmol) and
- 30 bis-(2-chloro-ethyl)-benzyl amine (6.44 g, 24 mmol). Yield 8 g (82%); yellow oil; MS: 488 $(M+H)^+$

- 1-Benzyl-4-[4-(2-ethyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid was prepared starting from 1-Benzyl-4-[4-(2-ethyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic ethyl ester (4.8 g, 10 mmol) dissolved in THF:methanol (3:1 150 ml) and 10
- 35 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 4 g (87%); Semi solid; MS: 460 $(M+H)^+$

- Starting from 1-Benzyl-4-[4-(2-ethyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic
- 40 acid (2.2 g, 5 mmol) and following the procedure outlined in example 83, 1.02 g of 1-

WO 99/42436

PCT/US98/17633

- 154 -

Benzyl-4-[4-(2-ethyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide was isolated as a HCl salt, an off white solid. Yield 40%; mp 114 °C; MS: 512 (M+H)⁺.

5

Example 260

4-(4-butoxy-benzenesulfonyl)-1-(3-methoxy-benzyl)-piperidine-4-carboxylic acid hydroxyamide

10 4-(4-Butoxy-benzenesulfonyl)-1-(3-methoxy-benzyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting from 4-(4-butoxy-benzenesulfonyl)-acetic acid ethyl ester (20 g, 77.5 mmol) and bis-(2-chloro-ethyl)-(3-methoxy-benzyl)-amine (34 g, 116 mmol). Yield 9.53 g (25%); brown oil; MS: 490.2 (M+H)⁺

15

4-(4-Butoxy-benzenesulfonyl)-1-(3-methoxy-benzyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-methoxy-benzenesulfonyl)-1-(3-methoxy-benzyl)-piperidine-4-carboxylic acid ethyl ester (2.61 g, 5.34 mmol) dissolved in THF:methanol (3:1 150 ml) and 10 N NaOH (15 ml). The resulting reaction mixture
20 was worked up as outlined in example 83. Yield 1 g (41%); brown solid; mp 175°C; MS: 462.0 (M+H)⁺

Starting from 4-(4-butoxy-benzenesulfonyl)-1-(3-methoxy-benzyl)-piperidine-4-carboxylic acid (900 mg, 1.95 mmol) and following the procedure outlined in example
25 83, 200 mg of 4-(4-butoxy-benzenesulfonyl)-1-(3-methoxy-benzyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as a HCl salt, a brown powder. Yield 20%; mp 137 °C; MS: 477.0 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.96 (t, J=7.11 Hz, 3H), 1.48 (m, 2H), 1.73 (m, 2H), 2.27 (m, 2H), 2.47 (m, 2H), 2.78 (m, 2H), 3.35 (m, 2H), 3.77 (s, 2H), 4.08 (t, J= 6.3 Hz, 3H), 4.32 (s, 2H), 7.03 (t, 2H), 7.15
30 (m, 3H), 7.36 (t, J= 7.8 Hz, 1H), 7.64 (d, J = 9 Hz, 2H), 9.36 (s, 1H), 10.22 (s, 1H).

WO 99/42436

PCT/US98/17633

- 155 -

Example 261

4-(4-Methoxy-benzenesulfonyl)-1-(4-thiophen-2-yl-benzyl)-piperidine-4-carboxylic acid hydroxyamide

5 4-(4-Methoxy-benzenesulfonyl)-1-(4-thiophen-2-yl-benzyl)-piperidine-4-carboxylic acid ethyl ester was prepared starting from 1-(4-Bromo-benzyl)-4-(4-methoxy-benzensulfonyl)-piperidine-4-carboxylic acid ethyl ester (3 g, 6.05 mmol) and 2-(tributylstannyl)-thiophene (6.8 g, 18.14 mmol) in the presence of tetrakis palladium
10 (0) in boiling toluene. Yield 1.58 g (52%); brown solid; mp 130°C; MS: 500 (M+H)⁺

4-(4-Methoxy-benzenesulfonyl)-1-(4-thiophen-2-yl-benzyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-methoxy-benzenesulfonyl)-1-(4-thiophen-2-yl-benzyl)-piperidine-4-carboxylic acid ethyl ester (1.3 g, 2.61 mmol) dissolved in
15 THF:methanol (3:1 150 ml) and 10 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 950 mg (77%); brown solid; mp 235°C; MS: 471.8 (M+H)⁺

Starting from 4-(4-methoxy-benzenesulfonyl)-1-(4-thiophen-2-yl-benzyl)-piperidine-4-carboxylic acid (920 mg, 1.95 mmol) and following the procedure outlined in example
20 83, 510 mg of 4-(4-methoxy-benzenesulfonyl)-1-(4-thiophen-2-yl-benzyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as a HCl salt, a brown solid. Yield 50%; mp 166 °C; MS: 486.9 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 2.12-2.21 (m, 2H), 2.50 (m, 2H), 2.78 (m, 2H), 3.39 (m, 2H), 3.87 (s, 3H), 4.29 (d, 2H), 7.17
25 (m, 3H), 7.54-7.75 (m, 8H), 9.36 (s, 1H), 10.07 (s, 1H)

Example 262

4-(4-methoxy-benzenesulfonyl)-1-(4-pyridin-2-yl-benzyl)-piperidine-4-carboxylic acid
30 hydroxyamide

4-(4-Methoxy-benzenesulfonyl)-1-(4-pyridin-2-yl-benzyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 261. Starting from 1-(4-Bromo-benzyl)-4-(4-methoxy-benzensulfonyl)-piperidine-4-carboxylic acid ethyl ester (4.65 g, 9.38 mmol) and 2-(tributylstannyl)-pyridine (12.08
35 g, 32.8 mmol). Yield 2.79 g (60%); brown oil; MS: 495.1 (M+H)⁺

WO 99/42436

PCT/US98/17633

- 156 -

4-(4-Methoxy-benzenesulfonyl)-1-(4-pyridin-2-yl-benzyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-methoxy-benzenesulfonyl)-1-(4-pyridin-2-yl-benzyl)-piperidine-4-carboxylic acid ethyl ester (1.83 g, 3.7 mmol) dissolved in THF:methanol (3:1 150 ml) and 10 N NaOH (10 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 1.38 g (80%); off white solid; mp 217°C; MS: 466.9 (M+H)⁺

Starting from 4-(4-methoxy-benzenesulfonyl)-1-(4-pyridin-2-yl-benzyl)-piperidine-4-carboxylic acid (1.32 g, 2.83 mmol) and following the procedure outlined in example 83, 480 mg of 4-(4-methoxy-benzenesulfonyl)-1-(4-pyridin-2-yl-benzyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as a HCl salt, a white powder. Yield 33%; mp 214 °C; MS: 482.0 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 2.30 (m, 2H), 2.80 (m, 2H), 3.42 (d, J= 12.5 Hz, 2H), 3.75 (m, 2H), 3.88 (s, 3H), 4.36 (s, 2H), 7.15 (d, J= 8.9 Hz, 2H), 7.59-7.74 (m, 4H), 7.84-7.95 (m, 3H), 8.55 (d, J= 8.1 Hz, 1H), 8.79 (s, 1H), 9.14 (s, 1H), 10.68 (s, 1H), 11.17 (s, 1H)

Example 263

1-(3,4-Dichlorobenzyl)-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide

4-(4-Butoxy-benzenesulfonyl)-1-(3,4-dichloro-benzyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83. Starting from (4-butoxy-benzenesulfonyl)acetic acid ethyl ester (13.2g 44 mmol) (3,4-dichloro-benzyl)-bis-(2-chloro-ethyl)-amine (14.3g, mmol). Yield 14.1g (60%), white solid, MP 86°C; MS: 527.9 (M+H)⁺

1-(3,4-dichlorobenzyl)-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-Butoxy-benzenesulfonyl)-1-(3,4-dichloro-benzyl)-piperidine-4-carboxylic acid ethyl ester (14.0 g, 26.5 mmol) dissolved in THF:Methanol (100: 50 ml) and 10 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 7.87 (60%); off white solid; mp 239°C; MS: 501.9 (M+H)⁺

Starting 1-(3,4-dichlorobenzyl)-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid (7.7g, 15.5 mmol) and following the procedure outlined in example 83, 4.05g of 1-(3,4-dichlorobenzyl)-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as a HCl salt, white solid. Yield 48%; mp 256.8 °C; MS:

WO 99/42436

PCT/US98/17633

- 157 -

514.9 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.94 (t, 3H), 1.38-1.48 (q, 2H), 1.68-1.75 (q, 2H), 2.27 (m, 4H), 2.72 (m, 2H), 4.10 (t, 2H), 4.24 (s, 2H), 7.12-7.15 (d, J=8.9, 2H), 7.51-7.53 (d, J=8.1, 1H), 7.63-7.65 (d, J=8.8, 2H), 7.72-7.75 (d, J=9.9, 2H), 7.87 (s, 1H), 9.36 (s, 1H), 10.5 (s, 1H), 11.2 (s, 1H)

5

Example 264

[4-(4-Chloro-benzyloxy)-benzenesulfonyl]-1-methylpiperidine-4-carboxylic acid hydroxamide

10

[4-(4-Chloro-benzyloxy)-benzenesulfonyl]-1-methylpiperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting from [4-(4-chloro-benzyloxy)-benzenesulfonyl]-acetic acid ethyl ester (13.79g, 37 mmol) and 4-(4-chloro-benzyloxy)-bis-(2-chloro-ethyl)-amine (8.7g, 45 mmol). Yield 10.9 (65%); brown oil; MS: 451.9 (M+H)⁺

15

[4-(4-Chloro-benzyloxy)-benzenesulfonyl]-1-methylpiperidine-4-carboxylic acid was prepared starting from [4-(4-Chloro-benzyloxy)-benzenesulfonyl]-1-methylpiperidine-4-carboxylic acid ethyl ester (10.7 g, 24 mmol) dissolved in THF:Methanol (75: 75 ml) and 10 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 4.9g (50%); off white solid; MS: 426.2 (M+H)⁺

20

Starting from [4-(4-Chloro-benzyloxy)-benzenesulfonyl]-1-methylpiperidine-4-carboxylic acid (4.9g, 12 mmol) and following the procedure outlined in example 83, 1.2g of [4-(4-Chloro-benzyloxy)-benzenesulfonyl]-1-methylpiperidine-4-carboxylic acid hydroxamide was isolated as a HCl salt, off white solid. Yield 24%; mp 117.8 °C; MS: 438.9 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 2.2 (m, 2H), 2.49 (m, 4H), 2.5 (s, 3 H), 2.6 (m, 2H), 5.2 (s, 2H), 7.25-7.23 (d, t=8.7, 2H), 7.5 (d, t=2.7, 4H), 7.68-7.71 (d, t=9.6, 2H), 9.33 (s, 1H), 10.11 (s, 1H)

25

30

Example 265

4-(4-Butoxy-benzenesulfonyl)-1-(3-phenoxy-benzyl)-piperidine-4-carboxylic acid hydroxamide

35

4-(4-Butoxy-benzenesulfonyl)-1-(3-phenoxy-benzyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting from 4-(4-butoxy-benzenesulfonyl)-acetic acid ethyl ester (10.1g, 34 mmol) and 1-(3-

WO 99/42436

PCT/US98/17633

- 158 -

phenoxy-benzyl)-bis-(2-chloro-ethyl)-amine (18g, 50 mmol). Yield 8.9 (49%); brown oil; MS: 552.1 (M+H)⁺

5 4-(4-Butoxy-benzenesulfonyl)-1-(3-phenoxy-benzyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-butoxy-benzenesulfonyl)-1-(3-phenoxy-benzyl)-piperidine-4-carboxylic ethyl ester (10.7 g, 24 mmol) dissolved in THF:Methanol (75: 50 ml) and 10 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 5.0g (76%); off white solid; MS: 524.3 (M+H)⁺

10 Starting from 4-(4-Butoxy-benzenesulfonyl)-1-(3-phenoxy-benzyl)-piperidine-4-carboxylic acid (5.9g, 11 mmol) and following the procedure outlined in example 83, 0.39g of 4-(4-butoxy-benzenesulfonyl)-1-(3-phenoxy-benzyl)-piperidine-4-carboxylic acid hydroxamide was isolated as a HCl salt, tan solid. Yield 11%; mp 92.5 °C; MS: 539.1 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.93-0.97 (t, J=3.6, 3H), 1.49 (m, 2H), 1.73 (m, 2H), 2.51 (m, 4H), 4.09 (t, 2H), 4.29 (bs, 2H), 7.06-7.10 (d, J=12, 2H), 7.13-7.15 (m, 3H), 7.39-7.42 (d, 2H), 7.63-7.66 (d, 2H), 9.50 (s, 1H), 9.98 (s, 1H)

Example 266

20 [4-(4-Chloro-benzyloxy)-benzenesulfonyl]- 1-(4-methylbenzyl)-piperidine-4-carboxylic acid hydroxamide

[4-(4-Chloro-benzyloxy)-benzenesulfonyl]- 1-(4-methylbenzyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting from [1-(4-chloro-benzyloxy)-benzenesulfonyl]-acetic acid ethyl ester (5.47g, 15 mmol) and 1-(4-methyl-benzyl)-bis-(2-chloro-ethyl)-amine (5.23g, 18 mmol). Yield 8.0 (96%); brown oil; MS: 542.0 (M+H)

30 [4-(4-Chloro-benzyloxy)-benzenesulfonyl]- 1-(4-methylbenzyl)-piperidine-4-carboxylic acid was prepared starting from [4-(4-Chloro-benzyloxy)-benzenesulfonyl]-1-(4-methylbenzyl)-piperidine-4-carboxylic acid ethyl ester (7.9 g, 124 mmol) dissolved in THF:Methanol (50: 50 ml) and 10 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield 4.6g (61%); off white solid, mp 204 °C; MS: 514.2 (M+H)⁺

Starting from [4-(4-Chloro-benzyloxy)-benzenesulfonyl]- 1-(4-methylbenzyl)-piperidine-4-carboxylic acid (4.2g, 8 mmol) and following the procedure outlined in

WO 99/42436

PCT/US98/17633

- 159 -

example 83, 1.3g of [4-(4-Chloro-benzyloxy)-benzenesulfonyl]-1-(4-methylbenzyl)-piperidine-4-carboxylic acid hydroxamide was isolated as a HCl salt, yellow solid. Yield 29%; mp 172 °C; MS: 528.9 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ

5

Example 267

4-(4-Butoxy-benzenesulfonyl)-1-(4-methylbenzyl)-piperidine-4-carboxylic acid hydroxamide

10 4-(4-Butoxy-benzenesulfonyl)-1-(4-methylbenzyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting 4-(4-Butoxy-benzenesulfonyl)-acetic acid ethyl ester (5.47g, 15 mmol) and 1-(4-methylbenzyl)-bis-(2-chloro-ethyl)-amine (15.3g, 51 mmol). Yield 10.1 (57%); white solid, MP 93°C; MS: 474.1 (M+H)⁺

15

4-(4-Butoxy-benzenesulfonyl)-1-(4-methylbenzyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-Butoxy-benzenesulfonyl)-1-(4-methylbenzyl)-piperidine-4-carboxylic ethyl ester (10.0 g, 22 mmol) dissolved in THF:Methanol (50:50 ml) and 10 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in

20 example 83. Yield 7.2g (72%); off white solid, mp 244 °C; MS: 446.3 (M+H)⁺

Starting from 4-(4-Butoxy-benzenesulfonyl)-1-(4-methylbenzyl)-piperidine-4-carboxylic acid (6.6g, 1.5 mmol) and following the procedure outlined in example 83, 2.06g of 4-(4-Butoxy-benzenesulfonyl)-1-(4-methylbenzyl)-piperidine-4-carboxylic acid hydroxamide was isolated as a HCl salt, yellow solid. Yield 28%; mp 137 °C; MS: 461.3 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ 0.91-0.964 (t, J=7.3, 3H), 1.41 (m, 2H), 1.70 (m, 4H), 1.79 (t, s, 3H), 2.52 (m, 2H), 2.76 (m, 2H), 3.33 (bd, 2H), 4.10 (t, 2H), 4.22 (d, 2H), 7.12-7.14 (d, J=8.7, 2H), 7.25-7.28 (d, J=8.1, 2H), 7.42-7.45 (d, J=7.8, 2H), 7.63-7.65 (d, J=8.7, 2H), 10.31 (s, 1H), 10.75 (bs, 1H)

30

Example 268

4-(4-Butoxy-benzenesulfonyl)-1-(4-cyano-benzyl)-piperidine-4- hydroxamide
35 carboxylic acid hydroxamide

4-(4-Butoxy-benzenesulfonyl)-1-(4-cyano-benzyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting 4-

WO 99/42436

PCT/US98/17633

- 160 -

(4-Butoxy-benzenesulfonyl)-acetic acid ethyl ester (5.29g, 17.6 mmol) and 1-(4-cyano-benzyl)-bis-(2-chloro-ethyl)-amine (6.19g, 21 mmol). Yield 6.8g (80%); tan oil; MS: 485.0 (M+H)⁺

- 5 4-(4-Butoxy-benzenesulfonyl)-1-(4-cyanobenzyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-Butoxy-benzenesulfonyl)-1-(4-cyano-benzyl)-piperidine-4-carboxylic ethyl ester (10.0 g, 124 mmol) dissolved in THF:Methanol (75: 50 ml) and 10 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in example 83. Yield .7g (11%); off white solid; MS: 456.0 (M+H)⁺

10

Starting from 4-(4-Butoxy-benzenesulfonyl)-1-(4-cyanobenzyl)-piperidine-4-carboxylic acid (.600g, 1.2 mmol) and following the procedure outlined in example 83, .21g of 4-(4-Butoxy-benzenesulfonyl)-1-(4-cyano-benzyl)-piperidine-4-carboxylic acid hydroxamide was isolated as a HCl salt, off white solid. Yield 34%; mp 241.6 °C;

- 15 MS: 472.0 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆): δ .915-.964 (t, J=7.2, 3H), 1.51 (q, 2H), 1.75 (q, 2H), 2.27 (m, 2H), 2.49 (m, 4H), 4.11-4.19 (t, 2H), 4.37 (s, 1H), 7.12-7.15 (d, J=8.7, 2H), 7.63-7.66 (d, J=9, 2H), 7.72-7.74 (d, J=7.8, 2H), 9.36 (s, 1H), 10.23 (s, 1H), 11.16 (s, 1H)

20

Example 269

4-(4-Butoxy-benzenesulfonyl)-1-pyridin-4-ylmethyl-piperidine-4-carboxylic acid hydroxamide

- 25 4-(4-Butoxy-benzenesulfonyl)-1-pyridin-4-ylmethyl-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method outlined in example 83 starting with 4-(4-Butoxy-benzenesulfonyl)-acetic acid ethyl ester (6.0g, 20.0 mmol) and pyridin-4-ylmethyl-bis-(2-chloro-ethyl)-amine. (4.89g, 21 mmol). Yield 4.5g (48%); Brown oil; MS: 461.0 (M+H)⁺

30

4-(4-Butoxy-benzenesulfonyl)-1-pyridin-4-ylmethyl-piperidine-4-carboxylic acid was prepared starting from 4-(4-Butoxy-benzenesulfonyl)-1-pyridin-4-ylmethyl-piperidine-4-carboxylic acid ethyl ester (3.0 g, 6.5 mmol) dissolved in THF:Methanol (75: 50 ml) and 10 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in

- 35 example 83. Yield 1.2g (42%); off white solid; MS: 433.0 (M+H)⁺

WO 99/42436

PCT/US98/17633

- 161 -

Starting from 4-(4-Butoxy-benzenesulfonyl)-1-pyridin-4-ylmethyl-piperidine-4-carboxylic acid (0.864 mg, 2.0 mmol) and following the procedure as outlined in example 83, 600 mg of 4-(4-Butoxy-benzenesulfonyl)-1-pyridin-4-ylmethyl-piperidine-4-carboxylic acid hydroxyamide was isolated as a HCl salt, off white solid.

- 5 Yield 67%; mp 118 °C; MS: 447.9 (M+H)⁺; ¹H NMR (300MHz, DMSO-d₆): δ 0.94 (t, 3H), 1.11 (t, 1H), 1.23 (t, 1H), 1.44 (m, 1H), 1.73 (m, 1H), 2.34 (m, 2H), 2.78 (m, 2H), 3.10 (m, 2H), 3.38 (m, 2H), 4.08 (t, 2H), 4.42 (br s, 2H), 7.13 (d, 2H), 7.64 (d, 2H), 7.94 (d, 2H), 8.82(d, 2H), 11.2 (br s, 1H), 11.4 (br s, 1H).

10 References:

- 1 Rickter, L. S.; Desai, M. C. *Tetrahedron Letters*, 1997, 38, 321-322.

The subject compounds of the present invention were tested for biological activity according to the following procedures.

15

In Vitro Gelatinase Assay

- The assay is based on the cleavage of the thiopeptide substrate ((Ac-Pro-Leu-Gly(2 mercapto-4 methyl-pentanoyl)-Leu-Gly-OEt), Bachem Bioscience) by the enzyme, 20 gelatinase, releasing the substrate product which reacts colorimetrically with DTNB ((5,5'-dithio-bis(2-nitro-benzoic acid)). The enzyme activity is measured by the rate of the color increase.

- The thiopeptide substrate is made up fresh as a 20 mM stock in 100% DMSO and the DTNB is dissolved in 100% DMSO as a 100 mM stock and stored in dark at room 25 temperature. Both the substrate and DTNB are diluted together to 1 mM with substrate buffer (50 mM HEPES pH 7.5, 5 mM CaCl₂) before use. The stock of human neutrophil gelatinase B is diluted with assay buffer (50 mM HEPES pH 7.5, 5 mM CaCl₂, 0.02% Brij) to a final concentration of 0.15 nM.

- The assay buffer, enzyme, DTNB/substrate (500 μM final concentration) and vehicle or 30 inhibitor are added to a 96 well plate (total reaction volume of 200 μl) and the increase in color is monitored spectrophotometrically for 5 minutes at 405 nm on a plate reader. The increase in OD₄₀₅ is plotted and the slope of the line is calculated which represents the reaction rate.

- The linearity of the reaction rate is confirmed ($r^2 > 0.85$). The mean ($x \pm \text{sem}$) of the 35 control rate is calculated and compared for statistical significance ($p < 0.05$) with drug-treated rates using Dunnett's multiple comparison test. Dose-response relationships can

WO 99/42436

PCT/US98/17633

- 162 -

be generated using multiple doses of drug and IC₅₀ values with 95% CI are estimated using linear regression (IPRED, HTB).

References: Weingarten, H and Feder, J., Spectrophotometric assay for vertebrate collagenase, Anal. Biochem. 147, 437-440 (1985).

5

In Vitro Collagenase Assay

The assay is based on the cleavage of a peptide substrate ((Dnp-Pro-Cha-Gly-Cys(Me)-His-Ala-Lys(NMa)-NH₂), Peptide International, Inc.) by collagenase releasing the
10 fluorescent NMa group which is quantitated on the fluorometer. Dnp quenches the NMa fluorescence in the intact substrate. The assay is run in HCBC assay buffer (50 mM HEPES, pH 7.0, 5 mM Ca²⁺, 0.02% Brij, 0.5% Cysteine), with human recombinant fibroblast collagenase (truncated, mw=18,828, WAR, Radnor). Substrate
15 is dissolved in methanol and stored frozen in 1 mM aliquots. Collagenase is stored frozen in buffer in 25 µM aliquots. For the assay, substrate is dissolved in HCBC buffer to a final concentration of 10 µM and collagenase to a final concentration of 5 nM. Compounds are dissolved in methanol, DMSO, or HCBC. The methanol and DMSO are diluted in HCBC to < 1.0%. Compounds are added to the 96 well plate containing enzyme and the reaction is started by the addition of substrate.

20

The reaction is read (excitation 340 nm, emission 444 nm) for 10 min. and the increase in fluorescence over time is plotted as a linear line. The slope of the line is calculated and represents the reaction rate.

25 The linearity of the reaction rate is confirmed ($r^2 > 0.85$). The mean ($x \pm \text{sem}$) of the control rate is calculated and compared for statistical significance ($p < 0.05$) with drug-treated rates using Dunnett's multiple comparison test. Dose-response relationships can be generated using multiple doses of drug and IC₅₀ values with 95% CI are estimated using linear regression (IPRED, HTB).

30

References: Bickett, D. M. et al., A high throughput fluorogenic substrate for interstitial collagenase (MMP-1) and gelatinase (MMP-9), Anal. Biochem. 212,58-64 (1993).

WO 99/42436

PCT/US98/17633

- 163 -

Procedure for Measuring TACE Inhibition

- Using 96-well black microtiter plates, each well receives a solution composed of 10 μL
- 5 TACE (Immunex, final concentration 1 $\mu\text{g/mL}$), 70 μL Tris buffer, pH 7.4 containing 10% glycerol (final concentration 10 mM), and 10 μL of test compound solution in DMSO (final concentration 1 μM , DMSO concentration <1%) and incubated for 10 minutes at room temperature. The reaction is initiated by addition of a fluorescent peptidyl substrate (final concentration 100 μM) to each well and then shaking on a
- 10 shaker for 5 sec.
- The reaction is read (excitation 340 nm, emission 420 nm) for 10 min. and the increase in fluorescence over time is plotted as a linear line. The slope of the line is calculated and represents the reaction rate.
- The linearity of the reaction rate is confirmed ($r^2 > 0.85$). The mean ($\bar{x} \pm \text{sem}$) of the
- 15 control rate is calculated and compared for statistical significance ($p < 0.05$) with drug-treated rates using Dunnett's multiple comparison test. Dose-response relationships can be generated using multiple doses of drug and IC_{50} values with 95% CI are estimated using linear regression
- 20 The results obtained following these standard experimental test procedures are presented in the following table.

WO 99/42436

PCT/US98/17633

- 164 -

Example	IC 50 (nM or % inhibition at 1 micromolar or 10micromolar (*))			
	MMP 1	MMP 9	MMP 13	TACE
1	NT	559.6	193.3	31.62%
2	NT	10.50%	0%	403
3	NT	308.9	169.4	27.43%
4	371	22.20%	17.10%	21%
5	NT	7.7	4.7	25%
6	267	21.4	15.6	40.43%
7	844	72.9	42.1	33%
8	NT	346	307.9	47%
9	313	107	NT	20.30%
10	8%	128	64	54.75%
11	18.80%	2925	319	942
12	100	10.8	11	15.50%
13	239	11	14	626
14	158	23	8	17.18%
15	285	17	4	137
16	325	9	24	180
17	238.6	8.9	1.4	41.00%
18	540	18.9	11.5	29.2%
19	446	95.8	4.8	33.1%
20	423	14.6	18.7	31%
21	318	13.2	15.3	39%
22	219	3.2	2.5	30%
23	593	7.9	4.0	40.6%
24	413	20.9	31.3	47.5
25	262	26.7	8.0	NT
26	304.6	6.3	3.2	34.6
27	629	106	30.1	NT
28	761	3.1	2.0	30.6%
29	297	4.3	3.6	41%
30	397	8.1	5.7	25.2%
31	162	15.2	5.7	688
32	13.7	3.7	1.0	NT
33	318	53.9	18.4	23.9%
34	519.8	34.7	26.1	28.1%
35	455.8	233.6	48.2	44.9
36	622	83.8	20.7	826
37	9%	31.6%	14.3%	87
38	48.3%	1.7%	5.8%	55.1%
39	29.4%	35.2%	26.6%	69.4
40	583	197	14	160
41	100	10.8	11	15.50%
42	262	50.9	6.2	36.5
43	66.1%	34.7%	55.5%	46.6%
44	47.1%	36.9%	39.5%	14.9%

WO 99/42436

PCT/US98/17633

- 165 -

Example	MMP 1	MMP 9	MMP 13	TACE
45	49%	48.6%	36.7%	20.4%
46	78.9%	79.12%	84.7%	1.4%
47	17.1%	12.9%	7.12%	3.3%
48	99.1%	79.1%	85.4%	51.1%
49	10.1%	23.7%	54.6%	NT
50	51.1	58.4	10.6	NT
51	178.1	10.4	13.1	48.14%
52	139.3	7.9	9.1	NT
53	647.9	27.80%	188	52.57%
54	110	66	21	55.10%
55	303	10	7	21.70%
56	299	16	12	65%
57	258	332	191	16.57%
58	211	35	39	7.70%
59	30.20%	447	141	24.86%
60	NT	184	NT	23.60%
61	258	38	22	17.21%
62	522	174	43	669
63	156	9	3	203
64	40.90%	25.60%	36.70%	29.70%
65	1000	63	13	42.21%
66	1600	131	226	42.33%
67	364	2.3	43.7	690
68	297	29	27	522
69	574.5	120.2	90	41.32%
70	1139	88.80%	127	764
71	1000	63	13	42.21%
72	117	11	1	51.64%
73	300	141	12	20.17%
74	138.1	9.2	4.3	47.86%
75	672.3	83.4	32.7	23.77%
76	805	NT	500	NT
77	205.5	NT	170	NT
78	262	560	34	24.58%
79	25	0.54	0.4	805
80	22.1%	26%	63.6%	191
81a	2036	230.9	43.9	27.1
81b	3765	154	15.7	228
82	237.6	19.4	5.1	34.5%
83	492	10.2	2.0	229
84	519	8.8	2.0	213
85	450	5.8	1.5	115
86	494	16.8	1.5	222
87	368	5.0	1.6	170.7
88	1329	12.8	3.1	610

WO 99/42436

PCT/US98/17633

- 166 -

Example	MMP 1	MMP 9	MMP 13	TACE
89	1389	38.6	7.0	49%
90	598	10.3	2.2	71.9
91	1929	13.3	10.8	503
92	59.6%	649	148	9.7
93	56.3%	452	38	15.8%
94	2640	138	28.6	22.9
95	3681	364	33.1	25.4%
96	4437	374	33.8	18.1
97	5109	484	43.7	20.20%
98	2383	3.8	1.2	154
99	656	16.2	2.4	250
100	4729	19.1	5.3	39.5%
101	642	12.3	2.1	197
102	662	33.7	1.9	53%
103	1306	45.1	8.8	470
104	2610	3.1	1.4	208
105	1214	44.2	4.1	50.2%
106	3788	5.1	0.9	631
107	629	26.8	2.5	293
108	2896	5.4	1.7	270
109	393	2.7	2.5	386
241	48.2% *	2.7	15.8	277
242	1950	2	1.3	581
243	2181	1.9	1.5	506
244	3417	9.8	1.5	594
245	7062	43.4	2.2	51.95%*
246	50.30%*	28.3	2.4	880
249	1412	2	1.6	270
250	1717	1.6	0.8	413
251	1067	0.8	0.9	301
252	801	1.1	0.9	278
255	2558	3.6	1.5	565
256	10000	7.2	2.9	43.01%*
259	3160	14.3	5.3	39%*
260	1495	2.9	1.3	272
261	513	10.9	2.7	273
262	422	6.1	2.3	298
263	3669	20.3	5.2	57.70%*
264	4293	2.9	3.1	182
265	1944	9.3	7.8	1037
266	4746	6	5.7	421
267	3620	5.4	2.3	508
268	2292	2.8	1.1	278
269	2071	2.2	1.4	296

WO 99/42436

PCT/US98/17633

- 167 -

Compounds prepared by solid phase synthesis Data: for Examples 110 to 240

Example No	MMP 1	MMP 9	MMP 13 % inhibition at 0.2 μ M (HTS)	MMP 13 % inhibition at 0.2 μ M (manual)	TACE % inhibition at 1 mM
110			75		17.6
111			10		40.4
112			50		33.7
113			0		13.1
114			0		0
115			0		0
116			0		9.1
117			7		8.1
118			24		16.7
119			0		7.8
120			31		19.9
121			0		6.1
122			0		3.1
123			0		2.5
124			0		0
125			5		2.3
126			25		10.4
127			47		29.2
128	1.9 mM	213 nM	91	255 nM	19.31
129			90		32.77
130			28		27.9
131			71		20.73
132			71		20.76
133			53		22.04
134			25		-9.31
135			79		42.67
136			89		42.69
137			83		13.35
138			20		5.284
139			8		28.05
140			29		-4.22
141			32		11.76
142			69		54.27
143			53		43.9
144			38		19.7
145			45		2.5
146			68		7.317
147			73		11.95
148			15		43.46
149			13		4.408
150			54		1.818
151			6		5.927
152			9		10.03
153			12		11.8
154			89		13.14
155			31		18.62
156			23		-2.09

WO 99/42436

PCT/US98/17633

- 168 -

Example No	MMP 1	MMP 9	MMP 13 % inhibition at 0.2 μ M (HTS)	MMP 13 % inhibition at 0.2 μ M (manual)	TACE % inhibition at 1 mM
157			19		13.7
158			33		-7.48
159			49		5.852
160			14		-3.57
161			0		12.7
162			13		0
163			84		9.515
164			74		62.69
165			71		73.7
166			9		4.16
167			27		8.961
168			21		3.688

Example No.	MMP 13 % inhibition at 36 nM (HTS)	MMP 13 % inhibition at 0.36 mM (HTS)	MMP 13 % inhibition at 3.6 mM (HTS)	TACE IC ₅₀ nM	TACE % inhibition at 1 mM
169	28	40	72		41.7
170	32	49	90		25.5
171	31	38	48		16.6
172	34	32	42		29.4
173	18	46	56		25.5
174	10	19	40		27.7
175	16	20	37		32.9
176	6	5	16		26.6
177	5	1	9		38.5
178	-10	74	39		26
179	12	32	60		42.7
180	14	19	45		34.4
181	6	35	62		15.7
182	-9	-8	7		28.6
183	-6	12	70		34.6
184	16	24	44		24.8
185	9	0	23		7.21
186	-14	-4	35		19.5
187	-14	-12	20		85.5
188	-27	-24	4		16.2
189	-30	-18	-9		14.
190	-35	-28	-13		38.3
191	-45	-3	22		2.9
192	-32	5	61		33.2
193	-32	-15	56		14.9
194	-17	-8	5		5.4
195	-9	-2	10		27.0
196	-18	1	11		35.7
197	-33	-26	-3		17.8
198	-39	-7	15		17.1
199	-10	-7	30		-1.0

WO 99/42436

PCT/US98/17633

- 169 -

Example No.	MMP 13 % inhibition at 36 nM (HTS)	MMP 13 % inhibition at 0.36 mM (HTS)	MMP 13 % inhibition at 3.6 mM (HTS)	TACE IC ₅₀ nM	TACE % inhibition at 1 mM
200					37.9
201					50.9
202					10.6
203					32.8
204					7.75
205					84.0
206					89.8
207					-6.3
208					67.7
209					31.2
210					52.2
211					20.7
212					56.0
213					-17.5
214					11.03
215				895	60.12
216					2.49
217					55.1
218				380	68.7
219					7.3
220				256	53.1
221				146	98.9
222				212	89.3
223				226	107.3
224				404	75.0
225				96.6	114.3
226	28	22	28		2.2
227	15	-16	-22		7.3
228	37	28	65		6.8
229	29	17	33		34.4
230	29	31	26	700	72.1
231	23	13	5		41.6
232	30	17	42		20.8
233	33	29	46		19.8
234	26	28	40		18.4
235	59	70	70		48.3
236	44	44	64		35
237	55	65	72		38.2
238	22	11	24	930	54.4
239	54	74	83		45.9
240	48	51	46		40.3

WO 99/42436

PCT/US98/17633

- 170 -

Pharmaceutical Composition

Compounds of this invention may be administered neat or with a pharmaceutical carrier to a patient in need thereof. The pharmaceutical carrier may be solid or liquid.

5 Applicable solid carriers can include one or more substances which may also act as flavoring agents, lubricants, solubilizers, suspending agents, fillers, glidants, compression aids, binders or tablet-disintegrating agents or an encapsulating material. In powders, the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier having
10 the necessary compression properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain up to 99% of the active ingredient. Suitable solid carriers include, for example, calcium phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, polyvinylpyrrolidone, low melting waxes
15 and ion exchange resins.

 Liquid carriers may be used in preparing solutions, suspensions, emulsions, syrups and elixirs. The active ingredient of this invention can be dissolved or suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both or pharmaceutically acceptable oils or fat. The liquid carrier
20 can contain other suitable pharmaceutical additives such as solubilizers, emulsifiers, buffers, preservatives, sweeteners, flavoring agents, suspending agents, thickening agents, colors, viscosity regulators, stabilizers or osmo-regulators. Suitable examples of liquid carriers for oral and parenteral administration include water (particularly containing additives as above, e.g., cellulose derivatives, preferable sodium
25 carboxymethyl cellulose solution), alcohols (including monohydric alcohols and polyhydric alcohols, e.g., glycols) and their derivatives, and oils (e.g., fractionated coconut oil and arachis oil). For parenteral administration the carrier can also be an oily ester such as ethyl oleate and isopropyl myristate. Sterile liquid carriers are used in sterile liquid form compositions for parenteral administration.

30 Liquid pharmaceutical compositions which are sterile solutions or suspensions can be utilized by, for example, intramuscular, intraperitoneal or subcutaneous injection. Sterile solutions can also be administered intravenously. Oral administration may be either liquid or solid composition form.

 The compounds of this invention may be administered rectally in the form of a
35 conventional suppository. For administration by intranasal or intrabronchial inhalation or insufflation, the compounds of this invention may be formulated into an aqueous or partially aqueous solution, which can then be utilized in the form of an aerosol. The

WO 99/42436

PCT/US98/17633

- 171 -

compounds of this invention may also be administered transdermally through the use of a transdermal patch containing the active compound and a carrier that is inert to the active compound, is non-toxic to the skin, and allows delivery of the agent for systemic absorption into the blood stream via the skin. The carrier may take any number of
5 forms such as creams and ointments, pastes, gels, and occlusive devices. The creams and ointments may be viscous liquid or semi-solid emulsions of either the oil in water or water in oil type. Pastes comprised of absorptive powders dispersed in petroleum or hydrophilic petroleum containing the active ingredient may also be suitable. A variety of occlusive devices may be used to release the active ingredient into the blood stream
10 such as a semipermeable membrane covering a reservoir containing the active ingredient with or without a carrier, or a matrix containing the active ingredient. Other occlusive devices are known in the literature.

The dosage to be used in the treatment of a specific patient suffering from a disease or condition in which MMPs and TACE are involved must be subjectively
15 determined by the attending physician. The variables involved include the severity of the dysfunction, and the size, age, and response pattern of the patient. Treatment will generally be initiated with small dosages less than the optimum dose of the compound. Thereafter the dosage is increased until the optimum effect under the circumstances is reached. Precise dosages for oral, parenteral, nasal, or intrabronchial administration
20 will be determined by the administering physician based on experience with the individual subject treated and standard medical principles.

Preferably the pharmaceutical composition is in unit dosage form, e.g., as tablets or capsules. In such form, the composition is sub-divided in unit dose containing appropriate quantities of the active ingredient; the unit dosage form can be
25 packaged compositions, for example packed powders, vials, ampoules, prefilled syringes or sachets containing liquids. The unit dosage form can be, for example, a capsule or tablet itself, or it can be the appropriate number of any such compositions in package form.


WO 99/42436

PCT/US98/17633

- 172 -

What is claimed is:

1. A compound according to formula I



5 wherein:

R¹ is alkyl of 1 to 18 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

10 alkenyl of 3 to 18 carbon atoms having 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;
alkynyl of 3 to 18 carbon atoms having 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;
aryl of 6 to 10 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

15 cycloalkyl of 3 to 8 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;
saturated or unsaturated 5 to 10 membered mono or bicyclic heterocycle containing one heteroatom selected from O, S or NR⁷, optionally substituted with one or two groups selected independently from R⁵;
20 or heteroaryl-(CH₂)₀₋₆ wherein the heteroaryl group is 5 to 6 membered with one or two heteroatoms selected independently from O, S, and N and may be optionally substituted with one or two groups selected independently from R⁵;

A is -S-, -SO- or SO₂-;

25 R² and R³, taken with the carbon atom to which they are attached, form a 5 to 7 membered heterocyclic ring containing O, S or N-R⁷ optionally having one or two double bonds;

R⁴ is hydrogen,

30 alkyl of 1 to 6 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;
alkenyl of 3 to 18 carbon atoms having 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;

WO 99/42436

PCT/US98/17633

- 173 -

- alkynyl of 3 to 18 carbon atoms having 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵; phenyl or naphthyl optionally substituted with one or two groups selected independently from R⁵;
- 5 C₃ to C₈ cycloalkyl or bicycloalkyl optionally substituted with one or two groups selected independently from R⁵;
- saturated or unsaturated 5 to 10 membered mono or bicyclic heterocycle containing one heteroatom selected from O, S or NR⁷, optionally substituted with one or two groups selected independently from R⁵;
- 10 R⁵ is H, C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁ to C₁₂ alkyl, C₂ to C₁₂ alkenyl, C₂-C₁₂ alkynyl, F, Cl, Br, I, CN, CHO, C₁-C₆ alkoxy, aryloxy, heteroaryloxy, C₃-C₆ alkenyloxy, C₃-C₆ alkynyloxy, C₁-C₆ alkoxyaryl, C₁-C₆ alkoxyheteroaryl, C₁-C₆ alkylamino-C₁-C₆ alkoxy, C₁-C₂ alkylene dioxy, aryloxy-C₁-C₆ alkyl amine, C₁-C₁₂ perfluoro alkyl, S(O)_n-C₁-C₆ alkyl, S(O)_n-aryl where n is 0, 1
- 15 or 2; OCOO C₁-C₆ alkyl, OCOOaryl, OCONR⁶, COOH, COO C₁-C₆ alkyl, COOaryl, CONR⁶R⁶, CONHOH, NR⁶R⁶, SO₂NR⁶R⁶, NR⁶SO₂aryl, -NR⁶CONR⁶R⁶, NHSO₂CF₃, SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl, CONHSO₂aryl, SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl, CONHSO₂aryl, NH₂, OH, aryl, heteroaryl, C₃ to C₈ cycloalkyl; or saturated or
- 20 unsaturated 5 to 10 membered mono or bicyclic heterocycle containing one heteroatom selected from O, S or NR⁷, wherein C₁-C₆ alkyl is straight or branched, heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2
- 25 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;
- R⁶ is H, C₁ to C₁₈ alkyl optionally substituted with OH; C₃ to C₆ alkenyl, C₃ to C₆ alkynyl, C₁ to C₆ perfluoro alkyl, S(O)_n-C₁-C₆ alkyl S(O)_n aryl where n is 0, 1 or 2; or COheteroaryl, wherein heteroaryl is a 5-10 membered mono or
- 30 bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;

WO 99/42436

PCT/US98/17633

- 174 -

and R⁷ is C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁-C₁₂ perfluoro alkyl, S(O)_n-C₁-C₆-alkyl, S(O)_n-aryl where n is 0, 1 or 2; COO-C₁-C₆-alkyl, COOaryl, CONHR⁶, CONR⁶R⁶, CONHOH, SO₂NR⁶R⁶, SO₂CF₃, SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO-C₁-C₆-alkyl, CONHSO₂aryl, aryl, or heteroaryl,

5 where aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected independently from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy; and heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or N-C₁-C₆ alkyl;

10 alkyl of 1 to 18 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

 alkenyl of 3 to 18 carbon atoms having from 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;

 alkynyl of 3 to 18 carbon atoms having from 1 to 3 triple bonds, optionally

15 substituted with one or two groups selected independently from R⁵;

 arylalkyl of 7 to 16 carbon atoms, wherein aryl is optionally substituted with one or two groups selected independently from R⁵;

 heteroarylalkyl wherein alkyl is from 1 to 6 carbon atoms and heteroaryl contains 1 or 2 heteroatoms selected from O, S or N and is optionally

20 substituted with one or two groups selected independently from R⁵;

 biphenylalkyl of 13 to 18 carbon atoms, wherein biphenyl is optionally substituted with one or two groups selected independently from R⁵;

 arylalkenyl of 8 to 16 carbon atoms, wherein aryl is optionally substituted with one or two groups selected independently from R⁵;

25 cycloalkylalkyl or bicycloalkylalkyl of 4 to 12 carbon atoms, wherein the cycloalkyl or bicycloalkyl group is optionally substituted with one or two groups selected independently from R⁵;

 saturated or unsaturated mono or bicyclic heterocycle containing one heteroatom selected from O, S or N-C₁-C₆ alkyl, optionally substituted with one or

30 two groups selected independently from R⁵; or

 R⁸R⁹N-C₁-C₆-alkoxyaryl-C₁-C₆-alkyl where R⁸ and R⁹ are independently selected from C₁-C₆ alkyl or R⁸ and R⁹ together with the interposed nitrogen forms a 5-7 membered saturated heterocyclic ring optionally

WO 99/42436

PCT/US98/17633

- 175 -

containing an oxygen atom, wherein the aryl group is phenyl or naphthyl;
or a pharmaceutically acceptable salt thereof.

5 2. A compound according to claim 1 wherein:

R¹ is alkyl of 1 to 18 carbon atoms, optionally substituted with one or two groups
selected independently from R⁵;

alkenyl of 3 to 18 carbon atoms having 1 to 3 double bonds, optionally
substituted with one or two groups selected independently from R⁵;

10 alkynyl of 3 to 18 carbon atoms having 1 to 3 triple bonds, optionally
substituted with one or two groups selected independently from R⁵;

aryl of 6 to 10 carbon atoms, optionally substituted with one to two groups
selected independently from R⁵;

15 cycloalkyl of 3 to 8 carbon atoms, optionally substituted with one to two groups
selected independently from R⁵;

saturated or unsaturated mono or bicyclic heterocycle of from 5 to 10 members
containing one heteroatom selected from O, S or NR⁷, optionally
substituted with one to two groups selected independently from R⁵;

20 or heteroaryl-(CH₂)₀₋₆ wherein the heteroaryl group is 5 to 6 membered with
one or two heteroatoms selected independently from O, S, and N and
may be optionally substituted with one or two groups selected
independently from R⁵;

A is -S-, -SO- or SO₂-;

25 R² and R³, taken with the carbon atom to which they are attached, form a 5 to 7
membered heterocyclic ring containing O, S or N-R⁷ optionally having one or
two double bonds;

R⁴ is hydrogen,

alkyl of 1 to 6 carbon atoms, optionally substituted with one or two groups
selected independently from R⁵;

30 alkenyl of 3 to 18 carbon atoms having 1 to 3 double bonds, optionally
substituted with one or two groups selected independently from R⁵;

alkynyl of 3 to 18 carbon atoms having 1 to 3 triple bonds, optionally
substituted with one or two groups selected independently from R⁵;

WO 99/42436

PCT/US98/17633

- 176 -

phenyl or naphthyl optionally substituted with one or two groups selected independently from R⁵;

C₃ to C₈ cycloalkyl or bicycloalkyl optionally substituted with one or two groups selected independently from R⁵;

- 5 R⁵ is H, F, Cl, Br, I, CN, CHO, C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁ to C₁₂ alkyl, C₂ to C₁₂ alkenyl, C₂-C₁₂ alkynyl, C₁-C₆ alkoxy, aryloxy, heteroaryloxy, C₃-C₆ alkenyloxy, C₃-C₆ alkynyloxy, C₁-C₆ alkoxyaryl, C₁-C₆ alkoxyheteroaryl, C₁-C₆-alkylamino-C₁-C₆ alkoxy, C₁-C₂-alkylene dioxy, aryloxy-C₁-C₆ alkyl amine, C₁-C₁₂ perfluoro alkyl, S(O)_n-C₁-C₆ alkyl, S(O)_n-aryl where n is 0, 1
 10 or 2; OCOO-C₁-C₆ alkyl, OCOOaryl, OCONR⁶, COOH, COO-C₁-C₆ alkyl, COOaryl, CONR⁶R⁶, CONHOH, NR⁶R⁶, SO₂NR⁶R⁶, NR⁶SO₂aryl, NR⁶CONR⁶R⁶, NHSO₂CF₃, SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl, CONHSO₂aryl, SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl, CONHSO₂aryl, NH₂, OH, aryl, heteroaryl, C₃ to C₈ cycloalkyl; or saturated or
 15 unsaturated 5 to 10 membered mono or bicyclic heterocycle containing one heteroatom selected from O, S or NR⁷;
 wherein heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2
 20 groups selected independently from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;

R⁶ is H, C₁ to C₁₈ alkyl optionally substituted with OH; C₃ to C₆ alkenyl, C₃ to C₆ alkynyl, C₁ to C₆ perfluoro alkyl, S(O)_n alkyl or aryl where n is 0, 1, or 2; or COheteroaryl;

- 25 wherein heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;

- 30 and R⁷ is C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁-C₁₂ perfluoro alkyl, S(O)_n-alkyl, S(O)_n-aryl where n is 0, 1 or 2; COOalkyl, COOaryl, CONHR⁶, CONR⁶R⁶, CONHOH, SO₂NR⁶R⁶, SO₂CF₃, SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO₂alkyl, CONHSO₂aryl, aryl, heteroaryl; wherein C₁-C₆ alkyl is

WO 99/42436

PCT/US98/17633

- 177 -

straight or branched, heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;

5 alkyl of 1 to 18 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

alkenyl of 3 to 18 carbon atoms having from 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;

10 alkynyl of 3 to 18 carbon atoms having from 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;

arylalkyl of 7 to 16 carbon atoms, wherein aryl is optionally substituted with one or two groups selected independently from R⁵;

heteroarylalkyl wherein alkyl is from 1 to 6 carbon atoms and heteroaryl

15 contains 1 or 2 heteroatoms selected from O, S or N and is optionally substituted with one or two groups selected independently from R⁵;

biphenylalkyl of 13 to 18 carbon atoms, wherein biphenyl is optionally substituted with one or two groups selected independently from R⁵;

arylalkenyl of 8 to 16 carbon atoms, wherein aryl is optionally substituted with one or two groups selected independently from R⁵;

20 cycloalkylalkyl or bicycloalkylalkyl of 4 to 12 carbon atoms, wherein cycloalkyl or bicycloalkyl is optionally substituted with one or two groups selected independently from R⁵;

saturated or unsaturated mono or bicyclic heterocycle containing one heteroatom selected from O, S or N-C₁-C₆ alkyl, optionally substituted with one or

25 two groups selected independently from R⁵;

R⁸R⁹N-C₁-C₆-alkoxyaryl-C₁-C₆-alkyl where R⁸ and R⁹ are independently selected from C₁-C₆ alkyl or R⁸ and R⁹ together with the interposed nitrogen forms a 5-7 membered saturated heterocyclic ring optionally

30 containing an oxygen atom, wherein the aryl group is phenyl or naphthyl;

or a pharmaceutically acceptable salt thereof.

3. A compound according to claim 2 wherein

WO 99/42436

PCT/US98/17633

- 178 -

- R¹ is phenyl, naphthyl, alkyl of 1-18 carbon atoms or heteroaryl such as pyridyl, thienyl, imidazolyl or furanyl, optionally substituted with C₁-C₆ alkyl, C₁-C₆ alkoxy, C₆-C₁₀ aryloxy, heteroaryloxy, C₃-C₆ alkenyloxy, C₃-C₆ alkynyloxy, halogen; or S(O)_n-C₁-C₆alkyl C₁-C₆ alkoxyaryl or C₁-C₆ alkoxyheteroaryl;
- 5 A is -S-, -SO- or -SO₂-;
- R² and R³, taken with the carbon atom to which they are attached, form a 5 to 7 membered heterocyclic ring containing O, S or N-R⁷ optionally having one or two double bonds;
- R⁴ is hydrogen,
- 10 alkyl of 1 to 6 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;
- alkenyl of 3 to 18 carbon atoms having 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;
- alkynyl of 3 to 18 carbon atoms having 1 to 3 triple bonds, optionally
- 15 substituted with one or two groups selected independently from R⁵;
- phenyl or naphthyl optionally substituted with one or two groups selected independently from R⁵;
- C₃ to C₈ cycloalkyl or bicycloalkyl optionally substituted with one or two groups selected independently from R⁵;
- 20 R⁵ is H, C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁ to C₁₂ alkyl, C₂ to C₁₂ alkenyl, C₂-C₁₂ alkynyl, F, Cl, Br, I, CN, CHO, C₁-C₆ alkoxy, aryloxy, heteroaryloxy, C₃-C₆ alkenyloxy, C₃-C₆ alkynyloxy, C₁-C₆ alkylamino-C₁-C₆ alkoxy, C₁-C₂ alkylene dioxy, aryloxy-C₁-C₆ alkyl amine, C₁-C₁₂ perfluoro alkyl, S(O)_n-C₁-C₆ alkyl, S(O)_n-aryl where n is 0, 1 or 2; OCOO C₁-C₆ alkyl, OCOOaryl,
- 25 OCONR⁶, COOH, COO C₁-C₆ alkyl, COOaryl, CONR⁶R⁶, CONHOH, NR⁶R⁶, SO₂NR⁶R⁶, NR⁶SO₂aryl, -NR⁶CONR⁶R⁶, NHSO₂CF₃, SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl, CONHSO₂aryl, SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl, CONHSO₂aryl, NH₂, OH, aryl, heteroaryl, C₃ to C₈ cycloalkyl; saturated or unsaturated 5 to 10 membered
- 30 mono or bicyclic heterocycle containing one heteroatom selected from O, S or NR⁷, wherein C₁-C₆ alkyl is straight or branched, heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or NR⁷ and aryl is phenyl or

WO 99/42436

PCT/US98/17633

- 179 -

naphthyl, optionally substituted by 1 or 2 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;

- 5 R⁶ is H, C₁ to C₁₈ alkyl optionally substituted with OH; C₃ to C₆ alkenyl, C₃ to C₆ alkynyl, C₁ to C₆ perfluoro alkyl, S(O)_n alkyl or aryl where n is 0, 1 or 2; or COheteroaryl; wherein heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;
- 10 and R⁷ is C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁-C₁₂ perfluoro alkyl, S(O)_n-alkyl, S(O)_n-aryl where n is 0, 1 or 2; COOalkyl, COOaryl, CONHR⁶, CONR⁶R⁶, CONHOH, SO₂NR⁶R⁶, SO₂CF₃, SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO₂alkyl, CONHSO₂aryl, aryl, or heteroaryl; where aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected independently from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy; and heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or N-C₁-C₆ alkyl;
- 15 alkyl of 1 to 18 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;
- 20 alkenyl of 3 to 18 carbon atoms having from 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;
- alkynyl of 3 to 18 carbon atoms having from 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;
- 25 arylalkyl of 7 to 16 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;
- heteroarylalkyl wherein alkyl is from 1 to 6 carbon atoms and heteroaryl contains 1 or 2 heteroatoms selected from O, S or N and is optionally substituted with one or two groups selected independently from R⁵;
- 30 biphenylalkyl of 13 to 18 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;
- arylalkenyl of 8 to 16 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

WO 99/42436

PCT/US98/17633

- 180 -

- cycloalkylalkyl or bicycloalkylalkyl of 4 to 12 carbon atoms, optionally substituted with one or two groups selected independently from R⁵; saturated or unsaturated mono or bicyclic heterocycle containing one heteroatom selected from O, S or NR-C₁-C₆ alkyl, optionally substituted with one or two groups selected independently from R⁵;
- 5 R⁸R⁹N-C₁-C₆-alkoxyaryl-C₁-C₆-alkyl where R⁸ and R⁹ are independently selected from C₁-C₆ alkyl or R⁸ and R⁹ together with the interposed nitrogen forms a 5-7 membered saturated heterocyclic ring optionally containing an oxygen atom, wherein the aryl group is phenyl or
- 10 naphthyl;
- or a pharmaceutically acceptable salt thereof.
4. A compound according to claim 1 which is 1-benzyl-4-(4-methoxy-benzene-sulfonyl)-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutically acceptable
- 15 salt thereof.
5. A compound according to claim 1 which is 4-(4-methoxy-benzenesulfonyl)-1-(3-methoxy-benzyl)-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutically acceptable salt thereof.
- 20 6. A compound according to claim 1 which is 1-(3,4-dichlorobenzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxamide or a pharmaceutically acceptable salt thereof.
- 25 7. A compound according to claim 1 which is 4-(4-methoxy-benzenesulfonyl)-1-(4-methylbenzyl)-piperidine-4-carboxylic acid hydroxamide or a pharmaceutically acceptable salt thereof.
- 30 8. A compound according to claim 1 which is 4-(4-methoxy-benzene-sulfonyl)-1-napthalene-2-yl-methylpiperidine-4-carboxylic acid hydroxamide or a pharmaceutically acceptable salt thereof.
- 35 9. A compound according to claim 1 which is 1-biphenyl-4-ylmethyl-4-(4-methoxy-benzenesulfonyl)piperidine-4-carboxylic acid hydroxamide or a pharmaceutically acceptable salt thereof.

WO 99/42436

PCT/US98/17633

- 181 -

10. A compound according to claim 1 which is 4-(4-methoxy-benzene-sulfonyl)-1-(3-methyl-but-2-enyl)piperidine-4-carboxylic acid hydroxamide or a pharmaceutically acceptable salt thereof.
- 5 11. A compound according to claim 1 which is 1-(4-bromo-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutically acceptable salt thereof.
- 10 12. A compound according to claim 1 which is 4-(4-methoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid hydroxy amide or a pharmaceutically accepted salt thereof.
- 15 13. A compound according to claim 1 which is 1-benzyl-4-(4-benzyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical salt thereof.
- 20 14. A compound according to claim 1 which is 4-(4-butoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical salt thereof.
- 25 15. A compound according to claim 1 which is 4-(4-butoxy-benzenesulfonyl)-1-[3-(2-morpholinyl-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical salt thereof.
- 30 16. A compound according to claim 1 which is 1-methyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical salt thereof.
17. A compound according to claim 1 which is 1-ethyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical salt thereof.
18. A compound according to claim 1 which is 1-n-butyl-4-(4-butoxy-benzene-sulfonyl)-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical salt thereof.
- 35 19. A compound according to claim 1 which is 4-[4-(4-chloro-phenoxy)-benzenesulfonyl]1-methyl-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical salt thereof.

WO 99/42436

PCT/US98/17633

- 182 -

20. A compound according to claim 1 which is 4-[4-(4-chloro-phenoxy)-benzene-sulfonyl]1-ethyl-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical salt thereof.
- 5 21. A compound according to claim 1 which is 1-butyl-4-[4-(4-chloro-phenoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical salt thereof.
22. A compound according to claim 1 which is 1-benzyl-4-[4-(4-chloro-phenoxy)-
10 benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical salt thereof.
23. A compound according to claim 1 which is 1-benzyl-4-[4-(3-methyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical salt
15 thereof.
24. A compound according to claim 1 which is 1-butyl-4-[4-(3-methyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical salt thereof.
- 20 25. A compound according to claim 1 which is 1-benzyl-4-[4-(2-ethyl-butoxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical salt thereof.
- 25 26. A compound according to claim 1 which is 4-(4-butoxy-benzenesulfonyl)-1-(3-methoxy-benzyl)-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical salt thereof.
27. A compound according to claim 1 which is 4-(4-methoxy-benzenesulfonyl)-1-(4-
30 thiophen-2-yl-benzyl)-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical salt thereof.
28. A compound according to claim 1 which is 4-(4-methoxy-benzenesulfonyl)-1-(4-pyridin-2-yl-benzyl)-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical
35 salt thereof.
29. A compound according to claim 1 which is 1-(3,4-dichlorobenzyl)-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide or a pharmaceutical salt thereof.

40

WO 99/42436

PCT/US98/17633

- 183 -

30. A compound according to claim 1 which is [4-(4-chloro-benzyloxy)-benzenesulfonyl]-1-methylpiperidine-4-carboxylic acid hydroxamide or a pharmaceutical salt thereof.
- 5 31. A compound according to claim 1 which is 4-(4-butoxy-benzenesulfonyl)-1-(3-phenoxy-benzyl)-piperidine-4-carboxylic acid hydroxamide or a pharmaceutical salt thereof.
- 10 32. A compound according to claim 1 which is [4-(4-chloro-benzyloxy)-benzenesulfonyl]-1-(4-methylbenzyl)-piperidine-4-carboxylic acid hydroxamide or a pharmaceutical salt thereof.
- 15 33. A compound according to claim 1 which is 4-(4-butoxy-benzenesulfonyl)-1-(4-methylbenzyl)-piperidine-4-carboxylic acid hydroxamide or a pharmaceutical salt thereof.
- 20 34. A compound according to claim 1 which is 4-(4-butoxy-benzenesulfonyl)-1-(4-cyano-benzyl)-piperidine-4-carboxylic acid hydroxamide or a pharmaceutical salt thereof.
- 25 35. A compound according to claim 1 which is 4-(4-Butoxy-benzenesulfonyl)-1-pyridin-4-ylmethyl-piperidine-4-carboxylic acid hydroxamide or a pharmaceutical salt thereof.
- 30 36. A compound according to claim 1 or a pharmaceutically acceptable salt thereof which is selected from the group of compounds consisting of:
4-(4-methoxy-benzenesulfonyl)-1-(3-phenyl-propyl)-piperidine-4-carboxylic acid hydroxyamide,
1-tert-butyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide,
1-butyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide,
1-cyclooctyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide,
1-ethyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide,
35 1-isopropyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide,
1-methyl-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide,
1-benzyl-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide,


WO 99/42436

PCT/US98/17633

- 184 -

- 1-(4-fluoro-benzyl)-4-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid
hydroxyamide,
1-(4-fluoro-benzyl)-4-(4-butoxy-benzenesulfonyl)-piperidine-4-carboxylic acid
hydroxyamide,
5 4-(4-methoxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid
hydroxyamide,
4-(4-methoxy-benzenesulfonyl)-1-[2-(4-methoxyphenyl)-ethyl]-piperidine-4-carboxylic
acid hydroxyamide,
4-(4-methoxy-benzenesulfonyl)-1-(2-phenyl-ethyl)-piperidine-4-carboxylic acid
10 hydroxyamide,
4-(4-n-butoxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid
hydroxyamide,
4-(4-methoxy-benzenesulfonyl)-1-(3-phenoxy-propyl)-piperidine-4-carboxylic acid
hydroxyamide,
15 4-(4-n-butoxy-benzenesulfonyl)-1-(3-phenoxy-propyl)-piperidine-4-carboxylic acid
hydroxyamide,
4-(4-methoxy-benzenesulfonyl)-1-(2-phenoxy-ethyl)-piperidine-4-carboxylic acid
hydroxyamide,
4-(4-n-butoxy-benzenesulfonyl)-1-(2-phenoxy-ethyl)-piperidine-4-carboxylic acid
20 hydroxyamide, and
4-(4-methoxy-benzenesulfonyl)-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-
carboxylic acid hydroxy amide.

37. A method of inhibiting pathological changes mediated by matrix metalloproteinases
25 in mammals which comprises administration to a mammal in need thereof a
therapeutically effective amount of a matrix metalloproteinase inhibiting compound of
the formula



- 30 wherein:

R¹ is alkyl of 1 to 18 carbon atoms, optionally substituted with one or two groups
selected independently from R⁵;
alkenyl of 3 to 18 carbon atoms having 1 to 3 double bonds, optionally
substituted with one or two groups selected independently from R⁵;

WO 99/42436

PCT/US98/17633

- 185 -

- alkynyl of 3 to 18 carbon atoms having 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;
aryl of 6 to 10 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;
5 cycloalkyl of 3 to 8 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;
saturated or unsaturated 5 to 10 membered mono or bicyclic heterocycle containing one heteroatom selected from O, S or NR⁷, optionally substituted with one or two groups selected independently from R⁵;
10 or heteroaryl-(CH₂)₀₋₆ wherein the heteroaryl group is 5 to 6 membered with one or two heteroatoms selected independently from O, S, and N and may be optionally substituted with one or two groups selected independently from R⁵;
A is -S-, -SO- or SO₂-;
15 R² and R³, taken with the carbon atom to which they are attached, form a 5 to 7 membered heterocyclic ring containing O, S or N-R⁷ optionally having one or two double bonds;
R⁴ is hydrogen,
alkyl of 1 to 6 carbon atoms, optionally substituted with one or two groups
20 selected independently from R⁵;
alkenyl of 3 to 18 carbon atoms having 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;
alkynyl of 3 to 18 carbon atoms having 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;
25 phenyl or naphthyl optionally substituted with one or two groups selected independently from R⁵;
C₃ to C₈ cycloalkyl or bicycloalkyl optionally substituted with one or two groups selected independently from R⁵;
saturated or unsaturated 5 to 10 membered mono or bicyclic heterocycle
30 containing one heteroatom selected from O, S or NR⁷, optionally substituted with one or two groups selected independently from R⁵;
R⁵ is H, C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁ to C₁₂ alkyl, C₂ to C₁₂ alkenyl, C₂-C₁₂ alkynyl, F, Cl, Br, I, CN, CHO, C₁-C₆ alkoxy, aryloxy, heteroaryloxy, C₃-C₆

WO 99/42436

PCT/US98/17633

- 186 -

- alkenyloxy, C₃-C₆ alkynyloxy, C₁-C₆ alkoxyaryl, C₁-C₆ alkoxyheteroaryl, C₁-C₆ alkylamino-C₁-C₆ alkoxy, C₁-C₂ alkylene dioxy, aryloxy-C₁-C₆ alkyl amine, C₁-C₁₂ perfluoro alkyl, S(O)_n-C₁-C₆ alkyl, S(O)_n-aryl where n is 0, 1 or 2; OCOO C₁-C₆ alkyl, OCOOaryl, OCONR⁶, COOH, COO C₁-C₆ alkyl, 5 COOaryl, CONR⁶R⁶, CONHOH, NR⁶R⁶, SO₂NR⁶R⁶, NR⁶SO₂aryl, - NR⁶CONR⁶R⁶, NHSO₂CF₃, SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl, CONHSO₂aryl, SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl, CONHSO₂aryl, NH₂, OH, aryl, heteroaryl, C₃ to C₈ cycloalkyl; or saturated or unsaturated 5 to 10 membered mono or bicyclic heterocycle containing one 10 heteroatom selected from O, S or NR⁷, wherein C₁-C₆ alkyl is straight or branched, heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl, 15 C₁-C₆ alkoxy, or hydroxy;
- R⁶ is H, C₁ to C₁₈ alkyl optionally substituted with OH; C₃ to C₆ alkenyl, C₃ to C₆ alkynyl, C₁ to C₆ perfluoro alkyl, S(O)_n-C₁-C₆ alkyl S(O)_n aryl where n is 0, 1 or 2; or COheteroaryl, wherein heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected 20 independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;
- and R⁷ is C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁-C₁₂ perfluoro alkyl, S(O)_n-C₁-C₆-alkyl, S(O)_n-aryl where n is 0, 1 or 2; COO-C₁-C₆-alkyl, COOaryl, CONHR⁶, 25 CONR⁶R⁶, CONHOH, SO₂NR⁶R⁶, SO₂CF₃, SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO-C₁-C₆-alkyl, CONHSO₂aryl, aryl, or heteroaryl, where aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected independently from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy; and heteroaryl is a 5-10 30 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or N-C₁-C₆ alkyl;
- alkyl of 1 to 18 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

WO 99/42436

PCT/US98/17633

- 187 -

- alkenyl of 3 to 18 carbon atoms having from 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;
 alkynyl of 3 to 18 carbon atoms having from 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;
 5 arylalkyl of 7 to 16 carbon atoms, wherein aryl is optionally substituted with one or two groups selected independently from R⁵;
 heteroarylalkyl wherein alkyl is from 1 to 6 carbon atoms and heteroaryl contains 1 or 2 heteroatoms selected from O, S or N and is optionally substituted with one or two groups selected independently from R⁵;
 10 biphenylalkyl of 13 to 18 carbon atoms, wherein biphenyl is optionally substituted with one or two groups selected independently from R⁵;
 arylalkenyl of 8 to 16 carbon atoms, wherein aryl is optionally substituted with one or two groups selected independently from R⁵;
 cycloalkylalkyl or bicycloalkylalkyl of 4 to 12 carbon atoms, wherein the
 15 cycloalkyl or bicycloalkyl group is optionally substituted with one or two groups selected independently from R⁵;
 saturated or unsaturated mono or bicyclic heterocycle containing one heteroatom selected from O, S or N-C₁-C₆ alkyl, optionally substituted with one or two groups selected independently from R⁵; or
 20 R⁸R⁹N-C₁-C₆-alkoxyaryl-C₁-C₆-alkyl where R⁸ and R⁹ are independently selected from C₁-C₆ alkyl or R⁸ and R⁹ together with the interposed nitrogen forms a 5-7 membered saturated heterocyclic ring optionally containing an oxygen atom, wherein the aryl group is phenyl or naphthyl;
 25 or a pharmaceutically acceptable salt thereof.

38. A method according to claim 37 wherein the condition treated is atherosclerosis, atherosclerotic plaque formation, reduction of coronary thrombosis from atherosclerotic plaque rupture, restenosis, MMP-mediated osteopenias, inflammatory diseases of the
 30 central nervous system, skin aging, angiogenesis, tumor metastasis, tumor growth, osteoarthritis, rheumatoid arthritis, septic arthritis, corneal ulceration, abnormal wound healing, bone disease, proteinuria, aneurysmal aortic disease, degenerative cartilage loss following traumatic joint injury, demyelinating diseases of the nervous system, cirrhosis of the liver, glomerular disease of the kidney, premature rupture of fetal
 35 membranes, inflammatory bowel disease, or periodontal disease.

WO 99/42436


PCT/US98/17633

- 188 -

39. A method according to claim 37 wherein the condition treated is age related macular degeneration, diabetic retinopathy, proliferative vitreoretinopathy, retinopathy of prematurity, ocular inflammation, keratoconus, Sjogren's syndrome, myopia, ocular tumors, ocular angiogenesis/neovascularization and corneal graft rejection.

5

40. A method of inhibiting pathological changes mediated by TNF- α converting enzyme (TACE) in mammals which comprises administration to a mammal in need thereof a therapeutically effective amount of a TACE inhibiting compound of the formula



10

wherein:

R¹ is alkyl of 1 to 18 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

15

alkenyl of 3 to 18 carbon atoms having 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;

alkynyl of 3 to 18 carbon atoms having 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;

aryl of 6 to 10 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

20

cycloalkyl of 3 to 8 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

saturated or unsaturated 5 to 10 membered mono or bicyclic heterocycle containing one heteroatom selected from O, S or NR⁷, optionally

25

substituted with one or two groups selected independently from R⁵;
or heteroaryl-(CH₂)₀₋₆- wherein the heteroaryl group is 5 to 6 membered with one or two heteroatoms selected independently from O, S, and N and may be optionally substituted with one or two groups selected independently from R⁵;

30

A is -S-, -SO- or SO₂-;

R² and R³, taken with the carbon atom to which they are attached, form a 5 to 7 membered heterocyclic ring containing O, S or N-R⁷ optionally having one or two double bonds;

WO 99/42436

PCT/US98/17633

- 189 -

R⁴ is hydrogen,

alkyl of 1 to 6 carbon atoms, optionally substituted with one or two groups
selected independently from R⁵;

alkenyl of 3 to 18 carbon atoms having 1 to 3 double bonds, optionally

5 substituted with one or two groups selected independently from R⁵;

alkynyl of 3 to 18 carbon atoms having 1 to 3 triple bonds, optionally

substituted with one or two groups selected independently from R⁵;

phenyl or naphthyl optionally substituted with one or two groups selected
independently from R⁵;

10 C₃ to C₈ cycloalkyl or bicycloalkyl optionally substituted with one or two
groups selected independently from R⁵;

saturated or unsaturated 5 to 10 membered mono or bicyclic heterocycle

containing one heteroatom selected from O, S or NR⁷, optionally

substituted with one or two groups selected independently from R⁵;

15 R⁵ is H, C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁ to C₁₂ alkyl, C₂ to C₁₂ alkenyl, C₂-C₁₂
alkynyl, F, Cl, Br, I, CN, CHO, C₁-C₆ alkoxy, aryloxy, heteroaryloxy, C₃-C₆
alkenyloxy, C₃-C₆ alkynyloxy, C₁-C₆ alkoxyaryl, C₁-C₆ alkoxyheteroaryl,
C₁-C₆ alkylamino-C₁-C₆ alkoxy, C₁-C₂ alkylene dioxy, aryloxy-C₁-C₆ alkyl
amine, C₁-C₁₂ perfluoro alkyl, S(O)_n-C₁-C₆ alkyl, S(O)_n-aryl where n is 0, 1
20 or 2; OCOO C₁-C₆ alkyl, OCOOaryl, OCONR⁶, COOH, COO C₁-C₆ alkyl,
COOaryl, CONR⁶R⁶, CONHOH, NR⁶R⁶, SO₂NR⁶R⁶, NR⁶SO₂aryl, -
NR⁶CONR⁶R⁶, NHSO₂CF₃, SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO₂-
C₁-C₆ alkyl, CONHSO₂aryl, SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl,
CONHSO₂aryl, NH₂, OH, aryl, heteroaryl, C₃ to C₈ cycloalkyl; or saturated or
25 unsaturated 5 to 10 membered mono or bicyclic heterocycle containing one
heteroatom selected from O, S or NR⁷, wherein C₁-C₆ alkyl is straight or
branched, heteroaryl is a 5-10 membered mono or bicyclic heteroaryl
group having 1 to 3 heteroatoms selected independently from O, S or
NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2
30 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl,
C₁-C₆ alkoxy, or hydroxy;

R⁶ is H, C₁ to C₁₈ alkyl optionally substituted with OH; C₃ to C₆ alkenyl, C₃ to C₆
alkynyl, C₁ to C₆ perfluoro alkyl, S(O)_n-C₁-C₆ alkyl S(O)_n aryl where n is 0,

WO 99/42436

PCT/US98/17633

- 190 -

- 1 or 2; or COheteroaryl, wherein heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;
- 5 and R⁷ is C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁-C₁₂ perfluoro alkyl, S(O)_n-C₁-C₆-alkyl, S(O)_n-aryl where n is 0, 1 or 2; COO-C₁-C₆-alkyl, COOaryl, CONHR⁶, CONR⁶R⁶, CONHOH, SO₂NR⁶R⁶, SO₂CF₃, SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO-C₁-C₆-alkyl, CONHSO₂aryl, aryl, or heteroaryl,
- 10 where aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected independently from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy; and heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or N-C₁-C₆ alkyl;
- 15 alkyl of 1 to 18 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;
- alkenyl of 3 to 18 carbon atoms having from 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;
- alkynyl of 3 to 18 carbon atoms having from 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;
- 20 arylalkyl of 7 to 16 carbon atoms, wherein aryl is optionally substituted with one or two groups selected independently from R⁵;
- heteroarylalkyl wherein alkyl is from 1 to 6 carbon atoms and heteroaryl contains 1 or 2 heteroatoms selected from O, S or N and is optionally substituted with one or two groups selected independently from R⁵;
- 25 biphenylalkyl of 13 to 18 carbon atoms, wherein biphenyl is optionally substituted with one or two groups selected independently from R⁵;
- arylalkenyl of 8 to 16 carbon atoms, wherein aryl is optionally substituted with one or two groups selected independently from R⁵;
- 30 cycloalkylalkyl or bicycloalkylalkyl of 4 to 12 carbon atoms, wherein the cycloalkyl or bicycloalkyl group is optionally substituted with one or two groups selected independently from R⁵;

WO 99/42436

PCT/US98/17633

- 191 -

saturated or unsaturated mono or bicyclic heterocycle containing one heteroatom selected from O, S or N-C₁-C₆ alkyl, optionally substituted with one or two groups selected independently from R⁵; or

5 R⁸R⁹N-C₁-C₆-alkoxyaryl-C₁-C₆-alkyl where R⁸ and R⁹ are independently selected from C₁-C₆ alkyl or R⁸ and R⁹ together with the interposed nitrogen forms a 5-7 membered saturated heterocyclic ring optionally containing an oxygen atom, wherein the aryl group is phenyl or naphthyl;


or a pharmaceutically acceptable salt thereof.

10

41. The method according to claim 40 wherein the condition treated is rheumatoid arthritis, graft rejection, cachexia, anorexia, inflammation, fever, insulin resistance, septic shock, congestive heart failure, inflammatory disease of the central nervous system, inflammatory bowel disease, or HIV infection.

15

42. A pharmaceutical composition comprising a pharmaceutical carrier and a therapeutically effective amount of a matrix metalloproteinase or TACE inhibiting compound according to the formula



20

wherein:

R¹ is alkyl of 1 to 18 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

25

alkenyl of 3 to 18 carbon atoms having 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;

alkynyl of 3 to 18 carbon atoms having 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;

aryl of 6 to 10 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

30

cycloalkyl of 3 to 8 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

saturated or unsaturated 5 to 10 membered mono or bicyclic heterocycle containing one heteroatom selected from O, S or NR⁷, optionally substituted with one or two groups selected independently from R⁵;

WO 99/42436

PCT/US98/17633

- 192 -

or heteroaryl-(CH₂)₀₋₆- wherein the heteroaryl group is 5 to 6 membered with one or two heteroatoms selected independently from O, S, and N and may be optionally substituted with one or two groups selected independently from R⁵;

5 A is -S-, -SO- or SO₂-;

R² and R³, taken with the carbon atom to which they are attached, form a 5 to 7 membered heterocyclic ring containing O, S or N-R⁷ optionally having one or two double bonds;

R⁴ is hydrogen,

10 alkyl of 1 to 6 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;

alkenyl of 3 to 18 carbon atoms having 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;

15 alkynyl of 3 to 18 carbon atoms having 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;

phenyl or naphthyl optionally substituted with one or two groups selected independently from R⁵;

C₃ to C₈ cycloalkyl or bicycloalkyl optionally substituted with one or two groups selected independently from R⁵;

20 saturated or unsaturated 5 to 10 membered mono or bicyclic heterocycle containing one heteroatom selected from O, S or NR⁷, optionally substituted with one or two groups selected independently from R⁵;

R⁵ is H, C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁ to C₁₂ alkyl, C₂ to C₁₂ alkenyl, C₂-C₁₂ alkynyl, F, Cl, Br, I, CN, CHO, C₁-C₆ alkoxy, aryloxy, heteroaryloxy, C₃-C₆ alkenyloxy, C₃-C₆ alkynyloxy, C₁-C₆ alkoxyaryl, C₁-C₆ alkoxyheteroaryl, C₁-C₆ alkylamino-C₁-C₆ alkoxy, C₁-C₂ alkylene dioxy, aryloxy-C₁-C₆ alkyl amine, C₁-C₁₂ perfluoro alkyl, S(O)_n-C₁-C₆ alkyl, S(O)_n-aryl where n is 0, 1 or 2; OCOO C₁-C₆ alkyl, OCOOaryl, OCONR⁶, COOH, COO C₁-C₆ alkyl, COOaryl, CONR⁶R⁶, CONHOH, NR⁶R⁶, SO₂NR⁶R⁶, NR⁶SO₂aryl, 25 -NR⁶CONR⁶R⁶, NHSO₂CF₃, SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl, CONHSO₂aryl, SO₂NHCOaryl, CONHSO₂-C₁-C₆ alkyl, CONHSO₂aryl, NH₂, OH, aryl, heteroaryl, C₃ to C₈ cycloalkyl; or saturated or unsaturated 5 to 10 membered mono or bicyclic heterocycle containing one 30

WO 99/42436

PCT/US98/17633

- 193 -

- heteroatom selected from O, S or NR⁷, wherein C₁-C₆ alkyl is straight or branched, heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;
- R⁶ is H, C₁ to C₁₈ alkyl optionally substituted with OH; C₃ to C₆ alkenyl, C₃ to C₆ alkynyl, C₁ to C₆ perfluoro alkyl, S(O)_n-C₁-C₆ alkyl S(O)_n aryl where n is 0, 1 or 2; or COheteroaryl, wherein heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or NR⁷ and aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy;
- and R⁷ is C₇-C₁₁ aroyl, C₂-C₆ alkanoyl, C₁-C₁₂ perfluoro alkyl, S(O)_n-C₁-C₆-alkyl, S(O)_n-aryl where n is 0, 1 or 2; COO-C₁-C₆-alkyl, COOaryl, CONHR⁶, CONR⁶R⁶, CONHOH, SO₂NR⁶R⁶, SO₂CF₃, SO₂NHheteroaryl, SO₂NHCOaryl, CONHSO-C₁-C₆-alkyl, CONHSO₂aryl, aryl, or heteroaryl, where aryl is phenyl or naphthyl, optionally substituted by 1 or 2 groups selected independently from halogen, cyano, amino, nitro, C₁-C₆ alkyl, C₁-C₆ alkoxy, or hydroxy; and heteroaryl is a 5-10 membered mono or bicyclic heteroaryl group having 1 to 3 heteroatoms selected independently from O, S or N-C₁-C₆ alkyl;
- alkyl of 1 to 18 carbon atoms, optionally substituted with one or two groups selected independently from R⁵;
- alkenyl of 3 to 18 carbon atoms having from 1 to 3 double bonds, optionally substituted with one or two groups selected independently from R⁵;
- alkynyl of 3 to 18 carbon atoms having from 1 to 3 triple bonds, optionally substituted with one or two groups selected independently from R⁵;
- arylalkyl of 7 to 16 carbon atoms, wherein aryl is optionally substituted with one or two groups selected independently from R⁵;
- heteroarylalkyl wherein alkyl is from 1 to 6 carbon atoms and heteroaryl contains 1 or 2 heteroatoms selected from O, S or N and is optionally substituted with one or two groups selected independently from R⁵;

WO 99/42436

PCT/US98/17633

- 194 -

- biphenylalkyl of 13 to 18 carbon atoms, wherein biphenyl is optionally substituted with one or two groups selected independently from R⁵;
- arylalkenyl of 8 to 16 carbon atoms, wherein aryl is optionally substituted with one or two groups selected independently from R⁵;
- 5 cycloalkylalkyl or bicycloalkylalkyl of 4 to 12 carbon atoms, wherein the cycloalkyl or bicycloalkyl group is optionally substituted with one or two groups selected independently from R⁵;
- saturated or unsaturated mono or bicyclic heterocycle containing one heteroatom selected from O, S or N-C₁-C₆ alkyl, optionally substituted with one or
- 10 two groups selected independently from R⁵; or
- R⁸R⁹N-C₁-C₆-alkoxyaryl-C₁-C₆-alkyl where R⁸ and R⁹ are independently selected from C₁-C₆ alkyl or R⁸ and R⁹ together with the interposed nitrogen forms a 5-7 membered saturated heterocyclic ring optionally containing an oxygen atom, wherein the aryl group is phenyl or
- 15 naphthyl;
- or a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/17633

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07C239/14 C07D211/20 C07D241/04 A61K31/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07C C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 606 046 A (CIBA-GEIGY AG) 13 July 1994 cited in the application see the whole document ---	1, 37, 40, 42
A	WO 96 00214 A (CIBA-GEIGY AG) 4 January 1996 see the whole document ---	1, 37, 40, 42
A	WO 95 35275 A (BRITISH BIOTECH PHARMACEUTICALS LIMITED) 28 December 1995 see the whole document ---	1, 37, 40, 42
A	WO 95 19961 A (BRITISH BIOTECH PHARMACEUTICALS) 27 July 1995 see the whole document ---	1, 37, 40, 42
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

21 December 1998

Date of mailing of the international search report

04/01/1999

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kyriakakou, G

INTERNATIONAL SEARCH REPORT

Inter. natl Application No
PCT/US 98/17633

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 90 05719 A (BRITISH BIOTECHNOLOGY LIMITED) 31 May 1990 see the whole document -----	1,37,40, 42

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/US 98/17633

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 606046	A	13-07-1994	US 5455258 A	03-10-1995
			AT 159012 T	15-10-1997
			AU 684255 B	11-12-1997
			AU 5265593 A	04-05-1995
			CA 2112779 A	07-07-1994
			DE 69314456 D	13-11-1997
			DE 69314456 T	26-02-1998
			DK 606046 T	04-05-1998
			ES 2107648 T	01-12-1997
			FI 940012 A	07-07-1994
			GR 3025611 T	31-03-1998
			HU 70536 A	30-10-1995
			JP 6256293 A	13-09-1994
			MX 9400276 A	29-07-1994
			NO 940038 A, B,	07-07-1994
			NZ 250517 A	26-10-1995
			SG 42933 A	17-10-1997
			US 5506242 A	09-04-1996
			US 5552419 A	03-09-1996
			US 5646167 A	08-07-1997
			US 5672615 A	30-09-1997
			ZA 9400048 A	11-08-1994
WO 9600214	A	04-01-1996	US 5506242 A	09-04-1996
			AU 692553 B	11-06-1998
			AU 2536995 A	19-01-1996
			CA 2192092 A	04-01-1996
			EP 0766672 A	09-04-1997
			FI 965156 A	20-12-1996
			HU 76548 A	29-09-1997
			NO 965568 A	17-02-1997
			US 5552419 A	03-09-1996
			US 5646167 A	08-07-1997
			US 5672615 A	30-09-1997
			US 5817822 A	06-10-1998
			ZA 9505206 A	27-12-1995
WO 9535275	A	28-12-1995	AU 695507 B	13-08-1998
			AU 2746595 A	15-01-1996
			AU 690703 B	30-04-1998
			AU 2746695 A	15-01-1996
			CA 2193691 A	28-12-1995
			CA 2193692 A	28-12-1995
			CN 1151157 A	04-06-1997
			EP 0766664 A	09-04-1997
			EP 0766665 A	09-04-1997
			FI 965153 A	20-12-1996
			WO 9535276 A	28-12-1995
			GB 2303850 A, B	05-03-1997
			GB 2303629 A, B	26-02-1997
			JP 10501806 T	17-02-1998
			JP 10507158 T	14-07-1998
			NO 965515 A	20-02-1997
			NZ 288299 A	26-06-1998
WO 9519961	A	27-07-1995	AU 678884 B	12-06-1997
			AU 1460395 A	08-08-1995
			AU 1654097 A	22-05-1997

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/17633

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9519961 A		CA 2181709 A	27-07-1995
		EP 0740655 A	06-11-1996
		FI 962905 A	19-07-1996
		GB 2300188 A,B	30-10-1996
		GB 2315750 A,B	11-02-1998
		HU 74511 A	28-01-1997
		JP 9508362 T	26-08-1997
		NO 963031 A	20-09-1996
		NZ 278632 A	27-04-1998
WO 9005719 A	31-05-1990	AU 644064 B	02-12-1993
		AU 4800390 A	12-06-1990
		CA 2003718 A,C	23-05-1990
		DE 68914687 D	19-05-1994
		DE 68914687 T	08-09-1994
		DK 96691 A	23-07-1991
		EP 0446267 A	18-09-1991
		ES 2055409 T	16-08-1994
		FI 100470 B	15-12-1997
		HU 9500245 A	28-09-1995
		JP 2565599 B	18-12-1996
		JP 4502008 T	09-04-1992
		NO 177701 B	31-07-1995
		US 5310763 A	10-05-1994
		US 5240958 A	31-08-1993